

FIAN/TD-30/00
ITEP/TH-64/00
hep-th/0011197

Integrable Many-Body Systems and Gauge Theories

A.Gorsky *

ITEP, Moscow 117259, Russia

and

A.Mironov [†]

Theory Department, Lebedev Physics Institute, Moscow 117924, Russia
and ITEP, Moscow 117259, Russia

Abstract

The review studies connections between integrable many-body systems and gauge theories. It is shown how the degrees of freedom in integrable systems are related with topological degrees of freedom in gauge theories. The relations between families of integrable systems and $\mathcal{N} = 2$ supersymmetric gauge theories are described. It is explained that the degrees of freedom in the many-body systems can be identified with collective coordinates of D-branes, solitons in string theory.

*E-mail: gorsky@heron.itep.ru

[†]E-mail: mironov@lpi.ru, mironov@itep.ru

Contents

1	Introduction	3
1.1	Moduli and nonperturbative degrees of freedom	3
1.2	Seiberg-Witten solution	6
1.3	Seiberg-Witten theory and theory of prepotential	8
1.4	Duality, branes and gauge theories	10
1.5	Seiberg-Witten solution and branes	11
2	Many-body systems and gauge theories in two and three dimensions	13
2.1	Calogero many-body system and Yang-Mills theory in two dimensions	13
2.2	Gauged G/G σ -model and Ruijsenaars systems	16
2.3	Elliptic Calogero model and gauge theories	19
3	Particle systems and supersymmetric gauge theories	21
3.1	Seiberg-Witten theory and integrable systems	22
3.2	$4d$ pure gauge theory: Toda chain	25
3.3	Adjoint hypermultiplet added: Calogero system	29
3.4	Fundamental matter added (SQCD): XXX spin chain	31
3.5	$5d$ SQCD: twisted XXZ chain	36
3.6	$5d$ pure gauge theory: relativistic Toda chain	39
3.7	$5d$ adjoint matter: Ruijsenaars-Schneider system	41
3.8	Fundamental matter in $6d$: XYZ chain	46
4	Integrability and $\mathcal{N} = 1$ supersymmetric YM theories	49
4.1	Superpotentials in $\mathcal{N} = 1$ theories from $\mathcal{N} = 2$ spectral curves	49
4.2	Superpotentials from nonperturbative configurations	52
4.3	$\mathcal{N} = 1$ theories and integrability	54
5	Perturbative prepotentials	55
5.1	Perturbative prepotentials in gauge theories	55
5.2	Perturbative prepotentials in integrable systems	57
5.3	$4d$ theory: a simple example of calculation	58
5.4	From $4d$ to higher dimensions	59
5.5	Perturbative prepotentials: fundamental matter	61
5.6	$SU(2)$ adjoint matter in $5d$	63
5.7	Perturbative prepotentials: adjoint matter	66
6	Theory of prepotential	68

6.1	From Seiberg-Witten theory to Whitham hierarchies: RG dynamics	68
6.2	Whitham hierarchy for the Toda chain	71
6.3	Solution to Whitham hierarchy and prepotential	74
6.4	Whitham hierarchy for $SU(2)$: an example	77
6.5	WDVV equations	80
6.6	WDVV equations in Seiberg-Witten theory	83
6.7	Covariance of the WDVV equations	87
7	Dualities in the many-body systems and gauge theories	90
7.1	D-branes, gauge theory and separation of variables	90
7.2	S-duality	91
7.3	“Mirror” symmetry in dynamical systems	92
7.4	Duality as the canonical transformation: two-body system	94
7.5	Two-body systems dual to Calogero and Ruijsenaars	96
7.6	Double-elliptic two-body system	99
7.7	Duality, τ -functions and Hamiltonian reduction	101
7.8	Dual Hamiltonians for many-body systems	104
7.9	Dual Hamiltonians perturbatively	105
7.10	Dell systems: 6d adjoint matter	108
7.11	Quantum duality	110
8	Branes as low energy degrees of freedom in supersymmetric gauge theories	112
8.1	Field theory on D-brane worldvolume	112
8.2	Branes and degrees of freedom in many-body systems	114
8.3	Branes in five dimensional theory	116
8.4	Branes and equations of motion in many-body systems	118
8.5	Branes and two Lax representations	120
8.6	Whitham dynamics via branes	121
9	Conclusion	122

1 Introduction

This review is devoted to the issue of hidden integrability in quantum field theory, i.e. integrability in the systems that naively have not enough many conserved integrals of motion. This situation differs considerably from the picture familiar for the experts in the theory of integrable systems. In the typical integrable system, one deals with a well-defined (field theory) Lagrangian, with the fields given on a base space-time manifold. The fields at some fixed moment of time, i.e. given on a space slice of the base manifold, provide the dynamical degrees of freedom, while integrable flows are identified with the time direction in the base manifold.

On the contrary, in the systems we are interested in, the phase space for the integrable dynamics is provided by a moduli space essential in the problem. The collective coordinates or moduli arise in the course of consideration of various issues and have essentially different origins. In particular, they can describe effective degrees of freedom important in one or another regime of the field theory not completely integrable by itself. The proper regime can be determined by the kinematics of the process or by the choice of specific topologically nontrivial data. In the cases when there are several collective degrees of freedom, a hidden integrability in the theory emerges as a rule. Its origin lies in symmetry properties of the moduli space.

As a particular example, it appears that a hidden integrability governs the low-energy effective actions of $\mathcal{N} = 2$ supersymmetric gauge theories, which in four dimensions were described by Seiberg and Witten [1, 2].

The role of this integrability is twofold; it allows one to uncover the hidden symmetries and to define proper degrees of freedom. In what follows, we shall show how the low-energy sector of supersymmetric gauge theories is described via finite dimensional integrable systems.

Restricted frames of the review does not allow us to provide the reader with all necessary information from the theory of integrable systems, supersymmetric gauge theories and nonperturbative string theory which might be relevant to the issue. This kind of problem arises since the integrability appears at the intersection of different branches of theoretical physics which use different tools and approaches. Therefore, we just recommend several (rather accidental) review papers, where the issues used in what follows as well as the proper list of references can be found. Other references can be found in the proper parts of the present review.

The finite gap integration is perfectly presented in [3], Whitham hierarchies are discussed in [4, 5], while background of the Seiberg-Witten approach is described in [6]. Nonperturbative string theory is reviewed in [7]. Necessary background from algebraic geometry used in the brane approach is described in [8], while a generic point of view on integrability in (nonperturbative) field/string theory can be found in [9]. An intensive discussion of integrability in Seiberg-Witten theory and related structures (Whitham hierarchies and WDVV equations) can be found in the book [10] that contains recent comprehensive reviews. Note also that the very detailed review [11] may serve a good complimentary text to the present paper.

1.1 Moduli and nonperturbative degrees of freedom

Description of the strong coupling regime remains one of the most important problems of quantum field theory. Since the perturbation theory does not work in this region, one can mostly reply on constructing some effective degrees of freedom which would allow one to develop a new perturbation theory with a small parameter distinct from the coupling constant of the initial theory. A nice example

of the existence of different descriptions at different energy scales is quantum chromodynamics (QCD). At high energies, quarks and gluons giving the proper degrees of freedom are described by the Yang-Mills Lagrangian coupled to fermions in the fundamental representation, while at low energies it is a chiral (σ -model) Lagrangian that describes the low-energy effective action, with the proper degrees of freedom given by the colorless mesons and baryons. Note that the parameters of the chiral Lagrangian can not be calculated from the first principles (at the moment) and are fixed phenomenologically.

The general structure of low-energy actions is fixed by the symmetries. It is expected that the effective actions enjoy a complete set of the Ward identities of the initial field theory. Some of the Ward identities may be anomalous. However, since the anomalies are related to the level crossing phenomena, not renormalized and, therefore, can be equally well described in the UV and IR terms [12], the anomalous Ward identities could still impose restrictions on the structure of effective theories. For instance, the chiral anomalous Ward identities fix the Lagrangian of mesons, while the conformal Ward identities yield the effective Lagrangian for the dilaton in ordinary as well as in $\mathcal{N} = 1$ supersymmetric QCD [13].

Another phenomenon that has much to do with the issues discussed is universality. Effective actions typically enjoy this feature which implies that several theories different in the UV regime can flow to the same theory in the infrared region. The reason for such an universality to occur is the rigidity of effective actions due to the symmetries. Moreover, the symmetry in the IR regime may be even higher than that in the initial UV theory.

The symmetry nature of effective actions results in nontrivial consequences. Presence of an underlying integrable systems seems to be the most important of all. As a rule, the partition function calculated in the effective theory appears to be the τ -function of some integrable system. On the other hand, the τ -function is a generating function for the conservation laws in integrable system and simultaneously the generating function of correlators in some quantum field theory [9, 14].

Identification of the variables in the integrable system describing the effective action is a difficult problem. An invariant way to introduce these variables in non-topological theories is still missing. However, there are several examples of two-dimensional theories when these variables can be identified with the transition amplitudes between vacuum states induced by nonperturbative configurations. The couplings and sources serve as the “space-time variables” in integrable systems. Examples of the systems related to the $2d$ Toda lattice and similar hierarchies (KP, KdV etc) can be borrowed from $2d$ σ -models [15] and $2d$ gravity [16, 17, 18, 19, 20, 9]. Selecting the concrete solution to the nonlinear integrable equations is a separate problem. It is typically done by an additional equation, string equation [16]¹ which is one of the Ward identities imposed on the theory [17, 22].

Although one can fix the general structure of effective action from the symmetry arguments, it is also interesting to evaluate it by the direct integration over heavy modes in the initial field theory. It is implied that there is some energy scale and all modes above this scale are integrated out. This integration should be done over the perturbative configurations and the scale serves a cutoff in the corresponding Feynman loop diagrams. It is also necessary to sum over the nonperturbative configurations, their typical sizes being restricted by this scale. The nonperturbative configurations important in different dimensions are: instantons, monopoles, vortices as well as solitons of different codimension in string theory.

The important feature of nonperturbative effects is that they give rise to the moduli spaces of the corresponding solutions. The very reason for the moduli space to emerge is the high symmetry of the

¹The string equation does not fix the possible discrete ambiguity of solutions [21].

theory. Therefore, the moduli space itself encodes symmetry properties. For instance, the dimension of the instanton moduli space is fixed by the space-time symmetries – translations and rotations as well as by the gauge symmetries [23, 24]. There are many other important moduli spaces, for instance, the moduli spaces: of the complex structures of Riemann surfaces, of the flat connections on a fixed surface, of the holomorphic vector bundles etc. Each of these appears when considering some nonperturbative configurations and can be derived via the reduction procedure from the instanton moduli space. Thus, the universal instanton moduli space comes as a most general object arising in field theories in four or less dimensions.

As we explained, the integration over moduli spaces involved into the derivation of effective actions should ultimately give rise to integrable theories. The same kind of moduli spaces have much to do with integrable systems also from another point of view. Namely, the phase spaces of integrable systems typically coincide with moduli spaces or cotangent bundles to the moduli spaces of the same type. Simple examples are given by solutions of the KdV hierarchy related to the moduli space of the complex structures or by solutions of the $2d$ Toda lattice hierarchy related to the moduli of the flat connections. Another important example is given by the Chern-Simons theory defined on a surface, whose phase space is the moduli space of the flat connections on this surface [25], while any gauge invariant observable can serve as a Hamiltonian. In terms of integrable systems, the problem of calculation of the nonperturbative contributions into effective actions is nothing but the calculations of some vacuum expectation values of observables on moduli spaces (see [25] as an example).

A large class of theories related with the moduli spaces is known as topological theories. They were introduced into quantum field theory by Witten [27] to obtain invariants of the vacuum manifolds in field theories. The first examples of topological theories have been elaborated in four dimensions for σ -models and gauge theories. In the σ -model case, the topological theory provides invariants of the holomorphic maps, while in the gauge theory it gives invariants of the instanton moduli space [28]. At the moment, there are a lot of examples of topological theories in different dimensions, e.g., σ -models in $d = 2$, Chern-Simons theory in $d = 3$, topological Lagrangians in $d = 4$ etc.

In all the cases, the topological theories can be obtained from the nontopological $\mathcal{N} = 2$ SUSY theories via the twisting procedure [29]. The partition function of the topological theories appears to be some invariant of the corresponding moduli space.

There are also some examples of field theories with the topological subsectors related solely with the vacuum states. An example is given by the $\mathcal{N} = 1$ supersymmetric gauge theory, where the perturbative β function [30] as well as some correlators can be calculated exactly [31]. At last, let us mention that the associativity (WDVV) equations which typically emerge in topological theories also appear in $\mathcal{N} = 2$ SUSY theories. This implies some hidden topological structures (subsectors) in these theories as well [32].

Since the moduli spaces arising in different applications are typically finite dimensional, the topological theories that provide finite dimensional moduli spaces are of the prime interest. It is these moduli spaces where the dynamics of integrable many-body systems with nontrivial conservation laws is developing. Gauge invariant observables with appropriate couplings serve the Hamiltonians.

Note that the partition function for the perturbed topological theory, i.e. the generating function for the correlators is the most interesting and informative object. It is simultaneously the generating function for topological invariants of the moduli space manifold [27] and the τ -function of an integrable hierarchy [19, 26].

1.2 Seiberg-Witten solution

New possibilities to deal with nonperturbative physics at strong coupling come from the exact derivation of the low energy effective action for the $\mathcal{N} = 2$ SYM theory [1, 2]. In principle, to get the exact answer one needs to perform the summation of the infinite instanton series. Therefore, the result [1, 2] due to N.Seiberg and E.Witten was the first example of the exact derivation of the total instanton sum.

However, the authors did not calculate instantonic series, using instead indirect and quite sophisticated methods. The calculation of the exact mass spectrum of stable particles was their second essentially new result. What is important, this theory is non-trivial in one loop and belongs to the class of asymptotically free theories. In fact, the situation is a little bit unusual, since, despite the fact that the direct instanton summation has not been performed yet, there are no doubts in the results derived which survives under different tests.

In order to get the effective action unambiguously, the authors of [1, 2] used the three ideas: holomorphy [33], duality [34] and their compatibility with the renormalization group flows. Holomorphy implies that the low energy effective action can be only of the form $\int \text{Im}\mathcal{F}(\Psi)$ depending on a single holomorphic function $\mathcal{F}$ called prepotential. The duality principle which provides the relation between the strong and weak coupling regimes is natural in the UV finite², say $\mathcal{N} = 4$ SUSY [35], theory, where the evident modular parameter built from the coupling constant and the θ -term

$$\tau = \frac{4\pi i}{g^2} + \frac{\theta}{2\pi}, \quad (1.1)$$

is not renormalized and coincides with its asymptotic value. To construct duality transformation in asymptotically free theory explicitly, one needs an artificial trick, to introduce into the game an additional object, a higher genus Riemann surface whose period matrix coincides with the matrix of couplings depending on the values of condensate in a given vacuum.

In the $\mathcal{N} = 2$ SYM theory this scheme is roughly realized as follows [1, 2]. First of all, in the Lagrangian there is a potential term for the scalar fields of the form

$$V(\phi) = \text{Tr}[\phi, \phi^+]^2, \quad (1.2)$$

where the trace is taken over adjoint representation of the gauge group (say, $SU(N_c)$). This potential gives rise to the valleys in the theory, when $[\phi, \phi^+] = 0$. The vacuum energy vanishes along the valleys, hence, the supersymmetry remains unbroken. One may always choose the v.e.v.'s of the scalar field to lie in the Cartan subalgebra $\phi = \text{diag}(a_1, \dots, a_n)$. These parameters a_i can not serve, however, good order parameters, since there is still a residual Weyl symmetry which changes a_i but leave the same vacuum state. Hence, one should consider the set of the gauge invariant order parameters $u_k = \langle \text{Tr}\phi^k \rangle$ that fix the vacuum state unambiguously. Thus, we obtain a moduli space parametrized by the vacuum expectation values of the scalars, which is known as the Coulomb branch of the whole moduli space of the theory. The choice of the point on the Coulomb branch is equivalent to the choice of the vacuum state and simultaneously yields the scale which the coupling is frozen on. At the generic point of the moduli space, the theory becomes effectively Abelian after the condensation of the scalar.

As soon as the scalar field acquires the vacuum expectation value, the standard Higgs mechanism works and there emerge heavy gauge bosons at large values of the vacuum condensate. To derive the effective low energy action one has to sum up the loop corrections as well as the multi-instanton

²In fact, one should require more: not renormalized and, therefore, non-running coupling constants.

contributions. The loop corrections are trivial, since they are vanishing beyond the one loop due to the supersymmetry. The explicit summation over instantons has not been done yet, however, additional arguments allow one to define the effective action indirectly.

The initial action of $\mathcal{N} = 2$ theory written in $\mathcal{N} = 2$ superfields has the simple structure $S(\Psi) = \text{Im}\tau \int \text{Tr}\Psi^2$ with $\tau = \frac{4\pi i}{g^2} + \frac{\theta}{2\pi}$. Now the $\mathcal{N} = 2$ supersymmetry implies that the low energy effective action gets renormalized only by holomorphic contributions so that it is ultimately given by a single function known as prepotential $S_{eff}(\Psi) = \text{Im} \int \mathcal{F}(\Psi)$. The prepotential is a holomorphic function of moduli u_k (or v.e.v.'s) except for possible singular points at the values of moduli where additional massless states can appear disturbing the low energy behaviour.

Thus, the problem effectively reduces to the determination of one holomorphic function. If one manage to fix its behaviour nearby singularities, the function can be unambiguously restored. One of the singularities, corresponding to large values of v.e.v.'s, i.e. to the perturbative limit is under control (since the theory is asymptotically free). All other singularities are treated with the use of duality and of the non-renormalization theorems for the central charges of the SUSY algebra. A combination of these two ideas allows one to predict the spectrum of the stable BPS states which become massless in the deep nonperturbative region and are in charge of all other singularities.

The duality transformation can be easily defined in the finite $\mathcal{N} = 4$ SUSY theory just as the modular transformations generated by $\tau \rightarrow \tau^{-1}$ and $\tau \rightarrow \tau + 1$. This makes a strong hint that the duality can be related with a modular space of some Riemann surfaces, where the modular group acts. In fact, the naive application of duality meets some serious difficulties. The reason is that, in the asymptotically free theory, one has to match the duality with the renormalization group³. This is non-trivial, since now τ depends on the scale which is supposedly involved into the duality transformation. The solution of this problem is that one is still able to connect the duality and modular transformations if considering the theory at different vacua connected by the duality. Then, the duality acts on the moduli space of vacua and this moduli space is associated with the moduli space of the auxiliary Riemann surface, where the modular transformations act.

At the next step, one has to find out proper variables whose modular properties fit the field theory interpretation. These variables are the integrals⁴ of a meromorphic 1-form dS over the cycles on the Riemann surface, a_i and a_i^D

$$\begin{aligned} a_i &= \oint_{A_i} dS, \\ a_D &= \oint_{B_i} dS, \end{aligned} \tag{1.3}$$

(where $i, j = 1, \dots, N_c - 1$ for the gauge group $SU(N_c)$).

³In fact, there are more problems with duality even in $\mathcal{N} = 2$ SUSY theories. In particular, it unifies into one duality multiplet the monopole and gauge boson supermultiplets, while they have different spins, see, e.g., [1, 6].

⁴ We define the symbols $\oint$ and res with additional factors $(2\pi i)^{-1}$ so that

$$\text{res}_0 \frac{d\xi}{\xi} = -\text{res}_\infty \frac{d\xi}{\xi} = \oint \frac{d\xi}{\xi} = 1$$

This explains the appearance of $2\pi i$ factors in the Riemann identities and in perturbative prepotentials in sections 5 and 6. Accordingly, the theta-functions are periodic with period $2\pi i$, and

$$\frac{\partial \theta(\vec{\xi}|\mathcal{T})}{\partial \mathcal{T}_{ij}} = i\pi \partial_{ij}^2 \theta(\vec{\xi}|\mathcal{T})$$

since periods of the Jacobi transformation $\xi_i \equiv \int^\xi d\omega_i$ belong to $2\pi i (\mathbb{Z} + \mathcal{T}\mathbb{Z})$.

These integrals play the two-fold role in the Seiberg-Witten approach. First of all, one may calculate the prepotential $\mathcal{F}$ and, therefore, the low energy effective action through the identification of a_D and $\partial\mathcal{F}/\partial a$ with a defined as a function of moduli (values of condensate) by formula (1.3). Then, using the property of the differential dS that its variations w.r.t. moduli are holomorphic one may also calculate the matrix of coupling constants

$$T_{ij}(u) = \frac{\partial^2 \mathcal{F}}{\partial a_i \partial a_j}, \quad (1.4)$$

The second role of formula (1.3) is that, as was shown these integrals define the spectrum of the stable states in the theory which saturate the Bogomolny-Prasad-Sommerfeld (BPS) limit. For instance, the formula for the BPS spectrum in the $SU(2)$ theory reads as

$$M_{n,m} = |na(u) + ma_D(u)|, \quad (1.5)$$

where the quantum numbers n, m correspond to the “electric” and “magnetic” states. The reader should not be confused with the spectrum derived from the low energy behaviour which fixes arbitrarily heavy BPS states. The point is that the BPS spectrum is related to the central charge of the extended SUSY algebra $Z_{\{m\},\{n\}} = a_i n^i + a_i^D m^i$ [36] and, therefore, has an anomaly origin. On the other hand, anomalies are not renormalized by the quantum corrections and can be evaluated in either of the UV and IR regions.

Note that the column (a_i, a_i^D) transforms under the action of the modular group $SL(2, \mathbb{Z})$ as a section of the linear bundle. Its global behaviour, in particular, the structure of the singularities is uniquely determined by the monodromy data. As we discussed earlier, the duality transformation connects different singular points. In particular, it interchanges “electric”, a_i and “magnetic”, a_i^D variables which describe the perturbative degrees of freedom at the strong and weak coupling regimes (of the theory at different vacua) respectively. Manifest calculations with the Riemann surface allow one to analyze the monodromy properties of dual variables when moving in the space of the order parameters. For instance, in the simplest $SU(2)$ case, on the u -plane of the single order parameter there are three singular points, and the magnetic and electric variables mix when encircling these points. Physically, in the theory with non-vanishing θ -term (not pure imaginary τ), the monopole acquires the electric charge, while the polarization of the instanton medium yields the induced dyons.

1.3 Seiberg-Witten theory and theory of prepotential

Although the both auxiliary objects, the Riemann surface and the differential dS have come artificially, an attempt to recognize them in SUSY gauge theories results into discovery of the integrable structures responsible for the Seiberg-Witten solutions.

Now let us briefly formulate the structures underlying Seiberg-Witten theory. Later on, we often refer to Seiberg-Witten theory as to the following set of data:

- Riemann surface $\mathcal{C}$
- moduli space $\mathcal{M}$ (of the curves $\mathcal{C}$), the moduli space of vacua of the gauge theory
- meromorphic 1-form dS on $\mathcal{C}$

How it was pointed out in [37], this input can be naturally described in the framework of some underlying integrable system.

To this end, first, we introduce bare spectral curve E that is torus $y^2 = x^3 + g_2x^2 + g_3$ for the UV-finite⁵ gauge theories with the associated holomorphic 1-form $d\omega = dx/y$. This bare spectral curve degenerates into the double-punctured sphere (annulus) for the asymptotically free theories (where dimensional transmutation occurs): $x \rightarrow w + 1/w$, $y \rightarrow w - 1/w$, $d\omega = dw/w$. On this bare curve, there are given a matrix-valued Lax operator $L(x, y)$. The corresponding dressed spectral curve $\mathcal{C}$ is defined from the formula $\det(L - \lambda) = 0$.

This spectral curve is a ramified covering of E given by the equation

$$\mathcal{P}(\lambda; x, y) = 0 \quad (1.6)$$

In the case of the gauge group $G = SU(N_c)$, the function $\mathcal{P}$ is a polynomial of degree N_c in λ .

Thus, we have the spectral curve $\mathcal{C}$, the moduli space $\mathcal{M}$ of the spectral curve being given just by coefficients of $\mathcal{P}$. The third important ingredient of the construction is the generating 1-form $dS \cong \lambda d\omega$ meromorphic on $\mathcal{C}$ (“ $\cong$ ” denotes the equality modulo total derivatives). From the point of view of the integrable system, it is just the shortened action “ pdq ” along the non-contractable contours on the Hamiltonian tori. This means that the variables a_i in (1.3) are nothing but the action variables in the integrable system. The defining property of dS is that its derivatives with respect to the moduli (ramification points) are holomorphic differentials on the spectral curve. This, in particular, means that

$$\frac{\partial dS}{\partial a_i} = d\omega_i \quad (1.7)$$

where $d\omega_i$ are the canonical holomorphic differentials⁶. Integrating this formula over B -cycles and using that $a_D = \partial\mathcal{F}/\partial a$, one immediately obtains (1.4).

So far we reckoned without matter hypermultiplets. In order to include them, one just needs to consider the surface $\mathcal{C}$ with punctures. Then, the masses are proportional to the residues of dS at the punctures, and the moduli space has to be extended to include these mass moduli. All other formulas remain in essence the same (see [39] for more details).

Note that it is still unknown whether the set of data above is identically equivalent to an integrable system. Inversely, it is associated only with specific integrable systems, which can be called analytic integrable systems. Although being wide enough, this class of integrable systems far does not cover the systems integrable in the Liouville sense.

The prepotential $\mathcal{F}$ and other “physical” quantities are defined in terms of the cohomology class of dS , formula (1.3). Note that formula (1.4) allows one to identify the prepotential with logarithm of the τ -function of the Whitham hierarchy [40]: $\mathcal{F} = \log \mathcal{T}_{\text{Whitham}}$. The other important property of the prepotential related to the feature of the Seiberg-Witten framework which merits the adjective

⁵The situation is still unclear in application to the case of fundamental matter with the number of matter hypermultiplets N_f equal to $2N_c$. In existing formulation for spin chains the bare coupling constant appears rather as a twist in gluing the ends of the chain together [38] (this parameter occurs only when $N_f = 2N_c$) and is not immediately identified as a modulus of a *bare* elliptic curve. This problem is a fragment of a more general puzzle: spin chains have not been described as Hitchin systems; only the “ 2×2 ” Lax representation is known for them, while its “dual” $N_c \times N_c$ one is not yet available.

⁶I.e. satisfying the conditions

$$\oint_{A_i} d\omega_j = \delta_{ij}, \quad \oint_{B_i} d\omega_j = T_{ij}$$

”topological” has much to do with the associative algebras [32]. Namely, it turns out that the prepotential of Seiberg-Witten theory satisfies a set of the Witten-Dijkgraaf-Verlinde-Verlinde (WDVV) equations. These equations have been originally presented in [41] (in a different form, see below)

$$F_i F_j^{-1} F_k = F_k F_j^{-1} F_i \quad (1.8)$$

where F_i ’s are matrices with the matrix elements that are the third derivatives of a unique function F of many variables a_i ’s (prepotential in the Seiberg-Witten theory) parameterizing a moduli space:

$$(F_i)_{jk} \equiv \frac{\partial^3 F}{\partial a_i \partial a_j \partial a_k}, \quad i, j, k = 1, \dots, n \quad (1.9)$$

Although generally there are a lot of solutions to the matrix equations (1.8), it is extremely non-trivial task to express all the matrix elements through the only function F . In fact, there have been only known the two different classes of non-trivial solutions to the WDVV equations, both being intimately related to the two-dimensional topological theories of type A (quantum cohomologies [42]) and of type B ($\mathcal{N} = 2$ SUSY Ginzburg-Landau theories that were investigated in a variety of papers, see, for example, [40] and references therein). Thus, the existence of a new class of solutions connected with the four-dimensional (and higher dimensional) theories looks quite striking.

1.4 Duality, branes and gauge theories

The progress in the description of the low energy SUSY gauge theories initiated the investigation of duality in string theory. It appeared that, in order to formulate the duality transformation in string theory, one has to introduce new objects called D-branes [43]. Their main property is that the open strings can end on the D-brane. Therefore, the Dirichlet boundary condition can be imposed. Moreover, the charge in the RR sector of the theory is attributed to the D-brane. Using D-branes, a stringy duality can be self-consistently described, and actually three different types of dualities can be introduced [44].

Since the superstring theory is anomaly free only in ten dimensions, the question about the role of remaining dimensions naturally arises. The results of the last years definitely indicate that these dimensions should be identified with the zero modes of the scalar fields [45] presented in $4d$ field theories. In the most symmetric situation in $d = 4$, which is the $\mathcal{N} = 4$ SUSY gauge theory there are 3 complex scalars whose vacuum expectation values should be considered as the six compactified dimensions. Generically the structure of the field space is very complicated and its metric is the subject of numerous investigations.

From the field theory point of view, $\mathcal{N} = 4$ is maximal admissible supersymmetry, since it is the case when the gravitational degrees of freedom are still decoupled [46] from the gauge sector of the theory, the case of our prime interest.

Now, in order to analyze a generic $4d$ gauge theory, one has to introduce an arbitrary gauge group, to ask for the proper SUSY and to define all the parameters in the theory such as masses and couplings. It appears that the account of the brane degrees of freedom allows one to fulfill this program at full. The important properties of branes which allow one to obtain these results are the presence of an $U(1)$ gauge field on their worldvolumes as well the existence of brane bound states [45].

Let us briefly describe how the $4d$ field theory can be formulated as a theory on the brane world volume of some dimension. At the first step, we start with the superstring theory in ten dimensions of type II, and branes are the BPS states breaking half of the supersymmetries (this gives us $\mathcal{N} =$

4 SUSY in $4d$). In one approach, the initial theory is considered in the flat space and SUSY is broken by the background brane configuration [47]. In the other approach, known as “geometrical engineering” [48] the supersymmetry is broken by a special choice of metric of the six dimensional compactification manifold. For instance, the metric of the $K3$ manifold gives rise to breaking of the half of supersymmetries.

As a corollary of definition, the strings ending on the D-brane induce an $U(1)$ gauge field living on its worldvolume. At the next step, one should produce the general gauge group, say $SU(N_c)$, which is absent in the original string theory. Once again two approaches are possible; in the first approach, the gauge group arises from the bound state of branes [45], while in the second approach it originates from choice of the compactification manifold with the singularity structure related to the gauge group root system [49]. The higher rank gauge group obtained from the brane bound state can be explained as follows. Each of the branes carries the $U(1)$ index while the open strings can stretch between them. Therefore, having the stack of parallel N_c branes, each string can be associated with an $U(N_c)$ matrix realizing the $U(N_c)$ gauge theory⁷. If one now splits the branes, there are massive modes of the gauge fields in the spectrum of open strings which correspond to the strings stretched between different branes and the corresponding masses are proportional to the distance between branes in the embedding space. This is a realization of the Higgs mechanism. When the distance between branes vanishes, rank of the corresponding “unbroken” gauge group increases. In the “geometrical engineering” approach the gauge group comes from the wrapping of branes over the manifold with the corresponding singularity structure [50].

Note that the scalars in field theory play the role of coordinates of the brane or the bound state of branes in the $6d$ space.

The next step of this rough (rather pictorial) construction of field theory is to introduce massive hypermultiplets and couplings. To this end, one should further break the supersymmetry⁸. It can be done by adding more, background branes, of different codimensions and non-parallel to the stack of N_c branes.

Say, one can introduce hypermultiplets by adding branes of other codimensions so that the coordinates of these branes in $6d$ compactification manifold correspond to the values of their masses [47]. Couplings are introduced similarly and their values are identified with the distance between the background branes in one picture or by the size of the compact manifold in the “geometrical engineering” approach [48].

1.5 Seiberg-Witten solution and branes

Realization of the Seiberg-Witten solution of the $\mathcal{N} = 2$ SUSY gauge theory in brane terms seems to be the most impressive realization of the approach described. In the type IIA framework, the theory is solved along the following line [51]. The $SU(N_c)$ gauge theory arises on the worldvolume of N_c parallel D4 branes embedded into the flat $10d$ manifold. The worldvolume of the branes is five dimensional and stretches along, say, x_0, x_1, x_2, x_3, x_6 -directions, the branes being of finite size in the x_4 -direction, with the first four coordinates identified with the coordinates in our (infinite) $4d$ world.

Now one has to describe the compact six dimensions in field theory terms. To this end, one first

⁷It can be shown that the total $U(1)$ factor decouples.

⁸Note that, due to the conformal invariance, the $\mathcal{N} = 4$ SUSY theory depends on the coupling constant trivially and can not contain massive hypermultiplets.

assumes that all the branes are placed at the origin with respect to three coordinates⁹, say, x_7, x_8, x_9 . Now, the two remaining directions not stretching along the brane, x_4 and x_5 form the holomorphic variable, $x_4 + ix_5$ which is associated with the vacuum expectation value of the scalar in the $\mathcal{N} = 2$ SUSY gauge theory, that is to say, the coordinate on the Coulomb branch of the moduli space. At last, the compact coordinate of brane, x_6 is just the coordinate on (real section of) the auxiliary Riemann surface.

Since the D4 branes are restricted along the x_6 -direction, one needs some other, background branes where the D4 branes should end on. It is to be two additional branes of the NS type. It is just this configuration which breaks the $\mathcal{N} = 8$ SUSY in ten dimensions (32 generators) down to the $\mathcal{N} = 2$ SUSY in four dimensions (8 generators). The distance between the background branes along the x_6 -direction is identified with the coupling constant in the gauge theory $\delta x_6 = \frac{1}{g^2}$, while the back reaction to the background brane can be attributed to the perturbative renormalization of the coupling. In order to incorporate the fundamental matter hypermultiplets into the theory, one has to introduce additional N_f D6 branes whose coordinates¹⁰ along $x_4 + ix_5$ are related with masses of the hypermultiplets. On the other hand, in order to add the adjoint matter, the (complexified) coordinate x_6 should be made periodic.

The configuration considered is singular, since it is impossible to describe the intersection of the D4 and background NS5 branes smoothly. To resolve the singularities, Witten has suggested to lift the configuration into $d = 11$ manifold which plays the same role for membrane theory as the $d = 10$ space for superstrings. The theory defined in $d = 11$ is known as M-theory and it has the $d = 11$ supergravity as its low energy limit [52]. It is important that there are solitons in M-theory, that is, M5 branes and membranes (M2 branes). In the non-singular M-theory picture of $\mathcal{N} = 2$ SUSY gauge theories, the eleventh dimension is assumed to be compact and there is a single M5 brane wrapped around it. This M5 brane is assumed to be topologically non-trivial and to have 2 compact and 4 infinite dimensions, these latter being associated with our $4d$ world. With three coordinates (x_7, x_8, x_9) being set to zero, the two compact dimensions of the M5 brane are embedded into the four-dimensional space (of x_4, x_5, x_6, x_{10} coordinates) in a tricky way. This embedding is, however, holomorphic and describes the auxiliary Riemann surface. Being projected onto ten dimensions, different pieces of this two-dimensional surface are associated with the D4 and NS5 branes.

Note now that a similar picture has been also developed in the M-theory framework for $\mathcal{N} = 1$ gauge theories [53] which manifest confinement due to the monopole condensation. This gives some hope for relevance of the brane picture also for the ordinary, non-supersymmetric QCD.

In the other IIB approach [54] related to the previous consideration via the T duality transformation [55], the embedding manifold is topologically non-trivial and branes are assumed to be wrapped around some cycles on the surfaces, for instance, on tori (elliptic curves) embedded into K3 manifold [56]. This approach is less transparent, although the topological theories on the curves which are associated with the integrable many-body systems arise within this framework automatically. Similarly to the way of resolving the singular IIA configuration by lifting it into higher dimensional M-theory, the IIB configuration has to be lifted to the $d = 12$ theory known as F-theory [57]. The additional two dimensions are assumed to be compactified onto the torus whose modular parameter is identified with the (imaginary and real parts of) UV values of the coupling constant and the θ -term in the gauge theory.

⁹This is necessary to preserve, at least, $\mathcal{N} = 2$ SUSY.

¹⁰More precisely, the distance to the D4 branes.

2 Many-body systems and gauge theories in two and three dimensions

In this section we demonstrate how the many-body systems naturally arise in two and three dimensional gauge theories and the phase space of the system of particles is interpreted in terms of the topological gauge degrees of freedom. From what follows it becomes clear that the $2d$ YM theory is closely related to the Calogero-Moser system [58, 59, 60], while the Chern-Simons theory to its relativistic counterpart [61, 62], the Ruijsenaars-Schneider system.

2.1 Calogero many-body system and Yang-Mills theory in two dimensions

Let us show how the Lagrangian formulation of the Calogero systems can be related to the YM theory on a cylinder [63]. Remind that the Calogero system is defined as an N particle system with the pairwise interaction [58, 59]:

$$V(x_1, \dots, x_N) = \sum_{i \neq j} \frac{\nu^2}{(x_i - x_j)^2}, \quad (2.1)$$

ν^2 is the coupling constant. As we already mentioned, the phase spaces of integrable systems are typically related with group manifolds. Therefore, it should not be surprising that the Lagrangians under consideration have a group origin.

Let us start with the central extension $\widehat{\mathfrak{g}}$ of the loop algebra $\mathcal{L}\mathfrak{g}$, where $\mathfrak{g}$ is a semi-simple Lie algebra. It is defined as the space of pairs (ϕ, c) , where ϕ is the $\mathfrak{g}$ -valued scalar field given on the circle and c is the central element, just a real number. The Lie algebra structure is introduced by formula

$$[(\phi_1(\varphi), c_1), (\phi_2(\varphi), c_2)] = \left([\phi_1(\varphi), \phi_2(\varphi)], \int_{S^1} \langle \phi_1, \partial_\varphi \phi_2 \rangle \right), \quad (2.2)$$

where $\langle \cdot, \cdot \rangle$ denotes the Killing form on the algebra $\mathfrak{g}$.

Now we set the cotangent bundle to this algebra, $T^*\widehat{\mathfrak{g}}$ to be a “bare” phase space. $T^*\widehat{\mathfrak{g}}$ is defined to be a direct sum of $\widehat{\mathfrak{g}}$ and $\widehat{\mathfrak{g}}^*$, i.e. the space of sets $(A, k; \phi, c)$, where A is the gauge field on the circle and k is dual to c . The pairing between $\widehat{\mathfrak{g}}$ and $\widehat{\mathfrak{g}}^*$ is given by

$$\langle (A, \kappa); (\phi, c) \rangle = \int_{S^1} \langle \phi, A \rangle + c\kappa, \quad (2.3)$$

On this cotangent bundle, one can define the natural symplectic structure

$$\Omega = \int_{S^1} Tr(\delta\phi \wedge \delta A) + \delta c \wedge \delta \kappa \quad (2.4)$$

The adjoint and coadjoint actions of the loop group $\mathcal{L}G$ on $\widehat{\mathfrak{g}}$ and $\widehat{\mathfrak{g}}^*$ are given by formulas

$$(\phi(\varphi), c) \rightarrow \left(g(\varphi)\phi(\varphi)g(\varphi)^{-1}, \int_{S^1} Tr(-\phi g^{-1}\partial_\varphi g) + c \right) \quad (2.5)$$

$$(A, \kappa) \rightarrow (gAg^{-1} + \kappa g\partial_\varphi g^{-1}, \kappa) \quad (2.6)$$

This action clearly preserves the symplectic structure and thus defines a moment map

$$\mu : T^*\widehat{\mathfrak{g}} \rightarrow \widehat{\mathfrak{g}}^*, \quad (2.7)$$

which sends $(\phi, c; A, \kappa)$ to $(\kappa\partial_\varphi\phi + [A, \phi], 0)$.

Note that the choice of level of the moment map has a simple physical meaning: in the theory of the angular momentum it just fixes the sector of the Hilbert space. In the gauge theory, one selects the gauge invariant states.

Let us now discuss how to choose the proper level of the moment map in order to get further the Hamiltonian reduction w.r.t. this level. Let us remind that in the case of finite-dimensional group $\mathbf{G}$ one takes an element J from $\mathfrak{g}^*$, which has the maximal stabilizer different from the whole $\mathbf{G}$. It is easy to show that the representative of the coadjoint orbit of this element has the following form

$$J_\nu = \nu \sum_{\alpha \in \Delta_+} (e_\alpha + e_{-\alpha}), \quad (2.8)$$

where ν is some real number, $e_{\pm\alpha}$ are the elements of the nilpotent subalgebras, $\mathfrak{n}_\pm \subset \mathfrak{g}$ corresponding to the root α , and Δ_+ is the set of positive roots. Let us denote the coadjoint orbit of J_ν as $\mathcal{O}_\nu$ and the representation of $\mathbf{G}$ arising upon the quantization of $\mathcal{O}_\nu$ as R_ν . For a generic $J \in \mathfrak{g}^*$, let $\mathbf{G}_J$ be the stabilizer of J in the coadjoint orbit $\mathcal{O}_J$, thus $\mathcal{O}_J = \mathbf{G}/\mathbf{G}_J$.

Now in order to deal with the affine case, one just needs to substitute the roots by their affine counterparts. There is also a more geometrical standpoint: because of the vanishing level $k = \mu$ of the moment map, the coadjoint orbit of its generic value is huge, while the stabilizer of the element $(J(\varphi); 0) \in \widehat{\mathfrak{g}}^*$ is “a continuous product” $\prod_{\varphi \in S^1} \mathbf{G}_{J(\varphi)}$ which is very small as compared with the whole loop group $\mathcal{L}\mathbf{G}$ unless the group $\mathbf{G}_{J(\varphi)}$ coincides with $\mathbf{G}$ for almost all $\varphi \in S^1$.

All this implies that one can reasonably choose the value of the moment map to be

$$\mu = (\mathcal{J}[\mu], 0) : \mathcal{J}[\mu](\varphi) = \delta(\varphi)J_\nu \quad (2.9)$$

The coadjoint orbit of μ is nothing but the finite dimensional orbit $\mathcal{O}_\nu$.

Now in order to perform the Hamiltonian reduction, one has to resolve the moment map¹¹

$$\kappa\partial_\varphi\phi + [A, \phi] = J_\mu(\varphi) = \delta(\varphi)J_\nu. \quad (2.10)$$

It can be done in the following way. First, we use the generic gauge transformation $g(\varphi)$ to make A to be a Cartan subalgebra valued constant one-form D . However, we are still left with the freedom to use the constant gauge transformations which do not touch D . Actually the choice of D is not unique and is parametrized by the conjugacy classes of the monodromy $\exp(\frac{2\pi}{\kappa}D) \in \mathbf{T} \subset \mathbf{G}$. Let us fix a conjugacy class and denote ix_i the elements of the matrix $D = iQ$. Decompose $\mathfrak{g}$ -valued function ϕ on the S^1 on the Cartan-valued $P(\varphi) \in \mathfrak{t}$ and nilpotent-valued $\phi_\pm(\varphi) \in \mathfrak{n}_\pm$ components. Let $\phi_\alpha = \langle \phi, e_\alpha \rangle$. Then (2.10) reads:

$$\kappa\partial_\varphi P = \delta(\varphi)[J_\nu^g]_\gamma \quad (2.11)$$

$$\kappa\partial_\varphi\phi_\alpha + \langle D, \phi_\alpha \rangle = \delta(\varphi)[J_\nu^g]_\alpha, \quad (2.12)$$

where $\widetilde{J}_\nu^g$ is $Ad_{g(0)}^*(J_\nu)$, $[J]_\gamma$ denotes the Cartan part of J and $[J]_\alpha = \langle J, e_\alpha \rangle$.

From (2.11) one gets $D = const$ and $[J_\nu^g]_\gamma = 0$. Therefore, one can twist back J_ν^g to J_ν . Moreover, (2.12) implies that (if $\varphi \neq 0$) $\phi_\alpha(\varphi)$ can be presented as

¹¹This is nothing but the Gauss law.

$$\phi_\alpha(\varphi) = \exp\left(-\frac{\varphi}{\kappa} < D, \alpha >\right) \times M_\alpha, \quad (2.13)$$

where M_α is a locally constant element in $\mathfrak{g}$. From (2.12) one sees that M_α jumps, when φ passes through 0. The jump is equal to

$$\left[\exp\left(-\frac{2\pi}{\kappa} < D, \alpha >\right) - 1\right] \times M_\alpha = [J_\nu^g]_\alpha. \quad (2.14)$$

The final conclusion is that the physical degrees of freedom are contained in $\exp(-\frac{2\pi i}{\kappa}Q)$ and P with the symplectic structure

$$\omega = \frac{1}{2\pi i} \text{Tr}(\delta P \wedge \delta Q) \quad (2.15)$$

and one has

$$\phi_\alpha(\varphi) = \nu \frac{\exp(-\frac{i\varphi}{\kappa} < Q, \alpha >)}{\exp(-\frac{2\pi i}{\kappa} < Q, \alpha >) - 1}$$

With this manifest expression, one easily restores the whole $\phi(\varphi)$ adding the constant Cartan part. Note that, while the diagonal elements q_i of Q determine particle coordinates, the constant Cartan part which can not be fixed from the moment map is parameterized by particle momenta p_i . In these variables, $\phi(\varphi)$ is nothing but the Calogero Lax matrix and the particles are the Calogero ones.

Now if one takes a simple Hamiltonian system on $T^*\widehat{\mathfrak{g}}$ with a Hamiltonian invariant under the action (2.5),(2.6), this induces a relatively complicated system on the reduced symplectic manifold, $T^*\mathbb{T}$. As a Hamiltonian on $T^*\widehat{\mathfrak{g}}$, one can choose

$$\mathcal{H} = \int_{S^1} d\varphi P(\phi) \quad (2.16)$$

where $P(\phi)$ is any invariant polynomial on the Lie algebra $\mathfrak{g}$. This is, indeed, the standard way to construct Hamiltonians from the Lax matrix in integrable system.

Let us choose the Hamiltonian to be the quadratic Casimir

$$\mathcal{H}_2 = \int_{S^1} d\varphi < \phi, \phi >, \quad (2.17)$$

and look at the action of this system. Taking into account the constraint, i.e. adding to the action the moment map with the Lagrangian multiplier, one obtains

$$S = \int_{S^1 \times Z} d\varphi dt (\text{Tr} A \partial_t \phi - < \phi, \phi > - \frac{1}{\kappa} \text{Tr} A_0 (\kappa \partial_\varphi \phi + [A, \phi] - \delta(\varphi) J_\nu)). \quad (2.18)$$

Denoting $A_1 \equiv A$ and $g_{YM} \equiv 1/\kappa$, one can rewrite this action in the form

$$S = \int_{S^1 \times Z} d^2x \left[-\text{Tr} (\phi F_{01} + \phi^2) + g_{YM} \text{Tr} \left(J_\nu \int_C A_\mu dx^\mu \right) \right] \quad (2.19)$$

where F_{01} is the component of the gauge field strength tensor and C is the contour passing through $\varphi = 0$ along the time axis. This action of induced gauge theory, after integrating over the scalar field ϕ can be rewritten as the action of the 2d YM theory with an additional Wilson line inserted

$$S = \int_{S^1 \times Z} d^2x \left[-\frac{1}{4} \text{Tr} F_{\mu\nu}^2 + g_{YM} \text{Tr} \left(J_\nu \int_C A_\mu dx^\mu \right) \right] \quad (2.20)$$

On the reduced manifold one obtains from the Hamiltonian (2.17) the trigonometric Calogero-Sutherland system [59] with the Hamiltonian

$$H_2 = \frac{1}{2} \text{Tr} P^2 + \sum_{\alpha \in \Delta_+} \frac{\nu^2}{\sin^2 \langle Q, \alpha \rangle}, \quad (2.21)$$

For instance, for $G = SU(N)$ one obtains the Hamiltonian with the pairwise interaction

$$V_{ij}^A = \frac{\nu^2}{\sin^2(q_i - q_j)} \quad (2.22)$$

while for $G = SO(2N)$ one has

$$V_{ij}^D = \nu_2^2 \left[\frac{1}{\sin^2(q_i - q_j)} + \frac{1}{\sin^2(q_i + q_j)} \right] + \nu_1^2 \left[\frac{1}{\sin^2(q_i)} + \frac{1}{\sin^2(q_j)} \right] \quad (2.23)$$

where ν_1, ν_2 are the coupling constants. In the general case, one has as many different coupling constants as many orbits the Weyl group has in the root system.

Generalization of the consideration to the generic Wilson line which amounts to the spin trigonometric Calogero system was considered in [64].

2.2 Gauged G/G σ -model and Ruijsenaars systems

In this subsection we show that the Calogero system have a natural relativization which also admits an interpretation in field theory terms. The corresponding field theory (given on the phase space before the reduction) is nothing but the gauged G/G sigma model which is equivalent to the Chern-Simons Lagrangian [65], the central charge of the Kac-Moody algebra being just the deformation parameter, from the loop group point of view.

Note that this relativization arises in different physical problems, for instance, in two dimensional systems [66] and in gauge theories in higher dimensions [67, 68, 69]. Hence, the unified Lagrangian description for these different physical problems is of evident importance.

As in the non-relativistic system, we perform the Hamiltonian reduction procedure, however now the unreduced phase space is the cotangent bundle to the central extended loop group $\widehat{G}$. As a symplectic manifold, it is presented by the following data: $(g : S^1 \rightarrow G, c \in U(1); A \in \Omega^1(S^1) \otimes \mathfrak{g}^*, \kappa \in \mathbf{R})$.

The group acts as

$$\begin{aligned} g &\rightarrow hgh^{-1}, \quad A \rightarrow hAh^{-1} + \kappa h \partial_\varphi h^{-1} \\ \kappa &\rightarrow \kappa, \quad c \rightarrow c \times \mathcal{S}(g, h), \end{aligned} \quad (2.24)$$

where $\mathcal{S}$ is constructed from the $U(1)$ -valued 2-cocycle $\Gamma(g, h)$ on the group $\mathcal{LG}$ ($\Gamma(g, g^{-1})$),

$$\mathcal{S}(g, h) = \Gamma(h, g)\Gamma(hg, h^{-1}). \quad (2.25)$$

Then, the group invariant symplectic structure looks as follows

$$\Omega = \int_{S^1} \text{Tr} [A(g^{-1}\delta g)^2 + \delta A \wedge g^{-1}\delta g] + \kappa \partial_\varphi g \dot{g}^{-1} (\delta g \dot{g}^{-1})^2 - \kappa \delta(\partial_\varphi g) \dot{g}^{-1} \delta g \dot{g}^{-1}] + c^{-1} \delta c \wedge \delta \kappa \quad (2.26)$$

The generalization of the Gauss law can be found from the group action

$$\mu(g, c; A, \kappa) = (gAg^{-1} + \kappa g \partial_\varphi g^{-1} - A, 0) \quad (2.27)$$

At the next step, we fix the level of the moment map. It is natural to assume that its color structure does not change keeping the integrability intact

$$\mu(g, c; A, \kappa) = i\nu \left(\frac{1}{N} \text{Id} - e \otimes e^+ \right) \delta(\varphi). \quad (2.28)$$

As before, we make A a diagonal matrix D by a generic gauge transformation $h(\varphi)$ modulo the action of the affine Weyl group.

The moment map equation now looks

$$gDg^{-1} + \kappa g \partial_\varphi g^{-1} - D = i\nu \left(\frac{1}{N} \text{Id} - f \otimes f^+ \right) \delta(\varphi) \quad (2.29)$$

where $f = h(\varphi)e$ is the vector from $\mathbf{C}^N$ with the unit norm $\langle f, f \rangle = 1$. Using the residual gauge symmetry, one can make $f \in \mathbf{R}^N$, and using (2.29) one immediately gets

$$g = \exp\left(\frac{\varphi}{\kappa} D\right) G(\varphi) \exp\left(-\frac{\varphi}{\kappa} D\right); \quad \partial_\varphi G = -\frac{J}{\kappa} G \delta(\varphi), \quad (2.30)$$

where $J = i\nu(\frac{1}{N} \text{Id} - f \otimes f^+)$. It is useful to introduce the notation for the monodromy of connection D : $Z = \exp(-\frac{2\pi}{\kappa} D) = \text{diag}(z_1, \dots, z_N)$, $\prod_i z_i = 1$, $z_i = \exp(\frac{2\pi i q_i}{\kappa})$. It satisfies the boundary condition

$$\tilde{G}^{-1} Z \tilde{G} = \exp\left(\frac{2\pi J}{\kappa}\right) Z \quad (2.31)$$

where $\tilde{G} = G(+0)$.

Solving this equation is a separate interesting issue, see [65]. The result looks as follows. First, introduce the characteristic polynomial $P(z)$ of the matrix Z

$$P(z) = \prod_i (z - z_i). \quad (2.32)$$

Define also

$$Q^\pm(z) = \frac{P(\lambda^{\pm 1} z) - P(z)}{(\lambda^{\pm N} - 1) z P'(z)}, \quad (2.33)$$

where $\lambda = e^{\frac{2\pi i \nu}{N\kappa}}$. At $\lambda \rightarrow 1$, the rational functions $Q^\pm(z)$ tend to $\frac{1}{N}$. In these notations, the matrix $\tilde{G}$ can be put in the form

$$\begin{aligned} \tilde{G}_{ij} &= -\lambda^{-\frac{N-1}{2}} \frac{\lambda^{-N} - 1}{\lambda^{-1} z_i - z_j} e^{ip_i} (Q^+(z_i) Q^-(z_j))^{1/2} = \\ &= e^{ip_i - \frac{\pi i}{\kappa} (q_i + q_j)} \frac{\sin(\frac{\pi \nu}{\kappa})}{\sin(\frac{\pi(q_{ij} - \frac{\nu}{N})}{\kappa})} \prod_{k \neq i, l \neq j} \frac{\sin(\frac{\pi(q_{ik} + \frac{\nu}{N})}{\kappa})}{\sin(\frac{\pi q_{ik}}{\kappa})} \frac{\sin(\frac{\pi(q_{il} - \frac{\nu}{N})}{\kappa})}{\sin(\frac{\pi q_{il}}{\kappa})} \end{aligned} \quad (2.34)$$

where p_i are the quantities that can not be determined from the moment map equation and, similarly to the Calogero case, are identified with particle momenta, while q_i are particle coordinates, the symplectic structure on the reduced manifold being $\sim \sum_i \delta p_i \wedge \delta q_i$.

In order to construct Hamiltonians, one may fix any Ad -invariant function $\chi(g)$ and define

$$H_\chi = \int_{S^1} d\varphi \chi(g). \quad (2.35)$$

Such a Hamiltonian of the dynamical system on $T^*\hat{G}$ is evidently invariant under the group action.

For instance, one may take $\chi_\pm(g) = \text{Tr}(g \pm g^{-1})$ (where the trace, Tr is taken over the fundamental representation of $SU(N)$) obtaining

$$H_\pm = \sum_i (e^{i\vartheta_i} \pm e^{-i\vartheta_i}) \prod_{j \neq i} f(q_{ij}) \quad (2.36)$$

where the function $f(q)$ is

$$f^2(q) = [1 - \frac{\sin^2(\pi\nu/\kappa N)}{\sin^2(\pi q/\kappa)}], \quad (2.37)$$

One may recognize in this Hamiltonian the Hamiltonian of the trigonometric Ruijsenaars-Schneider model [61, 62], while the solution for $g(\varphi)$ constructed above gives the Ruijsenaars Lax matrix. Redefining $q_i \rightarrow \kappa q_i$ and $\vartheta_i \rightarrow \vartheta_i/\kappa$, which does not change the symplectic form, and taking the limit $\kappa \rightarrow \infty$, we return back to the Calogero-Sutherland system.

Now we come to the question what is the field theory our system corresponds to. This time we choose the Hamiltonian to be zero so that the term $\int pdq$ serves as a free action and we also add to the action the constraint, the moment map with the Lagrangian multiplier

$$S(A, g) = \int d\varphi dt \text{Tr} [A_1 g^{-1} \partial_t g + \kappa \partial_t g \cdot g^{-1} \cdot \partial_\varphi g \cdot g^{-1} - \kappa d^{-1} (\partial_\varphi g \cdot g^{-1} (dg \cdot g^{-1})^2) - \frac{1}{\kappa} A_0 (\kappa g^{-1} \partial_\varphi g + g^{-1} A_1 g - A_1 - J\delta(\varphi))] \quad (2.38)$$

where we have omitted the term $\kappa c^{-1} \partial_t c$. This action is the action of the G/G gauged σ -model with the Wilson line inserted.

Now we present the equivalent representation of this theory in terms of the Chern-Simons theory with some Hamiltonian. The Chern-Simons theory is the gauge theory given on a three manifold which we choose to be the product of an interval and a torus $X = I \times T^2$, with the action

$$S_{CS} = \frac{i\kappa}{4\pi} \int_X d^3x \text{Tr} (A \wedge dA + \frac{2}{3} A \wedge A \wedge A). \quad (2.39)$$

We are going to consider the theory with an inserted Wilson line. The phase space of this theory is the moduli space of the flat connections on the torus with a marked point with prescribed conjugacy class of the monodromy of connection around the marked point. The path integral is of the form

$$\int \mathcal{D}A \langle v_1 | \text{Tr}(P \exp \int A) | v_2 \rangle \exp(-S_{CS}(A)). \quad (2.40)$$

The monodromy class U around the marked point is fixed by the highest weight $\mathfrak{h}$ of the representation R_ν , $U = \exp(\frac{2\pi i}{\kappa+N} \text{diag}(\mathfrak{h}_i))$. The monodromy condition on the torus $\Sigma = T^2$ that relates the monodromies along the cycles, g_A, g_B and that around the marked point g_C is

$$g_A g_B g_A^{-1} g_B^{-1} = g_C \quad (2.41)$$

With the condition on the conjugacy class of g_C and taking into account that for the representation R_ν the signature is like $\nu \cdot \text{diag}(N, 0, \dots, 0)$, one can get equation (2.31) (see [65] for detailed explanations).

Let us also point out the equivalence of the sector of the Chern-Simons theory on $\Sigma \times S^1$ with observables depending on A_0 , ∂_t being tangent to S^1 , and of the gauged $\mathbf{G}/\mathbf{G}$ theory. This can be proved, say, by gauge fixing A_0 constant and diagonal.

The equivalence provides one with a tool for computation of the spectrum of the Ruijsenaars-Schneider model which is contained in the spectrum of the relativistic free particles on the circle. In the pure Chern-Simons theory, the particles obey the Fermi statistics, since the phase space is $(\mathbf{T}^{N-1} \times \mathbf{T}^{N-1})/\mathcal{S}_N$, the moduli space of the flat connections on the torus. Here $\mathbf{T}^{N-1}$ is the Cartan torus in $SU(N)$, and $\mathcal{S}_N$ is the symmetric group. Since the wave function vanishes at the diagonal in the $N - 1$ -dimensional torus, these particles really behave like fermions.

The Shrödinger equation for the relativistic particle is the difference equation, since $\cos(\frac{2\pi i}{\kappa+N}\partial_q)$ is a finite-difference operator. Its eigenfunctions on the circle are $e^{2\pi i n q}$ with the eigenvalues

$$E_n = \cos\left(\frac{2\pi n}{\kappa+N}\right) \quad (2.42)$$

Then, the complete spectrum of the model reads

$$\sum_i E_{n_i} \quad (2.43)$$

with additional conditions that follow from the invariance $E_n = E_{n+\kappa+N}$ and the symmetry group action

$$(\kappa+N) > n_N > \dots > n_i > \dots > n_1 \geq 0 \quad (2.44)$$

$$\sum_i n_i < (\kappa+N) \quad (2.45)$$

2.3 Elliptic Calogero model and gauge theories

There is also another generalization of the trigonometric Calogero model, the elliptic model. In next sections we shall see that just the elliptic systems are responsible for nonperturbative dynamics in the vacuum sector of the softly broken $\mathcal{N} = 4$ SUSY gauge theories, i.e. of the $\mathcal{N} = 2$ SUSY gauge theories with the adjoint matter hypermultiplet.

In this subsection we show how the elliptic models are described via Hamiltonian reduction of the holomorphic theories in three dimensions [70]. Let us consider a Hamiltonian system whose phase space is the cotangent bundle to the algebra of $SL(N, \mathbb{C})$ -valued functions on the elliptic curve with modular parameter τ , $T^*\widehat{\mathbf{g}}^{\Sigma_\tau}$:

$$(\phi, c; \kappa \bar{\partial} + \bar{A}), \quad \phi : \Sigma_\tau \rightarrow sl_N(\mathbb{C}), \quad \bar{A} \in \Omega^{(0,1)}(\Sigma_\tau) \otimes sl_N(\mathbb{C}), \quad c, \kappa \in \mathbb{C}$$

Let us also denote through $d\omega$ the holomorphic differential with integrals over the A - and B -cycles equal to 1 and τ , respectively.

On this phase space acts naturally the current group $SL_N(\mathbb{C})^{\Sigma_\tau}$

$$\begin{aligned}
\phi(z, \bar{z}) &\rightarrow g(z, \bar{z})\phi(z, \bar{z})g(z, \bar{z})^{-1} \\
\bar{A}(z, \bar{z})d\bar{z} &\rightarrow g(z, \bar{z})\bar{A}(z, \bar{z})g(z, \bar{z})^{-1} + \kappa g(z, \bar{z})\bar{\partial}g(z, \bar{z})^{-1} \\
\kappa &\rightarrow \kappa, \quad c \rightarrow c + \int_{\Sigma_\tau} d\omega \wedge \text{Tr}(\phi g \bar{\partial} g)
\end{aligned} \tag{2.46}$$

preserving the holomorphic symplectic form Ω on $T^*\widehat{\mathbf{g}}^{\Sigma_\tau}$

$$\Omega = \delta c \wedge \delta \kappa + \int_{\Sigma_\tau} d\omega \text{Tr}(\delta \phi \wedge \delta \bar{A}) \tag{2.47}$$

Then, the moment map has the form

$$\mu = \kappa \bar{\partial} \phi + [\bar{A}, \phi] \tag{2.48}$$

In order to get the system of interacting particles, one is to enlarge the phase space by adding a finite dimensional coadjoint orbit of $SL(N, \mathbb{C})$. A generic complex orbit $SL(N, \mathbb{C}) / (SL(N-l, \mathbb{C}) \times (\mathbb{C}^*)^l)$ is given by a matrix f_{ij} of rank l which can be presented as

$$f_{ij} = \sum_{a=1}^l u_i^a v_j^a \tag{2.49}$$

Then, the moment map M on the enlarged manifold X_ν has the zero level and reads as

$$0 = \kappa \bar{\partial} \phi + [\bar{A}, \phi] - J_{ij} \delta(z, \bar{z}) dz \wedge d\bar{z} \omega. \tag{2.50}$$

where $J_{ij} \equiv (\delta_{ij} - 1)f_{ij}$. Using gauge transformations, one can obtain for the matrix elements of ϕ

$$\kappa \bar{\partial} \phi_{ij} + q_{ij} \phi_{ij} = J_{ij} \delta(z, \bar{z}), \tag{2.51}$$

where $a_{ij} \equiv a_i - a_j$, and $a_i \in \mathbb{C}$ are diagonal elements of $\bar{A}$. From this equation one obtains three consequences:

- $J_{ii} = 0$
- $\phi_{ii} = p_i = \text{const}, i = 1, \dots, N; \sum_i p_i = 0$
- $\phi_{ij} = \frac{1}{\kappa} J_{ij} \exp\left(\frac{a_{ij}(z - \bar{z})}{\kappa}\right) \frac{\theta_*\left(z + \frac{\pi a_{ij}}{\kappa \tau_2}\right)}{\theta_*(z) \theta_*\left(\frac{\pi a_{ij}}{\kappa \tau_2}\right)}$, for $i \neq j$, where $\tau_2 = \text{Im} \tau$, θ_* is the θ -function with odd characteristics¹².

The first two statements merely follow from the diagonal components of equation (2.51), with p_i being just arbitrary constants of integration (restricted by $\sum_i p_i = 0$, since we consider the $SL(N, \mathbb{C})$ group). To prove the third statement, let us make the substitution

$$\phi_{ij}(z, \bar{z}) = \exp\left(\frac{a_{ij}(z - \bar{z})}{\kappa}\right) \psi_{ij}(z, \bar{z}), \tag{2.52}$$

so that ψ_{ij} has the nontrivial monodromy

¹²We also use the notation θ_1 for this θ -function.

$$\begin{aligned}\psi_{ij}(z+1, \bar{z}+1) &= \psi_{ij}(z, \bar{z}) \\ \psi_{ij}(z+\tau, \bar{z}+\bar{\tau}) &= e^{-\frac{2\pi i \tau_2}{\kappa} a_{ij}} \psi_{ij}(z, \bar{z})\end{aligned}\tag{2.53}$$

According to (2.51), we have $\bar{\partial}\psi_{ij} = J_{ij}\delta(z, \bar{z})$, i.e. ψ_{ij} is a meromorphic section of the holomorphic linear bundle with the pole at the origin.

The matrix ϕ obtained is nothing but the Lax operator for the elliptic spin Calogero system [71, 72]. In order to get spinless Calogero case, one needs to choose the simplest orbit $l = 1$ (which describes the fundamental representation). In this case, the factor matrix J_{ij} in ϕ can be trivially gauged away to an arbitrary scalar $-\kappa\nu$. One should also introduce the new rescaled variables $q_i \equiv -\pi a_i/\kappa\tau_2$ identified with coordinates in the many-body system, p_i being the corresponding momenta, and one obtains finally the Lax operator for the elliptic Calogero system which being calculated at $z \neq 0$, coincides with the Lax matrix found by Krichever earlier [73].

Now invariants of the algebra yield Hamiltonians of the integrable system. The simplest one coincides, up to a constant, with the second Hamiltonian of the elliptic Calogero-Moser system

$$\begin{aligned}H &= \int_{\Sigma_\tau} d^2z \frac{1}{2} \text{Tr} \phi(z, \bar{z})^2 = \int_{\Sigma_\tau} d^2z \left\{ \sum_i \frac{1}{2} p_i^2 + \nu^2 \left[\sum_{i < j} \wp(q_{ij}) - \wp(z) \right] \right\} = \\ &= \left\{ \int_{\Sigma_\tau} d^2z \right\} \times \left\{ \sum_i \frac{1}{2} p_i^2 + \nu^2 \sum_{i < j} \wp(q_{ij}) \right\} - \frac{\nu^2 N(N-1)}{2} \int_{\Sigma_\tau} d^2z \wp(z)\end{aligned}\tag{2.54}$$

Let us remark that there is an essential difference between elliptic model and its degeneration at the quantum level. It appears that it is impossible to get the quantum spectrum of the elliptic model analytically. The only tool available is the Bethe ansatz approach which yields the equation for the eigenvalues [74]. Some quantum aspects of these systems within the representation theory framework were discussed in [75].

One may want to obtain an elliptic deformation of the two-dimensional YM theory. The action proposed in [70] is of the form

$$S_\tau = \int_{\Sigma_\tau \times S^1} \omega \wedge \text{Tr}(\phi F_{t\bar{z}} - \varepsilon \phi^2)\tag{2.55}$$

where S^1 is a time-like circle. One should certainly also insert a (vertical) Wilson line $P \exp(\nu \int A_t)$ in representation R_ν .

Let us emphasize that the elliptic Calogero-Moser model is an example of the Hitchin systems given on the moduli space of the holomorphic vector bundles [76, 77, 78, 79] for the surfaces with marked points. The elliptic Calogero-Moser system has a natural relativization – the elliptic Ruijsenaars-Schneider model which is generalization of the Hitchin systems. A description of the elliptic Ruijsenaars-Schneider system via Hamiltonian and Poissonian reductions can be found in [80].

Recently the Hamiltonian reduction procedure for the elliptic Calogero models considered above for A_n case has been extended to other classical groups in [81].

3 Particle systems and supersymmetric gauge theories

After having presented a manifest construction of the integrable systems, we are ready to show how these integrable many-body systems emerge in the consideration of effective low-energy $\mathcal{N} = 2$ su-

persymmetric gauge field theories. As it will be clear later, the role of the many-body systems is to provide the proper degrees of freedom in the vacuum sector of the theory. The fact that the complete set of data fixing the low-energy sector of the theory is governed by the many-body system definitely implies that the number of the effective degrees of freedom in the vacuum sector of the supersymmetric gauge theory is finite. In this section, we demonstrate how the corresponding particle systems can be recovered, while later we comment on the role of these degrees of freedom within the string theory context.

3.1 Seiberg-Witten theory and integrable systems

As explained in the Introduction, the set of data a Riemann surface, the corresponding moduli space and the differential dS that gives the physical quantities, in particular, prepotential is naturally associated with an integrable system. In the case of the $SU(2)$ pure gauge theory, which has been worked out in detail in [1] this correspondence can be explicitly checked [37]. In particular, one may manifestly check formulas (1.3), since all necessary ingredients are obtained in [1]. In this way, one can actually prove that this Seiberg-Witten theory corresponds to the integrable system (periodic Toda chain with two particles, see below). However, to repeat the whole Seiberg-Witten procedure for theories with more vacuum moduli is technically quite tedious if possible at all. Meanwhile, only in this way one could unquestionably prove the correspondence in other cases.

In real situation, the identification between the Seiberg-Witten theory and the integrable system comes via comparing the three characteristics

- number of the vacuum moduli and external parameters
- perturbative prepotentials
- deformations of the two theories

The number of vacuum moduli (i.e. the number of scalar fields that may have non-zero v.e.v.'s) on the physical side should be compared with dimension of the moduli space of the spectral curves in integrable systems, while the external parameters in gauge theories (bare coupling constants, hypermultiplet masses) should be also some external parameters in integrable models (coupling constants, values of Casimir functions for spin chains etc).

As for second item, we already pointed out the distinguished role of the prepotential in Seiberg-Witten theory (which celebrates a lot of essential properties which we shall discuss later). In the prepotential, the contributions of particles and solitons/monopoles (dyons) sharing the same mass scale, are still distinguishable, because of different dependencies on the bare coupling constant, *i.e.* on the modulus τ of the bare coordinate elliptic curve (in the UV-finite case) or on the Λ_{QCD} parameter (emerging after dimensional transmutation in UV-infinite cases). In the limit $\tau \rightarrow i\infty$ ($\Lambda_{QCD} \rightarrow 0$), the solitons/monopoles do not contribute and the prepotential reduces to the “perturbative” one, describing the spectrum of non-interacting *particles*. It is immediately given by the SUSY Coleman-Weinberg formula [39]:

$$\mathcal{F}_{pert}(a) \sim \sum_{\text{reps } R, i} (-)^F \text{Tr}_R(a + M_i)^2 \log(a + M_i) \quad (3.1)$$

Seiberg-Witten theory (actually, the identification of appropriate integrable system) can be used to construct the non-perturbative prepotential, describing the mass spectrum of all the “light” (non-

stringy) excitations (including solitons/monopoles). Switching on Whitham times [82] presumably allows one to extract some correlation functions in the “light” sector (see below).

The perturbative prepotential (3.1) has to be, certainly, added with the classical part,

$$\mathcal{F}_{class} = -\tau a^2 \quad (3.2)$$

since the bare action is $S(\Psi) = \text{Im}\tau \int \text{Tr}\Psi^2$. τ here is value of the coupling constant in the deeply UV region. It plays the role of modular parameter of the bare torus in the UV finite theories. On the other hand, in asymptotically free theories, where the perturbative correction behaves like $a^2 \log(a/\Lambda_{QCD})$, τ is just contained in Λ_{QCD} , since can not be invariantly defined neither in integrable nor in gauge theory.

As we already mentioned the problem of calculation of the prepotential in physical theory is simple only at the perturbative level, where it is given just by the leading contribution, since the β -function in $\mathcal{N} = 2$ theories is non-trivial only in one loop. However, the calculation of all higher (instantonic) corrections in the gauge theory can be hardly done at the moment¹³. Therefore, the standard way of doing is to make an identification of the Seiberg-Witten theory and an integrable system and then to rely on integrable calculations. This is why establishing the “gauge theories $\leftrightarrow$ integrable theories” correspondence is of clear practical (apart from theoretical) importance.

Now let us come to the third item in the above correspondence, namely, how one can extend the original Seiberg-Witten theory. There are basically three different ways to extend the original Seiberg-Witten theory .

First of all, one may consider other gauge groups, from other simple classical groups to those being a product of several simple factors. The other possibility is to add some matter hypermultiplets in different representations. The two main cases here are the matter in fundamental or adjoint representations. At last, the third possible direction to deform Seiberg-Witten theory is to consider 5- or 6-dimensional theories, compactified respectively onto the circle of radius R_5 or torus with modulus R_5/R_6 (if the number of dimensions exceeds 6, the gravity becomes obligatory coupled to the gauge theory).

Now we present a table of known relations between gauge theories and integrable systems, Fig.1.

Before coming into details, let us briefly describe different cells of the table.

The original Seiberg-Witten model, which is the $4d$ pure gauge $SU(N)$ theory (in fact, in their papers [1, 2], the authors considered the $SU(2)$ case only, but the generalization made in [36] is quite immediate), is the upper left square of the table. The remaining part of the table contains possible deformations. Here only two of the three possible ways to deform the original Seiberg-Witten model are shown. Otherwise, the table would be three-dimensional. In fact, the third direction related to changing the gauge group, although being of an interest is slightly out of the main line. Therefore, we only make several comments on it.

One direction in the table corresponds to matter hypermultiplets added. The most interesting is to add matter in adjoint or fundamental representations, although attempts to add antisymmetric and symmetric matter hypermultiplets were also done (see [85] for the construction of the curves and [86] for the corresponding integrable systems). Adding matter in other representations in the basic $SU(N)$ case leads to non-asymptotically free theories.

¹³In order to check the very ideology that the integrable systems lead to the correct answers, there were calculated first several corrections [83, 84]. The results proved to exactly coincide with the predictions obtained within the integrable approach.

SUSY physical theory	Pure gauge SYM theory, gauge group G	SYM theory with adj. matter	SYM theory with fund. matter
4d	inhomogeneous periodic Toda chain for the dual affine $\widehat{\mathfrak{g}}^\vee$ (<i>non-periodic Toda chain</i>)	elliptic Calogero model (<i>trigonometric Calogero model</i>)	inhomogeneous periodic XXX spin chain (<i>non-periodic chain</i>)
5d	periodic relativistic Toda chain (<i>non-periodic chain</i>)	elliptic Ruijsenaars model (<i>trigonometric Ruijsenaars</i>)	periodic XXZ spin chain (<i>non-periodic chain</i>)
6d	periodic “Elliptic” Toda chain (<i>non-periodic chain</i>)	Dell system (<i>dual to elliptic Ruijsenaars, elliptic-trig.</i>)	periodic XYZ (elliptic) spin chain (<i>non-periodic chain</i>)

Figure 1: SUSY gauge theories $\iff$ integrable systems correspondence. The perturbative limit is marked by the italic font (in parenthesis).

Columns: Matter in adjoint vs. fundamental representations of the gauge group. Matter in adjoint representation can be described in terms of a larger pure SYM model, either with higher SUSY or in higher dimensional space-time. Thus models with adjoint matter form a hierarchy, naturally associated with the hierarchy of integrable models $Toda\ chain \hookrightarrow Calogero \hookrightarrow Ruijsenaars \hookrightarrow Dell$ [37, 87, 88, 89, 90, 91, 69, 92, 93, 94]. Similarly, the models with fundamental matter [2, 95] are related to the hierarchy of spin chains originated from the Toda chain: $Toda\ chain \hookrightarrow \mathbf{XXX} \hookrightarrow \mathbf{XXZ} \hookrightarrow \mathbf{XYZ}$ [96, 97, 38, 98, 99].

Note that, while coordinates in integrable systems describing pure gauge theories and those with fundamental matter, live on the cylinder (i.e. the dependence on coordinates is trigonometric), the coordinates in the Calogero system (adjoint matter added) live on a torus¹⁴. However, when one takes the perturbative limit, the coordinate dependence becomes trigonometric.

Lines: Gauge theories in different dimensions. Integrable systems relevant for the description of vacua of $d = 4$ and $d = 5$ models are respectively the Calogero and Ruijsenaars ones (which possess the ordinary Toda chain and “relativistic Toda chain” as Inozemtsev’s limits [100]), while $d = 6$ theories are described by the double elliptic (Dell) systems. When we go from $4d$ (Toda, **XXX**, Calogero) theories to (compactified onto circle) $5d$ (relativistic Toda, **XXZ**, Ruijsenaars) theories the momentum-dependence of the Hamiltonians becomes trigonometric (the momenta live on the compactification circle) instead of rational. Similarly, the (compactified onto torus) $6d$ theories

¹⁴Since these theories are UV finite, they depend on an additional (UV-regularizing) parameter, which is exactly the modulus τ of the torus.

give rise to an elliptic momentum-dependence of the Hamiltonians, with momenta living on the compactification torus. Since adding the adjoint hypermultiplet elliptizes the coordinate dependence, the integrable system corresponding to $6d$ theory with adjoint¹⁵ matter celebrates both the coordinate- and momentum-dependencies elliptic. A candidate for “the elliptic Toda chain” was proposed in [101].

Further in this section we show how the Riemann surfaces and the meromorphic differentials arise when considering a concrete integrable many-body system and explain in more details different squares in the table of correspondence between Seiberg-Witten theories and integrable systems. However, of all $6d$ theories we describe here only the **XYZ** chain case, postponing the discussion of the most general adjoint matter system, Dell system (the right bottom cell of the table) till section 7 where this integrable model naturally emerges within the framework of duality.

3.2 $4d$ pure gauge theory: Toda chain

We start with the simplest case of the $4d$ pure gauge theory, which was studied in details in the original paper [1] for the $SU(2)$ gauge group and straightforwardly generalized to the $SU(N_c)$ case in [36].

This system is described by the periodical Toda chain with period N_c [37, 102, 103] whose equations of motion read

$$\frac{\partial q_i}{\partial t} = p_i \quad \frac{\partial p_i}{\partial t} = e^{q_{i+1}-q_i} - e^{q_i-q_{i-1}} \quad (3.3)$$

with the periodic boundary conditions $p_{i+N_c} = p_i$, $q_{i+N_c} = q_i$ imposed. In physical system, there are N_c moduli and there are no external parameters (Λ_{QCD} can be easily removed by rescaling). In the Toda system there are exactly N_c conservation laws. These conservation laws can be constructed from the Lax operator defined for any integrable system. The eigenvalues of the Lax operator do not evolve, thus, any function of the eigenvalues is an integral of motion.

There are two different Lax representation describing the periodic Toda chain. In the first one, the Lax operator is represented by the $N_c \times N_c$ matrix depending on dynamical variables

$$\mathcal{L}^{TC}(w) = \begin{pmatrix} p_1 & e^{\frac{1}{2}(q_2-q_1)} & 0 & we^{\frac{1}{2}(q_1-q_{N_c})} \\ e^{\frac{1}{2}(q_2-q_1)} & p_2 & e^{\frac{1}{2}(q_3-q_2)} & \dots & 0 \\ 0 & e^{\frac{1}{2}(q_3-q_2)} & -p_3 & & 0 \\ & & \dots & & \\ \frac{1}{w}e^{\frac{1}{2}(q_1-q_{N_c})} & 0 & 0 & & p_{N_c} \end{pmatrix} \quad (3.4)$$

The characteristic equation for the Lax matrix

$$\mathcal{P}(\lambda, w) = \det_{N_c \times N_c} (\mathcal{L}^{TC}(w) - \lambda) = 0 \quad (3.5)$$

generates the conservation laws and determines the spectral curve

$$w + \frac{1}{w} = 2P_{N_c}(\lambda) \quad (3.6)$$

¹⁵Let us point out once more that by adding the adjoint matter we always mean soft breaking of higher supersymmetries.

where $P_{N_c}(\lambda)$ is a polynomial of degree N_c whose coefficients are integrals of motion. If one restores the dependence on Λ_{QCD} in this spectral curve, it takes the form

$$w + \frac{\Lambda_{QCD}^{2N_c}}{w} = 2P_{N_c}(\lambda) \quad (3.7)$$

However, we typically re-scale it away, for exception of s.6.3, where we discuss the manifest RG dynamics w.r.t. Λ_{QCD} .

The spectral curve (3.6) is exactly the Riemann surface introduced in the context of $SU(N_c)$ gauge theory. It can be also put into the hyperelliptic form

$$2Y \equiv w - \frac{1}{w}, \quad Y^2 = P_{N_c}^2(\lambda) - 1 \quad (3.8)$$

The integrals of motion parameterize the moduli space of the complex structures of the hyperelliptic surfaces of genus $N_c - 1$, which is the moduli space of vacua in physical theory.

If one considers the case of two particles ($SU(2)$ gauge theory), $P(\lambda) = \lambda^2 - u$, $u = p^2 - \cosh q$, $p = p_1 = -p_2$, $q = q_1 - q_2$. Thus we see that the order parameter of the SUSY theory plays the role of Hamiltonian in integrable system. In the perturbative regime of the gauge theory, one of the exponentials in $\cosh q$ vanishes, and one obtains the non-periodic Toda chain. In the equation (3.6) the perturbative limit implies vanishing the second term in the l.h.s., i.e. the spectral curve becomes the sphere $w = 2P_{N_c}(\lambda)$.

The alternative Lax representation appeals to introducing the local operators at each site of the chain

$$L_i^{TC}(\lambda) = \begin{pmatrix} \lambda - p_i & e^{q_i} \\ -e^{-q_i} & 0 \end{pmatrix}, \quad i = 1, \dots, N_c \quad (3.9)$$

This Lax operator “makes a shift” to the neighbour site so that the auxiliary linear problem is

$$L_i(\lambda)\Psi_i(\lambda) = \Psi_{i+1}(\lambda) \quad (3.10)$$

where $\Psi_i(\lambda)$ is a two-component Baker-Akhiezer function. The periodic boundary conditions are easily formulated in terms of this Baker-Akhiezer function and read as

$$\Psi_{i+N_c}(\lambda) = w\Psi_i(\lambda) \quad (3.11)$$

where w is a free parameter (diagonal matrix). One can also introduce the transfer matrix that shifts i to $i + N_c$

$$T(\lambda) \equiv L_{N_c}(\lambda) \dots L_1(\lambda) \quad (3.12)$$

Then, the boundary conditions imply that $T(\lambda)\Psi_i(\lambda) = w\Psi_i(\lambda)$, i.e.

$$\det_{2 \times 2} (T_{N_c}(\lambda) - w) = 0 \quad (3.13)$$

providing us with the same spectral curve

$$\begin{aligned}
0 &= w^2 - w \operatorname{Tr} T_{N_c}(\lambda) + \det T_{N_c}(\lambda) = w^2 - w \operatorname{Tr} T_{N_c}(\lambda) + 1, \\
\text{i.e.: } \quad \mathcal{P}(\lambda, w) &= \operatorname{Tr} T_{N_c}(\lambda) - w - \frac{1}{w} = 2P_{N_c}(\lambda) - w - \frac{1}{w} = 0
\end{aligned} \tag{3.14}$$

The identification of the two representations can be also done at the level of their linear problems. Indeed, let us consider the Lax operator (3.9) with the linear problem (3.10) and the boundary conditions (3.11). If we parameterize $\Psi_i = \begin{pmatrix} \psi_i \\ \chi_i \end{pmatrix}$, then the linear problem (3.10) can be rewritten as

$$\psi_{i+1} + p_i \psi_i + e^{q_i - q_{i-1}} \psi_{i-1} = \lambda \psi_i, \quad \chi_i = -e^{q_{i-1}} \psi_{i-1} \tag{3.15}$$

and, along with the periodic boundary condition (3.11) reduces to the linear problem $\mathcal{L}(w)\Phi = \lambda\Phi$ for the $N_c \times N_c$ Lax operator (3.4) with the N_c -component Baker-Akhiezer function $\Phi = \{e^{-q_i/2} \psi_i\}$.

After having constructed the Riemann surface and the moduli space describing the $4d$ pure gauge theory, we turn to the third crucial ingredient of Seiberg-Witten theory that comes from integrable systems, the generating differential dS . The general construction was explained in the Introduction, this differential is in essence the “shorten” action $p dq$. Indeed, in order to construct action variables, a_i one needs to integrate the differential $\widetilde{dS} = \sum_i p_i dq_i$ over $N_c - 1$ non-contractable cycles in the Liouville torus which is nothing but the level submanifold of the phase space, i.e. the submanifold defined by values of all $N_c - 1$ integrals of motion fixed. On this submanifold, the momenta p_i are functions of the coordinates, q_i . The Liouville torus in Seiberg-Witten theory is just the Jacobian corresponding to the spectral curve (1.6) (or its factor over a finite subgroup).

In the general case of a g -parameter family of complex curves (Riemann surfaces) of genus g , the Seiberg-Witten differential dS is characterized by the property $\delta dS = \sum_{i=1}^g \delta u_i dv_i$, where $dv_i(z)$ are g holomorphic 1-differentials on the curves (on the fibers), while δu_i are variations of g moduli (along the base). In the associated integrable system, u_i are integrals of motion and π_i , some g points on the curve are momenta. The symplectic structure is

$$\sum_{i=1}^g da_i \wedge dp_i^{Jac} = \sum_{i,k=1}^g du_i \wedge dv_i(\pi_k) \tag{3.16}$$

The vector of the angle variables,

$$p_i^{Jac} = \sum_{k=1}^g \int^{\pi_k} d\omega_i \tag{3.17}$$

is a point of the Jacobian, and the Jacobi map identifies this with the g -th power of the curve, $Jac \cong \mathcal{C}^{\otimes g}$. Here $d\omega_i$ are *canonical* holomorphic differentials, $dv_i = \sum_{j=1}^g d\omega_j \oint_{A_j} dv_i$. Some details on the symplectic form on the finite-gap solutions can be found in [3, 104].

Technical calculation is, however, quite tedious. It is simple only in the 2-particle ($SU(2)$) case, when the Jacobian coincides with the curve itself. In this case, the spectral curve is

$$w + \frac{1}{w} = 2(\lambda^2 - u) \tag{3.18}$$

while $u = p^2 - \cosh q$. Therefore, one can write for the action variable

$$a = \oint pdq = \oint \sqrt{u - \cosh q} dq = \oint \lambda \frac{dw}{w}, \quad dS = \lambda \frac{dw}{w} \quad (3.19)$$

where we made the change of variable $w = e^q$ and used equation (3.18).

Now one can naturally assume that this expression for the differential dS is suitable for generic N_c . A long calculation (which can be borrowed, say, from the book by M.Toda [105]) shows that this is really the case. One can easily check that the derivatives of dS w.r.t. to moduli are holomorphic, up to total derivatives. Say, if one parameterizes¹⁶ $P_{N_c}(\lambda) = -\lambda^{N_c} + s_{N_c-2}\lambda^{N_c-2} + \dots$ and note that $dS = \lambda dw/w = \lambda dP_{N_c}(\lambda)/Y = P_{N_c}(\lambda)d\lambda/Y + \text{total derivatives}$, then

$$\frac{\partial dS}{\partial s_k} = \frac{\lambda^k d\lambda}{Y} \quad (3.20)$$

and these differentials are holomorphic if $k \leq N_c - 2$. Thus, $N_c - 1$ moduli gives rise to $N_c - 1$ holomorphic differentials which perfectly fits the genus of the curve (we use here that there is no modulus s_{N_c-1}).

It turns out that the form of $dS = \lambda dw/w$ is quite general and does not change even in more complicated cases of spin chains, which we discuss in the next subsection.

To conclude this subsection, let us briefly consider the theories with other gauge groups. First of all, we present an invariant algebraic formulation of the Toda Lax operator (3.4). It can be obtained much along the line discussed in the previous section, i.e. via the Hamiltonian reduction [60], or even obtained from the elliptic Calogero model by a degeneration [100] (we discuss it later). Here we just write down the answer which looks like [106, 60]

$$\mathcal{L}(w) = \sum_i^{N_c-1} [e^{\alpha_i \cdot \mathbf{q}} (\mathbf{e}_{\alpha_i} + \mathbf{e}_{\alpha_i}) + p_i \mathbf{h}_i] + w \mathbf{e}_{\alpha_0} e^{\alpha_0 \cdot \mathbf{q}} + \frac{1}{w} \mathbf{e}_{-\alpha_0} e^{\alpha_0 \cdot \mathbf{q}} \quad (3.21)$$

where the sum runs over all simple roots α_i of the affine algebra $A_{N_c-1}^{(1)}$, $\mathbf{e}_{\alpha_0}$ is the highest (long) root, $\mathbf{h}_i$ span the Cartan subalgebra and the vector $\mathbf{q}$ has as its components the particle coordinates. Note that the generators $\mathbf{e}_{\alpha_i}$ in (3.4) are taken in the simplest fundamental representation of the algebra. However, this requirement is not essential for the whole construction and can be removed.

This expression is straightforwardly extended to any simple affine Lie algebra¹⁷, $N_c - 1$ being substituted by the rank of the group. Now, in order to construct Seiberg-Witten theory with a simple gauge group G one needs to consider the Toda chain for the corresponding *dual* affine algebra $\widehat{\mathfrak{g}}^\vee$ [102]. Say, the gauge theory with the group $Sp(2n)$ is described by the affine Toda for $\left(C_n^{(1)}\right)^\vee$ etc.

Having the Lax operator, one may repeat the procedure of this subsection in order to get the spectral curve and the corresponding moduli space. As for the generating differential dS , it preserves the form $dS = \lambda dw/w$ for all groups.

Note that for any, at least, classical simple group there is also a 2×2 Lax representation generalizing (3.9). In this case, the Lax operator is given on the Dynkin diagram as a substitute of the closed chain above and the definition of transfer matrix is to be extended to include the reflection matrices [107]. Therefore, one can immediately construct some Seiberg-Witten theory in this way. We refer for the details to [108].

¹⁶The absence of the term λ^{N_c-1} is due to the $SU(N_c)$ group and corresponds to the total momentum equal to zero, i.e. to the center mass frame. This is important for further counting of holomorphic differentials.

¹⁷Sometimes one needs to add non-unit coefficients, depending on the root, in front of exponentials.

3.3 Adjoint hypermultiplet added: Calogero system

As far as there exist two different Lax representations, there are, at least, two possible generalizations of the periodic Toda chain (in fact, more as we shall see later). The first generalization is related to the $N_c \times N_c$ representation (3.4). On physical side, it corresponds to the simplest deformation of the $\mathcal{N} = 2$ SUSY gauge theory by adding the adjoint hypermultiplet [87, 90, 91, 89]. If the matter is massless, the theory is finite and the coupling constant can be defined at arbitrary scale. We mainly discuss the massive matter and find the corresponding dynamical system with the additional parameter. It appears that the deformation by the massive hypermultiplet can be described in terms of the elliptic Calogero model. The hypermultiplet mass plays the role of the coupling constant in the Calogero model, while the UV value of the field theory coupling constant defines the modulus of the torus where the Lax operator is defined. When one sends the mass of the hypermultiplet to infinity, in field theory the dimensional transmutation procedure is applied, which has its counterpart in the many-body system and is equivalent to the transition back from Calogero to the Toda system.

The matrix $N_c \times N_c$ Lax operator for the Calogero system was found in section 2 and read as

$$\mathcal{L}^{Cal}(\xi) = \left(\sum_i^{N_c} p_i \mathbf{h}_i + \sum_{\alpha \in \Delta} F(\mathbf{q}\alpha|\xi) \mathbf{e}_\alpha \right) = \begin{pmatrix} p_1 & F(q_1 - q_2|\xi) & \dots & F(q_1 - q_{N_c}|\xi) \\ F(q_2 - q_1|\xi) & p_2 & \dots & F(q_2 - q_{N_c}|\xi) \\ & & \dots & \\ F(q_{N_c} - q_1|\xi) & F(q_{N_c} - q_2|\xi) & \dots & p_{N_c} \end{pmatrix} \quad (3.22)$$

where the first sum runs over the Cartan subalgebra of $SU(N_c)$, while the second one runs over all roots (not only over the simple ones, in variance with the Toda case). The special functions that appear above are defined [73]

$$F(q|\xi) = \frac{\sigma(q - \xi)}{\sigma(q)\sigma(\xi)}, \quad (3.23)$$

where $\sigma(\xi)$ denotes the (Weierstrass) σ -function [109]. One identity used throughout the calculations with the Calogero (and Ruijsenaars below) system is

$$\frac{\sigma(u - v)\sigma(u + v)}{\sigma^2(u)\sigma^2(v)} = \wp(v) - \wp(u). \quad (3.24)$$

where the Weierstrass $\wp$ -function is defined as¹⁸

$$\wp(\xi) \equiv \partial^2 \log \sigma(\xi) = \sum_m \frac{1}{\sinh^2(\xi + m\tau)} \quad (3.25)$$

This Lax operator (3.22) can be just read off from the Calogero Lax operator obtained in s.2.3 (after a trivial gauge transformation). The only external parameter of the integrable system in this case, the coupling constant ν is proportional to the only external parameter of the physical theory, the hypermultiplet mass M^2 .

¹⁸In accordance with the standard definition, [109] $\wp(\xi)$ is the doubly periodic Weierstrass function with periods 2ω and $2\omega'$, and $\tau = \frac{\omega'}{\omega}$. We remark, however, that although the Weierstrass function depends on two periods, the homogeneity relation

$$\wp(tz|t\omega, t\omega') = t^{-2} \wp(z|\omega, \omega')$$

enable us to arbitrary scale one of these. All our results here are ultimately independent of such scaling and this allows us to choose the real period to be π (that is $\omega = \pi/2$). We make this choice throughout.

The spectral curve Σ^{Cal}

$$\det_{N_c \times N_c} (\mathcal{L}^{Cal}(\xi) - \lambda) = 0 \quad (3.26)$$

covers the “bare” spectral curve, torus $E(\tau)$

$$y^2 = (x - e_1)(x - e_2)(x - e_3) \quad (3.27)$$

manifestly parametrized by the variable ξ : $x = \wp(\xi)$, $y = -\frac{1}{2}\wp'(\xi)$.

Σ^{Cal} can be presented as a polynomial $\mathcal{P}(x, y; \lambda) = 0$ of degree N_c in λ , the manifest (quite involved) expression for it can be found in [87, 90, 110]. However, this manifest expression can be always obtained by a degeneration from the Ruijsenaars spectral curve discussed later. Amazingly, this latter curve can be written in a quite compact form.

In the simplest case of two particles, the curve Σ^{Cal} is

$$\mathcal{P}(x, y; \lambda) = \lambda^2 - u + \nu^2 \wp(\xi) = 0 \quad (3.28)$$

The generating differential dS on the Calogero curves has the form

$$dS = \lambda d\omega, \quad d\omega \equiv d\xi = -2 \frac{dx}{y} \quad (3.29)$$

Here $d\omega$ is the holomorphic 1-differential on the bare spectral torus and one can check that the derivatives w.r.t. moduli of this differential are holomorphic on Σ^{Cal} .

Now let us make the dimensional transmutation, i.e. reproduce the periodic Toda chain from the Calogero system [100, 89, 90, 91]. To this end, first, we degenerate the bare spectral torus $\tau \rightarrow i\infty$. Then, to provide the nearest neighbor interaction, we introduce the homogeneous coordinate “lattice” with the large distance Δ between sites

$$q_j = j\Delta + \phi_j \quad (3.30)$$

To see explicitly what kind of interaction emerges in this limit, it suffices to look at the Weierstrass function giving the Calogero potential, although the procedure can be easily repeated for the Lax operator. The Calogero potential has form (see (2.54))

$$V(x_{ij}) = \nu^2 \sum_{i,j} \wp(q_{ij}) \quad (3.31)$$

In the limit under consideration we introduce the renormalized coupling constant $\nu = \nu_0 \exp(\Delta)$ so that νu_0 will be ultimately nothing but Λ_{QCD} . Now, choosing $\Delta \sim \tau$ and taking the limit $\tau \rightarrow i\infty$, we see that only $m = 0$ term survives in the sum (3.25) in the potential (3.31) so that the resulting potential reads as

$$V_0(\phi_i) = \nu_0^2 \sum_{j=1}^{N_c-1} e^{\phi_{i+1} - \phi_i} \quad (3.32)$$

and describes the open (non-periodic) Toda chain.

In order to get the periodic Toda chain, one needs to fix $\Delta = \frac{\tau}{n}$. In this case, the $m = -1$ term in (3.25) also contributes into the sum and one finally obtains the potential

$$V_{TC}(\phi_i) = V_0 + \nu_0^2 e^{\phi_1 - \phi_n} \quad (3.33)$$

describing the periodic Toda chain.

Now, the bare torus (3.27) under the procedure described degenerates into a sphere with two punctures, at $w = 0$ and $w = \infty$: $y \rightarrow w - 1/w$, $x \rightarrow w + 1/w$, while the generating differential dS turns into $\lambda dw/w$ ($d\omega \rightarrow dw/w$) coinciding with (3.19).

Returning to physical theory, making the limit to the pure gauge theory one needs to send the hypermultiplet mass m to infinity keeping finite $\Lambda^{N_c} \sim M^{N_c} e^{2\pi i \tau}$. This is exactly the limit we just discussed in integrable system.

To conclude our consideration, let us note that the Calogero Lax operator considered above can be generalized to other groups [111], and the corresponding Seiberg-Witten theory can be constructed [112]. However, since the exact construction requires entering many technicalities we skip it here.

3.4 Fundamental matter added (SQCD): XXX spin chain

The second possible generalization is related to the 2×2 representation. This generalization on physical side corresponds considering theories with massive hypermultiplets in the fundamental representation. To preserve the asymptotic freedom one has to demand for the number of matter hypermultiplets, N_f not exceed $2N_c$. In particular, the theory with zero β -function, i.e. UV finite one corresponds to $N_f = 2N_c$. On integrable side, this theory is described by the inhomogeneous twisted **XXX** spin chain, while theories with $N_f < 2N_c$ are described by its degenerations [96, 38].

The periodic inhomogeneous $sl(2)$ **XXX** chain of length N_c is given by the 2×2 Lax matrices

$$L_i(\lambda) = (\lambda + \lambda_i) \cdot \mathbf{1} + \sum_{a=1}^3 S_{a,i} \cdot \sigma^a \quad (3.34)$$

σ^a being the standard Pauli matrices and λ_i being the chain inhomogeneities, and periodic boundary conditions are imposed. The linear problem in the spin chain has the same form (3.10) as in the Toda case.

One can also introduce the transfer matrix as in (3.12) which provides the spectral curve equation (3.13) and generates a complete set of integrals of motion.

Integrability of the spin chain follows from the *quadratic* r-matrix relations (see, e.g. [113])

$$\{L_i(\lambda) \otimes L_j(\lambda')\} = \delta_{ij} [r(\lambda - \lambda'), L_i(\lambda) \otimes L_i(\lambda')] \quad (3.35)$$

with the rational r -matrix

$$r(\lambda) = \frac{1}{\lambda} \sum_{a=1}^3 \sigma^a \otimes \sigma^a \quad (3.36)$$

The crucial property of this relation is that it is multiplicative and any product like (3.12) satisfies the same relation

$$\{T(\lambda) \otimes T(\lambda')\} = [r(\lambda - \lambda'), T(\lambda) \otimes T(\lambda')] \quad (3.37)$$

The Poisson brackets of the dynamical variables S_a , $a = 1, 2, 3$ (taking values in the algebra of functions) are fixed by (3.35) and are just

$$\{S_a, S_b\} = -i\epsilon_{abc}S_c \quad (3.38)$$

i.e. the vector $\{S_a\}$ plays the role of angular momentum (“classical spin”) giving the name “spin-chains” to the whole class of systems. Algebra (3.38) has an obvious Casimir function (an invariant, which Poisson commutes with all the spins S_a),

$$K^2 = \mathbf{S}^2 = \sum_{a=1}^3 S_a S_a \quad (3.39)$$

The spectral curve (3.13) is explicitly now

$$w + \frac{Q_{N_f}(\lambda)}{w} = 2P_{N_c}(\lambda), \quad 2P_{N_c}(\lambda) \equiv \text{Tr}T(\lambda), \quad Q_{N_f}(\lambda) \equiv \det T(\lambda) \quad (3.40)$$

or in the hyperelliptic parameterization

$$Y^2 = P_{N_c}^2(\lambda) - Q_{N_f}(\lambda) \quad (3.41)$$

Zeros of $Q_{N_f}(\lambda)$ define the masses of the hypermultiplets. Since

$$\det_{2 \times 2} L_i(\lambda) = (\lambda + \lambda_i)^2 - K_i^2 \quad (3.42)$$

one gets

$$\begin{aligned} Q_{N_f}(\lambda) &= \det_{2 \times 2} T(\lambda) = \prod_{i=1}^{N_c} \det_{2 \times 2} L_i(\lambda) = \prod_{i=1}^{N_c} ((\lambda + \lambda_i)^2 - K_i^2) = \\ &= \prod_{i=1}^{N_c} (\lambda - m_i^+)(\lambda - m_i^-) \end{aligned} \quad (3.43)$$

where we assumed that the values of spin K can be different at different sites of the chain, and

$$m_i^\pm = -\lambda_i \pm K_i. \quad (3.44)$$

Thus we obtain that N_f is generally equal to $2N_c$. Note that the hypermultiplet masses being external parameters in the gauge theory are also such in the spin chain. Indeed, there are still $N_c - 1$ integrals of motion that parameterize the spectral curve and the moduli space (i.e. parameterize the Coloumb branch of gauge theory), but the system also depends on $2N_c$ additional external parameters, N_c Casimir functions and N_c inhomogeneities.

While the determinant of the monodromy matrix (3.42) depends on dynamical variables only through the Casimirs K_i of the Poisson algebra, the dependence of the trace $\text{Tr}_{2 \times 2} T(\lambda)$ is less trivial. Still, it depends on $S_a^{(i)}$ only through Hamiltonians of the spin chain (which are not Casimirs but Poisson-commute with *each other*) – see further details in [96].

Let us note that we have some additional freedom in the definition of the spin chain and the spectral curve. Namely, note that r -matrix (3.36) is proportional to the permutation operator $P(X \otimes Y) = Y \otimes X$. Therefore, it commutes with any matrix of the form $U \otimes U$. Thus, one can multiply Lax operator of the spin chain by arbitrary constant matrix without changing the commutation relations and conservation laws. Moreover, one can also attach a constant (external magnetic field) matrix V

to the end of the chain (to the N_c -th site). This is the same as to consider more general boundary conditions – those with arbitrary matrix V^{-1} . This is why such a model is called twisted.

The described freedom allows one to fit easily the form of the spectral curve proposed in [2, 114]

$$w + \frac{Q(\lambda)}{w} = P(\lambda),$$

$$P(\lambda) = \prod_{i=1}^{N_c} (\lambda - \phi_i), \quad Q(\lambda) = h(h+1) \prod_{j=1}^{2N_c} \left(\lambda - m_j - \frac{2h}{n} \sum_i m_i \right), \quad (3.45)$$

$$h(\tau) = \frac{\theta_2^4}{\theta_4^4 - \theta_2^4}$$

where τ is the bare curve modular parameter and θ_i are the theta-constants.

It can be done, e.g., by choosing the matrices U and V to be¹⁹

$$U_i = \begin{pmatrix} 1 & 0 \\ 0 & \alpha_i \end{pmatrix}, \quad V = \begin{pmatrix} 1 & h(h+1) \\ \prod_i \alpha_i & 0 \end{pmatrix}, \quad (3.46)$$

$$\text{i.e.} \quad V^{-1} = \frac{1}{\det V} \begin{pmatrix} 0 & -h(h+1) \\ -\prod_i \alpha_i & 1 \end{pmatrix}$$

Now let us briefly consider degenerating our UV finite system to an $N_f < 2N_c$ case. This can be done in the standard way [2] by sending l masses $m_1, \dots, m_l$ to infinity while keeping $\Lambda_{QCD}^l \equiv e^{i\pi\tau} m_1 \dots m_l$ finite. After this procedure the modular forms disappear from (3.45) so that Λ_{QCD} emerges instead.

Degenerations of the system can be studied at a single site (for the sake of brevity, we omit the index of the site). Let us consider the Lax operator $\tilde{L} = UL$ (see (3.34), (3.46)) with spins satisfying the Poisson brackets (3.38). We are going to send α to zero. It still reserves two possibilities to get nontrivial Lax operator. The first possibility, when the both masses (3.44) disappear and one reaches the pure gauge theory, is described by the periodic Toda chain. In order to get it, one needs to redefine $S_+ \rightarrow \frac{1}{\alpha} S_+$, then to send α to zero and to remove, after this, the inhomogeneity by the shift of S_0 . This brings us to the Lax operator of the form (we introduce new notations $S_0 = S_3$, $S_{\pm} = S_1 \pm iS_2$)

$$\begin{pmatrix} \lambda + S_0 & S_- \\ S_+ & 0 \end{pmatrix} \quad (3.47)$$

so that the Poisson brackets are

$$\{S_{\pm}, S_0\} = \pm S_{\pm}, \quad \{S_+, S_-\} = 0 \quad (3.48)$$

This algebra is realized in new (Heisenberg) variables p and q

$$S_{\pm} = \pm e^{\pm q}, \quad S_0 = -p, \quad \{p, q\} = 1 \quad (3.49)$$

This leads us finally to the Toda chain Lax operator (3.9) and the Toda spectral curve (3.14).

¹⁹To fit (3.45), we also need to shift $\lambda_i \rightarrow \lambda_i - \frac{2h}{n} \sum_i m_i$.

Now let us return to the second possibility of the asymmetric degeneration, when one of the masses (3.44) remains in the spectrum while the second one goes to infinity. One can understand from (3.44) that, in contrast to the Toda case, this degeneration requires a special fine tuning of the Casimir function and inhomogeneity, so that both of them go to infinity but their sum (difference) remains finite. In the Lax operator, it can be done in the following way. Let us redefine $S_+ \rightarrow \frac{1}{\alpha}S_+$ and $S_0 \rightarrow \frac{1}{\alpha}S_0$. This means that the Poisson brackets take the form

$$\{S_+, S_-\} = -2S_0, \quad \{S_{\pm}, S_0\} = 0 \quad (3.50)$$

Now in order to preserve the finite Lax operator (3.34), one needs to take care of its element $L_{11}(\lambda)$. This can be done by rescaling $\lambda_i \rightarrow \frac{1}{\alpha}\lambda_i$ and fixing $\lambda_i + S_0$ to be $\alpha \cdot s_0$. This brings us to the Lax operator

$$L(\lambda) = \begin{pmatrix} \lambda + s_0 & S_- \\ S_+ & \lambda_i - S_0 \end{pmatrix} = \begin{pmatrix} \lambda + s_0 & S_- \\ S_+ & -2S_0 \end{pmatrix} \quad (3.51)$$

The determinant of this Lax operator is equal to $(\lambda - m)$ where m is the finite mass

$$m = 2s_0S_0 + S_+S_- \quad (3.52)$$

in perfect agreement with (3.44). Let us note that $2s_0S_0 + S_+S_-$ is also the Casimir function of the algebra (3.50), since $\{S_{\pm}, s_0\} = \pm S_{\pm}$.

Let us note that another (equivalent) way to count all possible degenerations [115] is to consider 2×2 Lax operator of the most general form linear in λ , which satisfies the Poisson brackets (3.35) with the rational r -matrix (3.36) and to determine all Casimir functions with respect to this Poisson brackets. Then, all possible degenerations are determined by vanishing the Casimirs functions.

Let us describe another possible interpretation of formula (3.44). Consider a free particle in $d = 4$ with the dispersion law

$$E^2 = \vec{p}^2 + m^2$$

and note that in the complex momentum space there is the level crossing at $\vec{p} = 0$ in the massless case and on the surface $p^2 = -m^2$ in the massive situation. According to general logic of the Berry phase phenomenon, if one considers the momentum space as a parameter space each level crossing point is associated with monopoles and each level crossing surface with “defects” of higher dimensions.

Therefore, the masses define the locations of singularities in the momentum space, indeed. In section 8, we shall see that at the points of the space of scalar field zero modes with coordinates equal to masses, there are branes of different dimensions located. This fits the above arguments, since, due to the gauge invariance, momenta and gauge fields enter the equations of motion combined in covariant derivatives.

The monopole singularity at $\vec{p} = 0$ in the massless case results in an additional term in the rotation operator in the momentum space, the phenomenon well-known for the rotation operator in the monopole background. Moreover, the monopole contribution to the rotation operator allows one to define a chirality transformation for the spin 0, 1, 2 particles in free theory as the rotation when carrying over closed contour in the momentum space. In the interacting theory, the chirality transformation is generically anomalous and, therefore, such a simple reasoning fails. However, the anomaly itself can be described in terms of the Berry phase framework in the space of fields both for

the external and internal anomalies. Then, one has again a satisfactory qualitative agreement, since the chiral invariance is broken either by mass or by anomaly: in both cases the violation is related with the appearance of singularities in the “generalized momentum space”. An analogy with the Peierls model discussed in this context in [116] provides an additional support for this point of view.

To conclude this long subsection, let us mention two other deformations of the original gauge theory. First of all, as in all other cases, one may consider the theory with a different gauge group G . This case is treated similarly to the Toda case by introducing reflection matrices and considering the spin chain on the Dynkin diagram of $\widehat{\mathfrak{g}} \vee$ [108]. Then, the spectral curves can be described by the general formula

$$\mathcal{P}(\lambda, w) = 2P(\lambda) - w - \frac{Q(\lambda)}{w} \quad (3.53)$$

Here $P(\lambda)$ is the characteristic polynomial of the group G for all $G \neq C_n$, i.e.

$$P(\lambda) = \det(g - \lambda I) = \prod_i (\lambda - \lambda_i) \quad (3.54)$$

where determinant is taken in the first fundamental representation and λ_i 's are the eigenvalues of the algebraic element g . For the pure gauge theories with the classical groups [102], $Q(\lambda) = \lambda^{2s}$ and²⁰

$$\begin{aligned} A_{n-1} : \quad P(\lambda) &= \prod_{i=1}^n (\lambda - \lambda_i), \quad s = 0; \\ B_n : \quad P(\lambda) &= \lambda \prod_{i=1}^n (\lambda^2 - \lambda_i^2), \quad s = 2; \\ C_n : \quad P(\lambda) &= \prod_{i=1}^n (\lambda^2 - \lambda_i^2) - \frac{2}{\lambda^2}, \quad s = -2; \\ D_n : \quad P(\lambda) &= \prod_{i=1}^n (\lambda^2 - \lambda_i^2), \quad s = 2 \end{aligned} \quad (3.55)$$

For exceptional groups, the curves arising from the characteristic polynomials of the dual affine algebras do not acquire the hyperelliptic form.

In order to include N_F massive hypermultiplets in the first fundamental representation one can just change λ^{2s} for $Q(\lambda) = \lambda^{2s} \prod_{l=1}^{N_F} (\lambda - m_l)$ if $G = A_n$ and for $Q(\lambda) = \lambda^{2s} \prod_{l=1}^{N_F} (\lambda^2 - m_l^2)$ if $G = B_n, C_n, D_n$ [114, 117, 108].

Another possibility of deformation, letting the gauge groups be products of simple factors and bi-fundamental matter be added, was proposed in [51]. This system is described by a higher, $sl(p)$ **XXX** magnet [38], with the Lax operator given by the $p \times p$ matrix at each site

$$L_i(\lambda) = K^{ab} S_{a,i} X_a + (\lambda + \lambda_i) \cdot \mathbf{1} \quad (3.56)$$

where $S_{a,i}$ are dynamical variables, X_a are generators of the $sl(p)$ algebra and K^{ab} is its Killing form.

Thus, we described the families of spectral curves for theories with fundamental matter. Note that, at any **XXX** spin chain, the generating differential has the same form as in the Toda theory (3.19), $dS = \lambda dw/w$.

²⁰In the symplectic case, the curve can be easily recast in the form with polynomial $P(\lambda)$ and $s = 0$.

3.5 5d SQCD: twisted XXZ chain

Now we describe the integrable system behind the 5d theory with fundamental matter hypermultiplets. The theory is considered with one dimension compactified onto the circle of radius R_5 , i.e. given on the space $\mathbf{R}^4 \times S^1$. We start from the UV finite theory, i.e. that with $N_f = 2N_c$ hypermultiplets. The corresponding integrable theory is the inhomogeneous **XXZ** spin chain [98, 99].

The Lax matrix for the **XXZ** ($sl(2)$) spin magnet has the form

$$L(\mu) = \begin{pmatrix} \mu e^{S_0} - \mu^{-1} e^{-S_0} & 2S_- \\ 2S_+ & \mu e^{-S_0} - \mu^{-1} e^{S_0} \end{pmatrix} \quad (3.57)$$

Thus, this Lax operator is given on a cylinder. In fact, from the point of view of integrable systems, some more natural spectral parameter is ζ : $\mu = e^\zeta$, and the Lax operator (3.57) becomes clearly trigonometric. This Lax operator is intertwined by the trigonometric r -matrix

$$r(\zeta) = \frac{i}{\sinh \pi \zeta} (\sigma_1 \otimes \sigma_1 + \sigma_2 \otimes \sigma_2 + \cosh \pi \zeta \sigma_3 \otimes \sigma_3) \quad (3.58)$$

so that the Poisson bracket of the Lax operators (3.35) gives rise to the Poisson brackets of the algebra of S_i 's:

$$\{S_+, S_0\} = \pm S_\pm; \quad \{S_+, S_-\} = \sinh 2S_0 \quad (3.59)$$

The second Casimir function of this algebra is

$$C_2 = \cosh 2S_0 + 2S_+ S_- \quad (3.60)$$

The non-linear commutation relations (3.59) are ones from the quantum deformed algebra $U_q(sl(2))$. This generalizes the fact that the **XXX** magnet is described by the Poisson brackets that reproduce the classical $sl(2)$ algebra. In these Poisson brackets, the Plank constant $\hbar$ turns out to be inessential and can be put equal to unity. In fact, it is proportional to the radius R_5 of the space-time circle in the corresponding 5d SUSY theory and can be easily restored with the replace of any generator S_i by $R_5 S_i$. Hereafter, we omit R_5 from all the formulas.

Following the standard procedure, now we consider the chain with N_c sites with the Lax operators (3.57) associated with each site and commuting with each other, introduce the inhomogeneities ζ_i which depend on the site of chain by the replace $\zeta \rightarrow \zeta + \zeta_i$ and impose the periodic boundary conditions. On this chain, the transfer matrix acts and the periodic boundary conditions generate the spectral curve and the conservation laws.

The manifest form of the curve in the **XXZ** case can be derived using explicit expression for the Lax matrix (3.57) and reads²¹

$$w + \frac{Q_{2N_c}(e^{2\zeta})}{e^{2N_c\zeta+2\sum_i \zeta_i} w} = 2e^{-N_c\zeta - \sum_i \zeta_i} P_{N_c}(e^{2\zeta}) = 2e^{-N_c\zeta - \sum_i \zeta_i} \left(e^{2N_c\zeta} + \dots + e^{2\sum_i \zeta_i} \right) \quad (3.61)$$

Changing the variables $e^{2\zeta} = \mu^2 \equiv \lambda$, $w \rightarrow e^{-N_c\zeta - \sum_i \zeta_i} w$, this curve can be recast in the hyperelliptic form in λ variables:

²¹In fact, the coefficient in front of P_{N_c} , instead of 2, is equal to $2 \cosh \prod_i^{N_c} e^{S_{0,i}}$. It is, however, the integral of motion that can be put equal to any number. We always fix $\sum_i^{N_c} S_{0,i}$ to be zero.

$$w + \frac{Q_{2N_c}(\lambda)}{w} = 2P_{N_c}(\lambda), \quad Y^2 = P_{N_c}^2(\lambda) - Q_{2N_c}(\lambda), \quad Y \equiv \frac{1}{2} \left(w - \frac{Q_{2N_c}(\lambda)}{w} \right) \quad (3.62)$$

while in terms of the “true” spectral parameter ζ this curve looks considerably more tricky. However, one can work with the variable λ taking into account that it lives on a cylinder or sphere with two marked points. In equation (3.62), the polynomial

$$Q_{2N_c}(\lambda) = \prod_i (\lambda^2 - 2K_i e^{2\zeta_i} \lambda + e^{4\zeta_i}) \equiv \prod_i \left(\lambda - e^{2m_i^{(+)}} \right) \left(\lambda - e^{2m_i^{(-)}} \right) \quad (3.63)$$

defines the masses $m_i^{(\pm)}$ of matter hypermultiplets (cf. with (3.44))

$$m_i^{(\pm)} = \zeta_i \pm \frac{1}{2} \log \left(K_i + \sqrt{K_i^2 - 1} \right) \quad (3.64)$$

Note that, being written in terms of the variable λ , the curve (3.62) is very similar to the curves arising for $4d$ theories. However, the difference coming from different generating 1-differentials dS turns out to be very crucial (see, for example, the discussion of residue formula and perturbative prepotentials in section 5).

It is easy to compare the curves (3.62) with other curves appeared in literature [118], rewriting them in the form

$$w + \frac{\prod_\alpha \sinh(\zeta - m_\alpha)}{w} = 2 \prod_i \sinh(\zeta - a_i) \quad (3.65)$$

where we rescaled $w \rightarrow e^{N_c \zeta + \sum_i \zeta_i} w$, denoted the roots of the polynomial $P_{N_c}(\lambda)$ (3.61) $\lambda_i \equiv e^{2a_i}$ and made use of formula (3.64) and the manifest form of the leading and the constant terms in this polynomial. Comparing (3.65) with (3.61), one finds, in particular, that

$$\sum a_i = \sum \zeta_i = \frac{1}{2} \sum m_\alpha \quad (3.66)$$

The condition (3.66) is not, however, absolutely necessary in the context of **XXZ** chains and $5d$ theories. It emerges only in the standard **XXZ** chain. In the context of Seiberg-Witten theory one needs rather to consider, similarly to the $4d$ case, the *twisted* **XXZ** model [119, 98, 99]. It is characterized by the Lax operator

$$L_i(\mu) = \begin{pmatrix} \mu e^{S_{0,i} - \zeta_i} - \alpha_i \mu^{-1} e^{-S_{0,i} + \zeta_i} & 2S_{-,i} \\ 2S_{+,i} & \mu e^{-S_{0,i} - \zeta_i} - \beta_i \mu^{-1} e^{S_{0,i} + \zeta_i} \end{pmatrix} \quad (3.67)$$

with generally non-unit constants α_i, β_i . These constants provide an arbitrary coefficient in front of the product in the r.h.s. of (3.65) (which can be also fixed by the integral of motion $\sum_i^{N_c} S_{0,i}$) and, thus, break the condition (3.66). They are also important for a careful treatment of dimensional transmutation (i.e. decreasing the number of massive hypermultiplets), see below.

The maximally degenerated case is related to the pure gauge theory when all the masses become infinite. In this case, one gets the relativistic Toda chain, s.3.6.

There certainly exist all intermediate degenerations when all but N_f masses are brought to infinity. This system, corresponding to the gauge theory with $N_f < 2N_c$ massive hypermultiplets, is described

by the chain with some of the sites degenerated and some of them not. The corresponding spectral curve has the form

$$w + \frac{Q_{N_f}(\lambda)}{w} = 2P_{N_c}(\lambda), \quad Q_{N_f}(\lambda) = \prod_{\alpha}^{N_f} (\lambda - e^{2m_{\alpha}}), \quad P_{N_c}(\lambda) = \prod_i^{N_c} (\lambda - e^{2a_i}) \quad (3.68)$$

or

$$w + \frac{\prod_{\alpha}^{N_f} \sinh(\zeta - m_{\alpha})}{w} = 2e^{(N_c - N_f/2)\zeta + \sum a_i - 1/2 \sum m_{\alpha}} \prod_i^{N_c} \sinh(\zeta - a_i) \quad (3.69)$$

(where we again rescaled $w \rightarrow e^{\frac{1}{2}(N_f\zeta + \sum m_{\alpha})}w$; this curve also coincides with that in [118]). As we already noted, the particular coefficient $e^{\sum a_i - 1/2 \sum m_{\alpha}}$ can be, in principle, replaced by any other number, say put equal to unity, which literally corresponds to [118]. It does not influence the result and we will discuss the reason below.

Note that the form of the spectral curve (3.65) is perfectly designed for taking the $4d$ limit. Indeed, one can restore R_5 -dependence in this formula multiplying each ζ_i and mass parameter by R_5 . In terms of algebra (3.59) it means that each generator has to be multiplied by R_5 and so does the spectral parameter. Then, one immediately reproduces the results of the previous subsection.

As in the $4d$ case, the description of gauge theory in terms of the spin chain allows an extension to the group product case. It is described by higher $sl(p)$ **XXZ** magnets given by the Lax operator

$$L(\mu) = \sum_{i,j} e_{ij} \otimes L_{ij} \quad (3.70)$$

and

$$L_{ii} = \mu e^{S_{0,i}} - \mu^{-1} e^{-S_{0,i}}, \quad L_{ij} = 2S_{ji}, \quad (i \neq j). \quad (3.71)$$

Here $S_{0,i}$ are associated with the vectors ϵ_i , realizing the simple roots $\alpha_i = \epsilon_i - \epsilon_{i+1}$, and e_{ij} has the only non-vanishing matrix element (i, j) . The generators S_i satisfy non-linear Poisson brackets that can be read off from the commutation relations of the quantum algebra $U_q(sl(p))$ like it has been done in (3.59). These Poisson brackets can be certainly obtained from the quadratic relation (3.35) with the trigonometric r -matrix for the $sl(p)$ case [113]. Note that, specially degenerating this system, one can easily reproduce the (p, q) -Web constructed in [120] that corresponds to different toric varieties.

Now let us say some words on the generating differential dS in $5d$ theories with fundamental matter.

The general scheme presented for $4d$ theory can be almost literally transferred to the $5d$ $\mathcal{N} = 2$ SUSY gauge models with one compactified dimension. It can be described through involving trigonometric r -matrices and L -operators instead of rational ones, i.e. coming from Yangian to affine algebras. It means that now it is natural to consider the *both* parameters $\zeta = \frac{1}{2}\log\lambda$ and $\log w$ as coordinates on the cylinder.

Similar arguments imply that, instead of differential $dS = \lambda d\log w$, one now has to consider the differential [67, 39]

$$dS = \zeta \frac{dw}{w} \sim \log\lambda \frac{dw}{w} \quad (3.72)$$

so that, despite the similarity of the $5d$ spectral curves with $4d$ ones, periods of $dS^{(5)}$ are different from those of $dS^{(4)}$. Note that the derivatives of this differential w.r.t. moduli again give holomorphic differentials.

In fact, one can interpret the $5d$ theory as the $4d$ theory with infinite number of (Kaluza-Klein) vector multiplets with masses $M_n = n/R_5$ and infinite number of analogous (Kaluza-Klein) fundamental hypermultiplets. Then, one can equally consider either the just described $5d$ picture, or the $4d$ picture that involves Riemann surface presented as an infinite order covering (see the spectral curve equation in terms of the variable ζ (3.61)) with infinitely many punctures. This latter picture can be effectively encapsulated in the usual hyperelliptic Riemann surface (3.62) with finite number of punctures, but, as a memory of infinitely many multiplets, the spectral parameter λ now lives on the cylinder. Meanwhile, the differential $dS^{(5)}$ now evidently should be of the form (3.72) which “remembers” of its $4d$ origin.

3.6 $5d$ pure gauge theory: relativistic Toda chain

We are going to get the pure gauge theory in $5d$ from the theory with fundamental matter via the dimensional transmutation, sending masses of all the hypermultiplets to infinity. Therefore, we study the maximal degeneration of the **XXZ** spin chain [98, 99]. This degeneration is obtained from the twisted **XXZ** chain introduced above and is described by the relativistic Toda chain [67] (see also [39]). The twisting means just introducing some new parameters into the Lax operator, which are central elements of the Poisson bracket algebra. A special fine tuning of these parameters allows one to match smoothly different regimes and limiting cases of the **XXZ** spin chain.

As it has been explained in [38], there are two equivalent ways to twist the integrable system. One of them, which was applied in the **XXX** case, is to multiply the Lax operator by an arbitrary constant matrix U . This is possible in the **XXX** case, since the corresponding (rational) r -matrix commutes with the tensor product $U \otimes U$. The situation is much more restrictive in the trigonometric case, when the r -matrix commutes only with the matrices U of the very special form (say, it can be proportional to any one of the Pauli matrices). Therefore, in the trigonometric case we apply the second way of doing [119, 115], that is, we consider the Lax matrix of the general form but with some prescribed dependence of matrix elements on the spectral parameter. Then, the Poisson bracket (3.35) dictates Poisson brackets of matrix elements and, in particular, fixes some coefficients to be centers of the Poisson bracket algebra.

More explicitly, we fix the Lax matrix to be of the form

$$L(\mu) = \begin{pmatrix} \mu A^{(+)} + \mu^{-1} A^{(-)} & B \\ C & \mu D^{(+)} + \mu^{-1} D^{(-)} \end{pmatrix} \quad (3.73)$$

and to be intertwined by the same r -matrix (3.58). Then, up to inessential total normalization, one can rewrite the Lax operator in the form

$$L(\mu) = \begin{pmatrix} \mu e^{S_0} - \rho \mu^{-1} e^{-S_0} & 2S_- \\ 2S_+ & \gamma \mu e^{-S_0} - \nu \mu^{-1} e^{S_0} \end{pmatrix} \quad (3.74)$$

where ρ , ν and γ are constants (centers of algebra).

In this case, the commutation relations are slightly changed to

$$\{S_+, S_0\} = \pm S_{\pm}; \quad \{S_+, S_-\} = \frac{1}{2} (\nu e^{2S_0} - \rho \gamma e^{-2S_0}) \quad (3.75)$$

and the second Casimir function is equal to

$$C_2 = \frac{1}{2} (\rho \gamma e^{-2S_0} + \nu e^{2S_0}) + 2S_+ S_- \quad (3.76)$$

thus, the determinant of the Lax operator (3.74) is still the quadratic polynomial with coefficient being the Casimir function.

Now we are ready to demonstrate how the limit to the relativistic Toda chain can be done. To make the Lax operator (3.74) looking more similar to the relativistic Toda Lax operator, we multiply it by the function e^{S_0} . In the whole chain it results into the factor $e^{\sum_i^{N_c} S_{0,i}}$ which is the integral of motion and can be put zero. In particular, in the relativistic Toda case this integral is equal to the full momentum of the system. Although being so inessential, this redefining of the Lax operator still requires to modify some expressions [115]. For instance, the new Lax operator

$$L(\mu) = \begin{pmatrix} \mu e^{2S_0} - \rho \mu^{-1} & 2S_- e^{S_0} \\ 2S_+ e^{S_0} & \gamma \mu - \nu \mu^{-1} e^{2S_0} \end{pmatrix} \quad (3.77)$$

is intertwined by the new r -matrix (here r is the r -matrix from (3.58))

$$r^{(tw)} = r + \frac{1}{2} (\mathbf{I} \otimes \sigma_3 - \sigma_3 \otimes \mathbf{I}) \quad (3.78)$$

which is called twisted. This name comes from the quantum generalization of this matrix which can be obtained from the standard quantum trigonometric R -matrix by twisting [115]

$$R^{(tw)}(u) = F_{12}(\gamma) R(u) F_{21}^{-1}(\gamma)' \quad F_{12} \equiv F_{21}^{-1} = \exp \left\{ \frac{1}{2} (\sigma_3 \otimes \mathbf{I} - \mathbf{I} \otimes \sigma_3) \right\} \quad (3.79)$$

An important property of the Lax operator (3.77) is that its determinant is no longer dependent on only the Casimir function but also on e^{S_0} . The product of this quantity over all the sites is still certainly an integral of motion.

Now one can consider the particular reduction [115]

$$\gamma = \nu = 0, \quad \rho = 1 \quad (3.80)$$

i.e.

$$L(\mu) = \begin{pmatrix} \mu e^{2S_0} - \mu^{-1} & 2S_- e^{S_0} \\ 2S_+ e^{S_0} & 0 \end{pmatrix} \quad (3.81)$$

Taking into account the commutation relations (3.75), that is, $\{S_+, S_-\} = 0$, one can realize the algebra as $S_{\pm} = e^{\pm q}$, $S_0 = p$, $\{p, q\} = 1$ and reproduce the standard Lax operator for the relativistic Toda chain [121].

The spectral curve corresponding to this degenerated case can be easily written

$$w + \frac{1}{w} = \frac{1}{\lambda^{N_c/2}} P_{N_c}(\lambda) \quad (3.82)$$

Now one can also obtain from the theory with $N_f = 2N_c$ the theory with less number of massive multiplets degenerating the general **XXZ** chain at several sites. The curves for such theories we already discussed in the previous subsection.

At last, let us note that the differential dS constructed for the **XXZ** spin chain (3.72) does not change when degenerating to the relativistic Toda chain.

3.7 5d adjoint matter: Ruijsenaars-Schneider system

Now we are going to study the system emerging upon adding in 5d theory the adjoint matter hypermultiplet. This UV finite system can be described differently as follows. By starting with a five dimensional model one may obtain four dimensional $\mathcal{N} = 2$ SUSY models (with fields only in the adjoint representation of the gauge group) by imposing non-trivial boundary conditions on half of the fields:

$$\phi(x_5 + R_5) = e^{2i\epsilon} \phi(x_5). \quad (3.83)$$

If $\epsilon = 0$ one obtains $\mathcal{N} = 4$ SUSY in four dimensions, but when $\epsilon \neq 2\pi n$ this is explicitly broken to $\mathcal{N} = 2$. The low-energy mass spectrum of the four dimensional theory contains two towers of Kaluza-Klein modes:

$$M_n = \frac{\pi n}{R_5} \quad \text{and} \quad M_n = \frac{\epsilon + \pi n}{R_5}, \quad n \in Z. \quad (3.84)$$

According to the proposal of [67], this five dimensional theory may be associated with the elliptic Ruijsenaars-Schneider integrable model [61, 62]. In various double-scaling limits it reduces to systems we already studied:

(a) If $R_5 \rightarrow 0$ (with finite ϵ) the (finite) mass spectrum (3.84) reduces to a single point $M_n = 0$. This is the standard four dimensional $\mathcal{N} = 2$ SUSY pure gauge model associated with the periodic Toda chain. As we already know, in this situation $\mathcal{N} = 2$ SUSY in four dimensions is insufficient to ensure UV-finiteness, thus $\tau \rightarrow i\infty$, but the phenomenon of dimensional transmutation occurs whereupon one substitutes the dimensionless τ by the new dimensional parameter $\Lambda_{QCD}^{N_c} = e^{2\pi i\tau} (\epsilon/R_5)^{N_c}$.

(b) If $R_5 \rightarrow 0$ and $\epsilon \sim MR_5$ for finite M , then UV finiteness is preserved. The mass spectrum (3.84) reduces to the two points $M_n = 0$ and $M_n = M$. This is the four dimensional YM model with $\mathcal{N} = 4$ SUSY softly broken to $\mathcal{N} = 2$. The associated finite-dimensional integrable system is, as we know, the elliptic Calogero model. Case (a) is then obtained from (b) by the double scaling limit.

(c) If $R \neq 0$ but $\epsilon \rightarrow i\infty$ the mass spectrum (3.84) reduces to a single Kaluza-Klein tower, $M_n = \pi n/R_5$, $n \in Z$. This compactification of the five dimensional model has $\mathcal{N} = 1$ SUSY and is not UV-finite. Here $\tau \rightarrow i\infty$ and $\epsilon \rightarrow i\infty$, such that $2\pi\tau - N_c\epsilon$ remains finite. This system as we discussed in the previous subsection is described by the relativistic Toda chain.

(d) Finally, when $R_5 \neq 0$ and ϵ and τ are both finite one distinguished case still remains: $\epsilon = \pi/2$.²² Here only periodic and anti-periodic boundary conditions occur in the compact dimension. This is the case analyzed in [122]. It is clearly special from both the point of view of Yang-Mills theory and integrable systems.

²²The case $\epsilon = 0$ of fully unbroken five dimensional $\mathcal{N} = 2$ supersymmetry is of course also distinguished, but trivial: there is no evolution of effective couplings (renormalisation group flows) and the integrable system is just that of N_c non-interacting (free) particles.

Thus, the elliptic Ruijsenaars-Schneider model is a remarkable completely integrable system whose various limits include the (finite dimensional) Toda and Calogero-Moser models. Here we shall review only a few of its salient features. More comprehensive accounts of its structure and applications may be found in [123, 124]. The $SU(N_c)$ model has an explicit Lax representation with Lax operator [61, 62, 125, 126]

$$\mathcal{L}_{ij} = c(\xi|\epsilon) e^{P_i} \frac{F(q_{ij}|\xi)}{F(q_{ij}|\epsilon)}. \quad (3.85)$$

Here

$$e^{P_i} = e^{p_i} \prod_{l \neq i} \sqrt{\wp(q_{il}) - \wp(\epsilon)}. \quad (3.86)$$

where the p_i and q_i are canonically conjugate momenta and coordinates, $\{p_i, q_j\} = \delta_{ij}$. The commuting Hamiltonians may be variously written

$$H_k = \sum_{\substack{J \subset \{1, \dots, n\} \\ |J|=k}} e^{\sum_{j \in J} p_j} \prod_{\substack{j \in J \\ k \in \{1, \dots, n\} \setminus J}} \frac{F(q_{ij}|\xi)}{F(q_{ij}|\epsilon)} = \sum_{1 \leq i_1 < \dots < i_k \leq N_c} e^{P_{i_1} + \dots + P_{i_k}} \prod_{a < b} \frac{1}{\wp(q_{i_a i_b}) - \wp(\epsilon)}. \quad (3.87)$$

(The final product is taken to be unity in the case $k = 1$.) These Hamiltonians arise in the description of the spectral curve. Note that, after trigonometric degeneration, these Hamiltonians are proportional to those obtained in s.2.2. We discuss this degeneration in more details in section 5.

The integrability of the model depends on the functional equations satisfied by F [61, 62]; the connection with functional equations and integrable systems is part of a larger story [127].

Some comment on the parameters ξ and ϵ in these formulae is called for. Here ξ is precisely the spectral parameter associated with the bare spectral curve, where the Lax operator is given. Further, the additional parameter ϵ in (3.85), (3.86) above is the same parameter we introduced in (3.83) characterizing the boundary conditions. Actually the integrability of (3.87) does not require ϵ to be real, but such a choice guarantees the reality of the the Hamiltonians. The identification of these two parameters has been simplified by our choice of the real period of $\wp$ being π so that both (3.83) and (3.85), (3.86) are manifestly π periodic. The “non-relativistic limit” $\epsilon \rightarrow 0$ which leads to the Calogero-Moser system means we can identify the mass $M = \epsilon/R_5$ of the gauge multiplet with the Calogero-Moser coupling constant. The special point $\epsilon = \pi/2$ ($= \omega$) singled out earlier is now a half-period of the $\wp$ -function, and at this point $\wp'(\frac{\pi}{2}) = 0$.

Now inserting the Lax operator (3.85) into the determinant (3.88)

$$\det_{N_c \times N_c} (\mathcal{L}^R(\xi) - \lambda) = 0 \quad (3.88)$$

one gets the spectral curve equation. Its coefficients are expressed through the Hamiltonians (3.87) and parameterize the corresponding moduli space.

To conclude constructing Seiberg-Witten theory we should determine the differential dS . It is evident from our previous discussions that it is to be of the form

$$dS = \frac{1}{R_5} \log \lambda d\xi \quad (3.89)$$

Indeed, on one hand, it should be proportional to $d\xi$ since describes the adjoint matter (compare with the Calogero case). On the other hand, the proportionality coefficient is to be $R_5^{-1}\log\lambda$, since the Ruijsenaars-Schneider system describes a $5d$ theory.

Now let us discuss more the spectral curve arising. From the point of view of classical integrability the overall normalization factor $c(\xi|\epsilon)$ of the Lax operator does not change the integrals of motion apart from scaling. This normalization factor does however lead to the rescaling of λ and this effects the explicit form of the 1-form dS .²³ We shall see below that significant simplifications occur with the choice

$$c(\xi|\epsilon) = c_0(\xi|\epsilon) \equiv \frac{1}{\sqrt{\wp(\xi) - \wp(\epsilon)}} = \frac{\sigma(\xi)\sigma(\epsilon)}{\sqrt{\sigma(\epsilon - \xi)\sigma(\epsilon + \xi)}} = \sqrt{c_+(\xi|\epsilon)c_-(\xi|\epsilon)} \quad (3.90)$$

where

$$c_{\pm}(\xi|\epsilon) \equiv \frac{\sigma(\xi)\sigma(\epsilon)}{\sigma(\epsilon \pm \xi)} = \mp \frac{1}{F(\epsilon|\mp \xi)}. \quad (3.91)$$

Similar issues of normalization enter into discussion of separation of variables [128].

Before generalizing to the higher rank situation it is instructive to first consider the case of gauge group $SU(2)$. In the formulas of the previous section this corresponds to setting $\mathcal{N} = 2$ and working in the center-of-mass frame $p_1 + p_2 = 0$. We define $p \equiv p_1$ and $q \equiv q_1 - q_2$.

Using the explicit form of the Lax operator the spectral curve (3.88) is found to be

$$\lambda^2 - cu\lambda + c^2(\wp(\xi) - \wp(\epsilon)) = \lambda^2 - cu\lambda + \frac{c^2}{c_0^2} = 0, \quad (3.92)$$

where

$$u \equiv H_1(p, q) = 2 \cosh p \sqrt{\wp(q) - \wp(\epsilon)} \quad (3.93)$$

is the Hamiltonian (3.87). For the choice $c(\xi|\mu) = c_0(\xi|\mu)$ this simplifies to yield

$$\lambda^2 - uc_0\lambda + 1 = 0, \quad (3.94)$$

or simply

$$c_0^{-1}(\xi) = \frac{u\lambda}{\lambda^2 + 1} = \frac{u}{\lambda + \lambda^{-1}}. \quad (3.95)$$

Observe that with our choice (3.90) this equation, which describes the Seiberg-Witten spectral curve, may be expressed in the form

$$u = H(\log\lambda, \xi). \quad (3.96)$$

Comparison with (3.89) shows that the generating differential $R_5 dS = \log\lambda d\xi$ takes the form pdq here.

We now consider the general $SU(N_c)$ model. The first step is to evaluate the spectral curve (3.88) for the Lax matrix (3.85). Expanding the determinant about the diagonal yields

²³ Note that although the choice of normalization $c(\xi|\epsilon)$ (or, equivalently, rescaling λ) can change the manifest expression for dS , it affects neither the symplectic 2-form $\frac{d\lambda}{\lambda} \wedge \frac{dw}{w}$, nor the period matrix, which is the second derivative of the prepotential.

$$\sum_{k=0}^{N_c} (-\lambda)^{N_c-k} c^k \left\{ \sum_{1 \leq i_1 < \dots < i_k \leq N_c} e^{P_{i_1} + \dots + P_{i_k}} \det_{(ab)} \frac{F(q_{i_a i_b} | \xi)}{F(q_{i_a i_b} | \epsilon)} \right\} = 0, \quad (3.97)$$

and Ruijsenaars [61, 62] expressed the determinants appearing here in terms of the Hamiltonians (3.87) by means of a generalized Cauchy formula. This expansion was re-obtained in [69] using a simple fact that generalized Cauchy formulae can be derived in terms of free-fermion correlators via Wick's theorem for fermions. The Ruijsenaars-Schneider model (and its spin generalizations) may be understood [126, 129] in terms of a reduction of the Toda lattice hierarchy. The Hirota bilinear identities of that hierarchy may be expressed in terms of free-fermion correlators [130, 20], and are the origin of those here. We believe that the free field expansions will ultimately lead to a better field theoretic understanding of the appearance of these integrable systems. The machinery of free-fermion correlators has already found use, for example, in calculating within the context of the Whitham hierarchy for these integrable systems [82].

The formulas for the determinants are

$$\tilde{D}_{I_k} \equiv \det_{(ab)} \frac{F(q_{i_a i_b} | \xi)}{F(q_{i_a i_b} | \epsilon)} = (-1)^{\frac{k(k-1)}{2}} \frac{\sigma^{k-1}(\xi - \epsilon) \sigma(\xi + (k-1)\epsilon)}{\sigma^k(\xi) \sigma^{k(k-1)}(\epsilon)} \prod_{a < b} \frac{\sigma^2(q_{i_a i_b}) \sigma^2(\epsilon)}{\sigma(\epsilon + q_{i_a i_b}) \sigma(\epsilon - q_{i_a i_b})}, \quad (3.98)$$

where $I_k = \{i_1, \dots, i_k\}$ denotes the set of k indices and $a, b = 1, \dots, k$. The right hand side of this expression is in fact a doubly periodic function of all the arguments (q_i , ξ and ϵ) and so may be rewritten in terms of Weierstrass functions. In particular, upon using (3.24), we find that

$$\prod_{a < b} \frac{\sigma^2(q_{i_a i_b}) \sigma^2(\epsilon)}{\sigma(\epsilon + q_{i_a i_b}) \sigma(\epsilon - q_{i_a i_b})} = \prod_{a < b} \frac{1}{\wp(q_{i_a i_b}) - \wp(\epsilon)}, \quad (3.99)$$

thus recovering the Hamiltonians (3.87) noted earlier.

Now we are ready to obtain the final formula for the spectral curve in the Ruijsenaars-Schneider system. Upon substitution of (3.98) into (3.97) the curve takes the form

$$\sum_{k=0}^{N_c} (-\lambda)^{N_c-k} c^k \left\{ \sum_{I_k} e^{P_{i_1} + \dots + P_{i_k}} \tilde{D}_{I_k} \right\} = \sum_{k=0}^N (-\lambda)^{N_c-k} c^k D_k(\xi | \epsilon) H_k = 0. \quad (3.100)$$

Here we have collected the q -independent factors in (3.98) into D_k where

$$D_k(\xi | \epsilon) = (-1)^{\frac{k(k-1)}{2}} \frac{\sigma^{k-1}(\xi - \epsilon) \sigma(\xi + (k-1)\epsilon)}{\sigma^k(\xi) \sigma^{k(k-1)}(\epsilon)} = \frac{1}{c_-^k} \cdot \frac{\sigma(\xi + (k-1)\epsilon)}{\sigma(\xi - \epsilon)} \cdot \frac{(-1)^{\frac{k(k-1)}{2}}}{\sigma^{k(k-2)}(\epsilon)}. \quad (3.101)$$

Again this is doubly periodic function in both ξ and ϵ and so is expressible in terms of the Weierstrass function and its derivative, though the explicit formulae are rather complicated. Although it may appear from the last expression in (3.101) that (3.100) simplifies when $c = c_-$, such a choice would break the double periodicity of our spectral curve and so is inappropriate for the fully elliptic model²⁴.

Part of the difficulty in dealing with the elliptic Ruijsenaars-Schneider and Calogero-Moser models is the complicated nature of these spectral curves. For example, in the case of $SU(3)$ the spectral curve (3.88) is

²⁴However, in the perturbative limit when one of the periods becomes infinite this choice is then available.

$$\begin{aligned}
& \lambda^3 - cu\lambda^2 + c^2v\lambda(\wp(\xi) - \wp(\epsilon)) + \\
& + c^3 \left(\frac{1}{2}\wp'(\xi)\wp'(\epsilon) - 3\wp(\xi)\wp^2(\epsilon) + \wp^3(\epsilon) + 2\alpha\wp(\xi)\wp(\epsilon) + 2\beta(\wp(\xi) + \wp(\epsilon)) + 2\gamma \right) = 0 \quad (3.102)
\end{aligned}$$

$$u \equiv H_1 = H_+, v \equiv H_2 = H_-$$

where

$$\begin{aligned}
H_{\pm} = & e^{\pm p_1} \sqrt{\wp(q_{12}) - \wp(\epsilon)} \sqrt{\wp(q_{13}) - \wp(\epsilon)} + e^{\pm p_2} \sqrt{\wp(q_{12}) - \wp(\epsilon)} \sqrt{\wp(q_{23}) - \wp(\epsilon)} + \\
& + e^{\pm p_3} \sqrt{\wp(q_{13}) - \wp(\epsilon)} \sqrt{\wp(q_{23}) - \wp(\epsilon)}, \quad p_1 + p_2 + p_3 = 0. \quad (3.103)
\end{aligned}$$

Again the issue is whether some choice of the normalization c might simplify matters. With our choice of (3.90) for example, the third term in (3.102) turns into just $v\lambda$, but no drastic simplification occurs in the constant term unless at the special point $\epsilon = \frac{\pi}{2}$ (where $\wp'(\epsilon = \pi/2) = 0$) when – modulo c – the whole equation becomes linear in $x = \wp(\xi)$.

To conclude the discussion of the Ruijsenaars-Schneider system, we separately consider the special point where ϵ equals the real half-period $\omega = \pi/2$. Using the fact that $\sigma(\xi + 2\omega) = -\sigma(\xi)e^{2\eta_1(\xi+\omega)}$ with $\eta_1 = \xi(\omega)$ [109], we find that

$$\frac{\sigma(\xi)\sigma(\omega)}{\sigma(\xi + \omega)} \equiv c_+(\xi|\omega) = \alpha c_0(\xi|\omega) = \alpha^2 c_-(\xi|\omega) \quad (3.104)$$

with $\alpha = e^{-\xi\eta_1}$. Then

$$D_k(\xi|\epsilon) = \begin{cases} c_0^{-k} \mathcal{H}_1^{-k(k-2)/2} & k \text{ even,} \\ c_0^{-k} \mathcal{H}_1^{-k(k-2)/2} c_0 \mathcal{H}_1^{-1/2} & k \text{ odd.} \end{cases} \quad (3.105)$$

Here $\mathcal{H}_1 = 2e_1^2 + e_2e_3$ is independent of ξ and we note that in the trigonometric limit $\mathcal{H}_1 = 1$. The choice $c \sim c_0$ now essentially removes all ξ -dependence in the spectral curve apart from a term linear in c_0 that multiplies the odd Hamiltonians. Similar to the $SU(3)$ case, we choose $c = -ic_0$. By absorbing the (inessential) constant factors into the Hamiltonians, $h_k = i^k \mathcal{H}_1^{-k(k-2)/2} H_k$ for even k and $h_k = i^{k-1} \mathcal{H}_1^{-1/2} \mathcal{H}_1^{-k(k-2)/2} H_k$ for odd k , one obtains the non-perturbative spectral curve (3.100) at the special point $\epsilon = \omega$:

$$ic_0^{-1}(\xi|\omega) = i\sqrt{\wp(\xi) - \wp(\omega)} = \frac{\sum_{\text{odd } k}^N h_k(-\lambda)^{N-k}}{\sum_{\text{even } k}^N h_k(-\lambda)^{N-k}} = \frac{P(\lambda) - (-)^N P(-\lambda)}{P(\lambda) + (-)^N P(-\lambda)}, \quad (3.106)$$

where

$$P(\lambda) = \sum_k h_k(-\lambda)^{N-k} = \lambda^N + \dots + 1 \quad (3.107)$$

These formulae may also be easily expressed in terms of the Jacobi elliptic functions. For example, $\wp(\xi) - \wp(\omega) = (e_1 - e_3) \frac{cn^2(u, k)}{sn^2(u, k)}$ and, therefore, $c_0(\xi|\omega) = \frac{1}{\sqrt{e_1 - e_3}} \frac{sn(u, k)}{cn(u, k)}$; here $sn(u, k)$ and $cn(u, k)$ are the Jacobi functions [109], $u = (e_1 - e_3)^{1/2} \xi$, k is elliptic modulus.

To conclude this subsection, note that one can immediately generalize the Ruijsenaars-Schneider model to other groups, see [131].

3.8 Fundamental matter in 6d: XYZ chain

At the last subsection of this long and detailed consideration of different Seiberg-Witten theories, we make the first step towards 6d theories, that is 6d theory with the fundamental matter. This is the theory with *two* extra compactified dimensions, of radii R_5 and R_6 . The same argument as we used for the transition from four to five dimensions, i.e. taking account of the Kaluza-Klein modes, allows one to predict the structure of results in the 6d case as well. Indeed, one expects that coming from 4d (5d) to 6d means one should replace the rational (trigonometric) expressions by the elliptic ones, the (imaginary part of) modular parameter being identified with the ratio of the compactification radii R_5/R_6 .

This would give rise to “elliptization” of an integrable systems. In terms of 2×2 Lax representation, the “elliptization” corresponds to replacing by elliptic r -matrices and L -operators the rational (3.34) and trigonometric (3.57) ones. In particular, it means that now it is natural to consider the parameter $\zeta = \frac{1}{2} \log \lambda$ as co-ordinate on a torus, i.e. one of the spectral parameters lives on a torus, while the other one remains on a cylinder.

In fact, we know the only system providing proper elliptization of the **XXX** spin chain, that is, the **XYZ** Sklyanin chain [132]. It was propose as a candidate for the integrable system behind the 6d theory in [97, 98, 99].

Now we briefly describe the **XYZ** chain. All necessary facts on the elliptic functions can be found in [109]. The Lax matrix is now defined on the elliptic curve $E(\tau)$ and is explicitly given by (see [113] and references therein):

$$L^{\text{SkI}}(\zeta) = S^0 \mathbf{1} + i \sum_{a=1}^3 W_a(\zeta) S^a \sigma_a \quad (3.108)$$

where

$$W_a(\zeta) = \sqrt{e_a - \wp(\zeta|\tau)} = i \frac{\theta'_*(0) \theta_{a+1}(\zeta)}{\theta_{a+1}(0) \theta_*(\zeta)} \quad (3.109)$$

To keep the similarities with [97, 98, 99], we redefine in this subsection the spectral parameter $\zeta \rightarrow i \frac{\zeta}{2K}$, where $K \equiv \int_0^{\frac{\pi}{2}} \frac{dt}{\sqrt{1-k^2 \sin^2 t}} = \frac{\pi}{2} \theta_1^2(0)$, $k^2 \equiv \frac{e_1 - e_2}{e_1 - e_3}$ so that $K \rightarrow \frac{\pi}{2}$ as $\tau \rightarrow i\infty$. This factor results into the additional multiplier π in trigonometric functions in the limiting cases.

The Lax operator (3.108) satisfies the Poisson relation (3.35) with the numerical *elliptic* r -matrix $r(\zeta) = -\frac{i}{2} \sum_{a=1}^3 W_a(\zeta) \sigma_a \otimes \sigma_a$, which implies that S^0, S^a form the (classical) Sklyanin algebra [132]:

$$\begin{aligned} \{S^a, S^0\} &= i(e_b - e_c) S^b S^c \\ \{S^a, S^b\} &= -i S^0 S^c \end{aligned} \quad (3.110)$$

with the obvious notation: abc is the triple 123 or its cyclic permutations.

One can distinguish three interesting limits of the Sklyanin algebra: the rational, trigonometric and double-scaling limits [97]. We are interested in only trigonometric limit here, since it describes degeneration to the 5d case. In this limit, $\tau \rightarrow +i\infty$ and the Sklyanin algebra (3.110) transforms to

$$\begin{aligned} \{\hat{S}^3, \hat{S}^0\} &= 0, \quad \{\hat{S}^1, \hat{S}^0\} = i \hat{S}^2 \hat{S}^3, \quad \{\hat{S}^2, \hat{S}^0\} = -2i \hat{S}^3 \hat{S}^1 \\ \{\hat{S}^1, \hat{S}^2\} &= -i \hat{S}^0 \hat{S}^3, \quad \{\hat{S}^1, \hat{S}^3\} = i \hat{S}^0 \hat{S}^2, \quad \{\hat{S}^2, \hat{S}^3\} = -i \hat{S}^0 \hat{S}^1 \end{aligned} \quad (3.111)$$

The corresponding Lax matrix is

$$L_{XXZ} = \widehat{S}^0 \mathbf{1} - \frac{1}{\sinh \pi \zeta} \left(\widehat{S}^1 \sigma_1 + \widehat{S}^2 \sigma_2 + \cosh \pi \zeta \widehat{S}^3 \sigma_3 \right) \quad (3.112)$$

and the r -matrix

$$r(\zeta) = \frac{i}{\sinh \pi \zeta} (\sigma_1 \otimes \sigma_1 + \sigma_2 \otimes \sigma_2 + \cosh \pi \zeta \sigma_3 \otimes \sigma_3) \quad (3.113)$$

Now one can note that the algebra (3.111) admits the identification $\widehat{S}_0 = e^{S_0^{trig}} + e^{-S_0^{trig}}$, $\widehat{S}_3 = e^{-S_0^{trig}} - e^{S_0^{trig}}$. With this identification and up to normalization of the Lax operator (3.112), we finally get the Lax operator (3.57) of the **XXZ** chain with the Poisson brackets (3.59).

The determinant $\det_{2 \times 2} L^{\text{Skl}}(\zeta)$ is equal to

$$\det_{2 \times 2} L^{\text{Skl}}(\zeta) = \widehat{S}_0^2 + \sum_{a=1}^3 e_a \widehat{S}_a^2 - \wp(\zeta) \sum_{a=1}^3 \widehat{S}_a^2 = K - M^2 \wp(\zeta) = K - M^2 x \quad (3.114)$$

where

$$K = \widehat{S}_0^2 + \sum_{a=1}^3 e_a(\tau) \widehat{S}_a^2 \quad M^2 = \sum_{a=1}^3 \widehat{S}_a^2 \quad (3.115)$$

are the Casimir functions of the Sklyanin algebra.

Now, in order to construct the spectral curve, note that the determinant of the monodromy matrix $T_{N_c}(\zeta)$ is

$$\begin{aligned} Q(\zeta) &= \det_{2 \times 2} T_{N_c}(\zeta) = \prod_{i=1}^{N_c} \det_{2 \times 2} L^{\text{Skl}}(\zeta - \zeta_i) = \prod_{i=1}^{N_c} (K_i - M_i^2 \wp(\zeta - \zeta_i)) = \\ &= Q_0 \frac{\prod_{\alpha=1}^{N_f} \theta_*(\zeta - m_\alpha)}{\prod_{i=1}^{N_c} \theta_*(\zeta - \zeta_i)^2}, \quad Q_0 = \prod_i A_i \end{aligned} \quad (3.116)$$

where we used the Appendix of [99] in order to rewrite $K_i - M_i^2 \wp(\zeta - \zeta_i)$ as $A_i \frac{\theta_*(\zeta - m_i^{(+)}) \theta_*(\zeta - m_i^{(-)})}{\theta_*^2(\zeta - \zeta_i)}$. Here

$$A_i = \frac{M_i^2 [\theta'_*(0)]^2}{\theta_*^2(\kappa_i)}, \quad m_i^{(\pm)} = \zeta_i \pm \kappa_i \quad (3.117)$$

and κ_i is a solution to the equation

$$\frac{\theta_4^2(\kappa_i)}{\theta_*^2(\kappa_i)} = \frac{K_i \theta_4^2(0)}{M_i^2 [\theta'_*(0)]^2} + \frac{\pi^2 \theta_4^2(0) [\theta_2^4(0) + \theta_3^4(0)]}{3 [\theta'_*(0)]^2} \quad (3.118)$$

Similarly to the $5d$ case,

$$\sum_{\alpha=1}^{N_f} m_\alpha = 2 \sum_{i=1}^{N_c} \zeta_i \quad (3.119)$$

As a particular example, let us consider the case of the *homogeneous* chain [97] (all $\zeta_i = 0$ in (3.116)). $\text{Tr} T_{N_c}(\zeta)$ is the combination of polynomials:

$$P(\zeta) = Pol_{\left[\frac{N_c}{2}\right]}^{(1)}(x) + y Pol_{\left[\frac{N_c-3}{2}\right]}^{(2)}(x), \quad (3.120)$$

where $\left[\frac{N_c}{2}\right]$ is integral part of $\frac{N_c}{2}$, and coefficients of $Pol^{(1)}$ and $Pol^{(2)}$ are Hamiltonians of the **XYZ** model. The spectral equation (3.13) for the **XYZ** model is:

$$w + \frac{Q(\zeta)}{w} = 2P(\zeta) \quad (3.121)$$

where for the homogeneous chain P and Q are polynomials in $x = \wp(\zeta)$ and $y = \frac{1}{2}\wp'(\zeta)$. Eq. (3.121) describes the double covering of the elliptic curve $E(\tau)$: with generic point $\zeta \in E(\tau)$ one associates the two points of Σ^{XYZ} , labeled by the two roots $w_{\pm}$ of equation (3.121). The ramification points correspond to $w_+ = w_- = \pm\sqrt{Q}$, or $Y = \frac{1}{2}\left(w - \frac{Q}{w}\right) = \sqrt{P^2 - Q} = 0$. Note that, for the curve (3.121), $x = \infty$ is *not* a branch point, therefore, the number of cuts on the both copies of $E(\tau)$ is N_c and the genus of the spectral curve is $N_c + 1$.²⁵

In the case of a generic *inhomogeneous* chain the trace of monodromy matrix looks more complicated, being expression of the form

$$\begin{aligned} \text{Tr} T_{N_c}(\zeta) &= \text{Tr} \prod_{N_c \geq i \geq 1}^{\widehat{}} L_i^{\text{Skl}}(\zeta - \zeta_i) = \text{Tr} \sum_{\{\alpha_i\}} \prod_{N_c \geq i \geq 1}^{\widehat{}} \sigma^{\alpha_i} S_i^{\alpha_i} W_{\alpha_i}(\zeta - \zeta_i) = \\ &= \sum_{\{\alpha_i\}} \text{Tr} \prod_{N_c \geq i \geq 1}^{\widehat{}} \sigma^{\alpha_i} S_i^{\alpha_i} W_{\alpha_i}(\zeta - \zeta_i) \end{aligned} \quad (3.124)$$

$$\alpha_i = 0, a_i; \quad a_i = 1, 2, 3; \quad W_0(\zeta) = 1$$

It is clear from (3.109) that each term in the sum (3.124) (and thus the whole sum) has simple poles in all inhomogeneities $\zeta = \zeta_i$. Thus, the trace of the monodromy matrix can be represented as

$$P(\zeta) = \frac{1}{2} \text{Tr} T_{N_c}(\zeta) = P_0 \frac{\prod_{i=1}^{N_c} \theta_*(\zeta - a_i)}{\prod_{i=1}^{N_c} \theta_*(\zeta - \zeta_i)} \quad (3.125)$$

$$\sum_i a_i = \sum_i \zeta_i = \frac{1}{2} \sum_{\alpha} m_{\alpha} \quad (3.126)$$

where a_i should be identified with (shifted) gauge moduli. The equation (3.121) can be finally rewritten as

$$w + Q_0 \frac{\prod_{\alpha=1}^{N_f} \theta_*(\zeta - m_{\alpha})}{w \prod_{i=1}^{N_c} \theta_*(\zeta - \zeta_i)^2} = P_0 \frac{\prod_{i=1}^{N_c} \theta_*(\zeta - a_i)}{\prod_{i=1}^{N_c} \theta_*(\zeta - \zeta_i)} \quad (3.127)$$

We see that in the $6d$ case it is necessary always to require $N_f = 2N_c$ in order to have a single-valued meromorphic functions on torus. We also meet again the constraint (3.126). In the $5d$ case, it was

²⁵Rewriting analytically Σ^{XYZ} as a system of equations

$$y^2 = \prod_{a=1}^3 (x - e_a), \quad Y^2 = P^2 - Q \quad (3.122)$$

the set of holomorphic 1-differentials on Σ^{XYZ} can be chosen as

$$v = \frac{dx}{y}, \quad V_{\alpha} = \frac{x^{\alpha} dx}{yY} \quad \alpha = 0, \dots, \left[\frac{N_c}{2}\right], \quad \widetilde{V}_{\beta} = \frac{x^{\beta} dx}{Y} \quad \beta = 0, \dots, \left[\frac{N_c-3}{2}\right] \quad (3.123)$$

with the total number of holomorphic 1-differentials $1 + \left(\left[\frac{N_c}{2}\right] + 1\right) + \left(\left[\frac{N_c-3}{2}\right] + 1\right) = N_c + 1$ being equal to the genus of Σ^{XYZ} .

removed, along with the condition $N_f = 2N_c$ twisting the spin chain. Since no twisted **XYZ** chain is known we are forced to preserve both constraints²⁶.

The curve (3.127) has no longer that simple expression in terms of polynomials of x and y similar to (3.120). Thus, in the inhomogeneous case it is not convenient to work in terms of “elliptic” variables x and y (analogous to λ in the $5d$ case) instead we deal with the variable $\zeta \in \mathbb{C}/\Gamma$.

Given the spectral curve and integrable system, we complete the description with writing down the generating 1-differential dS . It can be evidently chosen in the following way

$$dS^{XYZ} \cong \zeta \frac{dw}{w} \cong -\log w d\zeta \quad (3.128)$$

since, as any spin chain, it is to be proportional to dw/w and to ζ due to six dimensions (in 5 dimensions, ζ was just $\log \lambda$ and in 6 dimensions λ is replaced by the pair (x, y) related with ζ via $\wp(\zeta)$ and $\wp'(\zeta)$). Return to our oversimplified example of the homogeneous chain. Now, under the variation of moduli (which are all contained in P , while Q is moduli independent),

$$\delta(dS^{XYZ}) \cong \frac{\delta w}{w} d\zeta = \frac{\delta P(\zeta)}{\sqrt{P(\zeta)^2 - 4Q(\zeta)}} d\zeta = \frac{dx}{yY} \delta P \quad (3.129)$$

and, according to (3.123), the r.h.s. is a *holomorphic* 1-differential on the spectral curve (3.121).

4 Integrability and $\mathcal{N} = 1$ supersymmetric YM theories

4.1 Superpotentials in $\mathcal{N} = 1$ theories from $\mathcal{N} = 2$ spectral curves

In this section we shall discuss gauge theories with $\mathcal{N} = 1$ SUSY and their relation with the $\mathcal{N} = 2$ SUSY theories. For the detailed review of $\mathcal{N} = 1$ theories we recommend [133]. Here we explain some features of $\mathcal{N} = 1$ theories important for treating them within the integrable framework.

The key difference between theories with $\mathcal{N} = 1$ and $\mathcal{N} = 2$ SUSY is the absence of vacuum valleys in the $\mathcal{N} = 1$ case. Therefore, the Coulomb branch of the moduli space is absent and the number of the vacuum states is finite. The order parameter which distinguishes between the vacua can be identified with the gluino condensate. Due to the discreteness of vacua the spectrum of BPS states now is different. Instead of the particles saturating the BPS bound one has domain walls interpolating between vacua as well as stringy objects which can be identified with the strings themselves or with the domain wall intersections. These objects turn out to be related with the corresponding central charges in the algebra of $\mathcal{N} = 1$ supersymmetry [134, 135].

The main target is to be an object that presumably could be determined exactly. Instead of the prepotential in $\mathcal{N} = 2$ theory, the function which enjoys holomorphy is the superpotential $\mathcal{W}$; therefore, one could attempt to evaluate it. There are generically perturbative and nonperturbative contributions to the superpotential. As for the perturbative contribution, the β -function unlike the $\mathcal{N} = 2$ case gets contributions from all loops. However, the higher loops IR contributions can be calculated exactly [136]. There is also a crucial difference between nonperturbative contributions in $\mathcal{N} = 1$ and $\mathcal{N} = 2$ theories. Indeed, configurations of all possible instanton numbers contribute

²⁶Formula (3.126) implies that, while the sum of all the would-be gauge moduli a_i still remains a constant, it is no longer zero. Therefore, one would rather associate with the gauge moduli the quantities a_i shifted by the constant $\frac{1}{2N_c} \sum m_\alpha$. We do not know the proper field theory interpretation of this condition.

into the prepotential in the $\mathcal{N} = 2$ case, while only one-instanton terms contribute into the $\mathcal{N} = 1$ superpotential due to the non-anomalous R-symmetry.

Despite all this difference one may expect links between the exact results in $\mathcal{N} = 2$ theories and the superpotential. These expectations are true indeed and below we sketch the relation following [137] and especially [138]. The starting point is the Seiberg-Witten curve $P(x, y) = 0$ in $\mathcal{N} = 2$ theory with some matter content. Then, one should fix singular points on the Coulomb branch of the moduli space which correspond to the vacua in $\mathcal{N} = 1$ theory. These singular points are in one-to-one correspondence with vanishing of the discriminant of the curve. Then, comparing the equations in $\mathcal{N} = 2$ theory and the equation $\partial\mathcal{W} = 0$, where the derivative is taken w.r.t. the appropriated composite field, one observes that these equations coincide if the coordinate on the spectral curve is identified with some composite operator.

Consider $\mathcal{N} = 1$ supersymmetric $SU(2)$ gauge theories in four dimensions, with any possible content of matter superfields, such that the theory is either one-loop asymptotic free or conformal. This allows one to introduce $2N_f$ matter supermultiplets in the fundamental representation, Q_i^a , $i = 1, \dots, 2N_f$ and N_A supermultiplets in the adjoint representation, Φ_α^{ab} , $\alpha = 1, \dots, N_A$. Here a, b are fundamental representation indices, and $\Phi^{ab} = \Phi^{ba}$. The numbers N_f , N_A are limited by the condition to keep the theory asymptotically free or conformal:

$$b_1 = 6 - N_f - 2N_A \geq 0, \quad (4.130)$$

where $-b_1$ is the one-loop coefficient of the gauge coupling beta-function.

The effective superpotential of an $\mathcal{N} = 1$ supersymmetric $SU(2)$ gauge theory, with $2N_f$ doublets and N_A triplets is²⁷

$$\begin{aligned} \mathcal{W}_{N_f, N_A}(M, X, Z) = & -(4 - b_1) \left\{ \Lambda^{-b_1} \text{Pf}_{2N_f} X \left[\det_{N_A} (\Gamma_{\alpha\beta}) \right]^2 \right\}^{1/(4-b_1)} \\ & + \text{tr}_{N_A} \mu M + \frac{1}{2} \text{tr}_{2N_f} m X + \frac{1}{\sqrt{2}} \text{tr}_{2N_f} Y^\alpha Z_\alpha, \end{aligned} \quad (4.131)$$

where

$$\Gamma_{\alpha\beta}(M, X, Z) = M_{\alpha\beta} + \text{tr}_{2N_f} (Z_\alpha X^{-1} Z_\beta X^{-1}). \quad (4.132)$$

The first term in (4.131) is the exact non-perturbative superpotential, while the other terms give the tree-level superpotential. Λ here is a dynamically generated scale, while $\mu_{\alpha\beta}$, m_{ij} and Y_{ij}^α are the bare masses and Yukawa couplings, respectively ($\mu_{\alpha\beta} = \mu_{\beta\alpha}$, $m_{ij} = -m_{ji}$, $Y_{ij}^\alpha = Y_{ji}^\alpha$). The gauge singlets, X , M , Z , are given in terms of the $\mathcal{N} = 1$ superfield doublets, Q^a , the triplets, Φ^{ab} as follows:

$$\begin{aligned} X_{ij} &= Q_{ia} Q_j^a, a = 1, 2, ; i, j = 1, \dots, 2N_f, \\ M_{\alpha\beta} &= \Phi_{\alpha b}^a \Phi_{\beta a}^b; \alpha, \beta = 1, \dots, N_A; a, b = 1, 2, \\ Z_{ij}^\alpha &= Q_{ia} \Phi_{\alpha b}^a Q_j^b \end{aligned} \quad (4.133)$$

The gauge-invariant superfields X_{ij} may be considered as a mixture of $SU(2)$ “mesons” and “baryons”, while the gauge-invariant superfields Z_{ij}^α may be considered as a mixture of $SU(2)$ “meson-like” and “baryon-like” operators.

As an example, consider the theory with $N_f = 1$, $N_A = 1$. In this case, the superpotential reads

²⁷Throughout this section, for the sake of brevity we omit the index QCD from the notation of Λ_{QCD} .

$$\mathcal{W}_{1,1} = -\frac{\text{Pf}X}{\Lambda^3}\Gamma^2 + \mu M + \frac{1}{2}\text{tr}mX + \frac{1}{\sqrt{2}}\text{tr}YZ \quad (4.134)$$

Here m , X are antisymmetric 2×2 matrices, Y , Z are symmetric 2×2 matrices and

$$\Gamma = M + \text{tr}(ZX^{-1})^2. \quad (4.135)$$

To find the quantum vacua, we solve the equations: $\partial_M W = \partial_X W = \partial_Z W = 0$. The equations $\partial W = 0$ can be re-organized into the singularity conditions of an elliptic curve:

$$y^2 = x^3 + ax^2 + bx + c \quad (4.136)$$

(and some other equations), where the coefficients a, b, c are functions of only the field M , the scale Λ , the bare quark masses, m , and Yukawa couplings, Y . Explicitly,

$$a = -M, \quad b = \frac{\Lambda^3}{4}Pfm, \quad c = -\frac{\alpha}{16}, \quad (4.137)$$

where

$$\alpha = \frac{\Lambda^6}{4}\det Y. \quad (4.138)$$

The parameter x in the elliptic curve is given in terms of the composite field:

$$x \equiv \frac{1}{2}\Gamma. \quad (4.139)$$

$\mathcal{W}_{1,1}$ has $2 + N_f = 3$ vacua, namely, the three singularities of the elliptic curve. These are the three solutions, $M(x)$, of the equations: $y^2 = \partial y^2 / \partial x = 0$; the solutions for X , Z are given by the other equations of motion.

Similarly, if one considers a generic $SU(N_c)$ group, the spectral curve for the generic N_f , N_c can be put into the form (3.41)

$$y^2 = P_{N_c}^2(\lambda) + Q_{N_f}(\lambda) \quad (4.140)$$

with $\lambda = ZX^{-1}$ [137].

From this consideration, we can get an important lesson for the integrability approach; that is, one can make an identification of one of the coordinates defining the spectral curve. Indeed, the Lax operator of the spin chain (when acting on the eigenfunction, Baker-Akhiezer function) can be identified with some known composite operator. To get the pure gauge theory one has to decouple the fundamental matter sending its masses to infinity. Therefore, the Lax operator in the Toda system can also be identified as a composite operator which involves very heavy “regulator” fermions in the fundamental representation. The identification of the second coordinate involved in the equation of the spectral curve (4.140) is more tricky.

Let us note that the relation between $\mathcal{N} = 1$ superpotentials and $\mathcal{N} = 2$ curves can be established moving in the opposite direction as well [139]. To this aim one can consider the Affleck-Dine-Seiberg superpotential supplemented by the contribution from the heavy adjoint hypermultiplet and use the Konishi anomaly relating the vacuum expectation values of different operators

$$\begin{aligned}
\left\langle \frac{1}{2} \text{tr} m X + \frac{1}{\sqrt{2}} \text{tr} Y Z + \frac{1}{2} \frac{\text{Tr} W^2}{8\pi^2} \right\rangle &= 0 \\
\left\langle 2 \text{tr} \mu M + \frac{1}{\sqrt{2}} \text{tr} Y Z + 2 \frac{\text{Tr} W^2}{8\pi^2} \right\rangle &= 0
\end{aligned} \tag{4.141}$$

where the vacuum expectation value of $\text{Tr} W^2$ is the gluino condensate. The combination of the equation $\partial \mathcal{W} = 0$ and the Konishi relations amounts exactly to the condition of vanishing discriminant of the Seiberg-Witten curve.

Generically one can consider more specific degenerations of the Seiberg-Witten curves resulting into specific $\mathcal{N} = 1$ theories. For instance, one can consider all B -cycles on the $\mathcal{N} = 2$ curves vanishing simultaneously [140]. In this case, the very peculiar spectrum of strings emerges in $\mathcal{N} = 1$ theory [141]. One more interesting possibility comes from the phenomenon of collisions of vacua at some points in the parameter space [142] which is expected to correspond to nontrivial conformal field theories in four dimensions [143]. It was recently shown that, using the correspondence between $\mathcal{N} = 2$ and $\mathcal{N} = 1$ theories, one can identify the phase transition at these Argyres-Douglas points as the deconfinement phase transition [139]. The analysis of the Argyres-Douglas points within the integrability approach remains an interesting problem.

4.2 Superpotentials from nonperturbative configurations

We used the analysis above to recognize some meaning of the Lax operator. Now let us make the next step and identify the configuration space of the dynamical system under consideration in terms of the field theory. This can be done via calculating superpotentials from nonperturbative configurations [144, 145, 146]. In the brief description below we follow the paper [144]. To have the convenient starting point, one compactifies the four-dimensional theory to three dimensions on a circle of radius R . The most obvious modification is that the theory acquires new scalar degrees of freedom which come from the Wilson lines of the four-dimensional gauge field swapped around the compact dimension. Non-zero values for these scalars generically break the gauge group down to its maximal Abelian subalgebra, $SU(N_c) \rightarrow U(1)^{N_c-1}$. Hence, at the classical level, the compactified theory has a Coulomb branch. By a gauge rotation, the Wilson line can be chosen to lie in the Cartan subalgebra, $\phi = \phi^a H^a$. Each scalar ϕ^a is related to the corresponding Cartan component of the four-dimensional gauge field as $\phi^a = \oint_{S^1} A^a \cdot dx$. As a consequence, ϕ^a can be shifted by an integer multiple of 2π by performing a topologically non-trivial gauge transformation which is single-valued in $SU(N_c)/Z_{N_c}$. Thus we learn that each ϕ^a is a periodic variable with period 2π .

In addition to the N_c periodic scalars ϕ^a , the low energy theory on the Coulomb branch also includes N_c massless photons A_i^a , with $\sum_{a=1}^{N_c} A_i^a = 0$, where $i = 1, 2, 3$ is a three-dimensional Lorenz index. In three dimensions, a massless Abelian gauge field can be eliminated in favour of a scalar by a duality transformation. Thus we can eliminate the gauge fields A_i^a , in favour of N_c scalars σ^a subject to $\sum_{i=1}^{N_c} \sigma^a = 0$. In particular, the fields σ^a appear in the effective action as Lagrange multipliers for the Bianchi identity: $\mathcal{L}_\sigma \sim \sigma \varepsilon^{ijk} \partial_i F_{jk}$. Specifically, the normalization of σ^a is chosen so that the resulting term in the action is precisely $-i\sigma^a k^a$ where k^a is the magnetic charge in the $U(1)$ subgroup corresponding to the Cartan generator H^a . The Dirac quantization condition implies that each magnetic charge is quantized in integer units: $k^a \in \mathbb{Z}$, and so the path integral is invariant under shifts of the form $\sigma^a \rightarrow \sigma^a + 2\pi n^a$ with $n^a \in \mathbb{Z}$. Thus, like ϕ^a , each σ^a is a periodic variable with period 2π .

It is natural to combine the real scalars ϕ and σ to form a complex scalar $X = -i(\sigma + \tau\phi)$ [147]

where $\tau = 4\pi i/g^2 + \theta/2\pi$ is the usual complexified coupling. Let us now concentrate on the Euclidean theory with gauge group $SU(2)$. Gauge field configurations in the compactified theory are labelled by two distinct kinds of topological charge. The first is the Pontryagin number carried by instantons in four-dimensions,

$$p = \frac{1}{8\pi^2} \int_{R^3 \times S^1} \text{Tr} [F \wedge F] \quad (4.142)$$

The $4d$ instanton number appears in the microscopic action as the term $-ip\theta$. An important feature of the compactified theory which differs from the theory on R^4 is that p is not quantized in integer units. However, for each integer value of p , the theory has solutions with action $S_{cl} = 8\pi^2|p|/g^2 - ip\theta$. Thus we have $S_{cl} = 2\pi ip\tau$ for $p > 0$ and $S_{cl} = -2\pi ip\bar{\tau}$ for $p < 0$. When the instanton scale size is much less than the compactification radius, these solutions are close to their counterparts on R^4 .

In addition to the usual four-dimensional instantons, the compactified theory also has finite action configurations which carry three-dimensional magnetic charge,

$$k = \frac{1}{8\pi} \int_{R^3} \vec{\nabla} \cdot \vec{B} \in Z \quad (4.143)$$

where $B_i = \varepsilon_{ijk} F^{jk}/2$ is the Abelian magnetic field of the low-energy theory. As these configurations do not depend either on the time or on the coordinate of the compactified spatial dimension, we will refer to them as three-dimensional instantons. In addition to the integer-valued magnetic charge k , these $3d$ instantons also carry fractional Pontryagin number $p = \phi/2\pi$. The magnetic charge appears in the action as the term $-ik\sigma$ while, as above, $4d$ instanton number appears as the term $-ip\theta$. Including these term the action of a $3d$ instanton can be written as $S_{3d} = -i\tau\phi|k| + ik\sigma$. Thus we have $S_{3d} = kX$ for $k > 0$, and $S_{3d} = k\bar{X}$ for $k < 0$.

Putting everything together we can write out the most general possible superpotential which can be generated at leading semi-classical order,

$$\begin{aligned} \mathcal{W} = & m_1 m_2 m_3 \sum_{p=1}^{\infty} a_p q^p + m_1 m_2 m_3 \sum_{k=1}^{\infty} b_k \exp(-kX) + \\ & + m_1 m_2 m_3 \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} c_{k,n} q^{kn} \exp(-kX) + d_{k,n} q^{kn} \exp(+kX) \end{aligned} \quad (4.144)$$

The dependence of the superpotential on the masses of the adjoint scalars m_i is fixed by the zero modes counting.

The superpotential of the $SU(2)$ theory is a holomorphic function on the chiral superfield $X = -i(\sigma + \tau\phi)$. The weak coupling arguments indicate that this function should respect the classical periodicity of the variables ϕ and σ . Thus, $\mathcal{W}(X)$ is a holomorphic function defined on the complex torus E . The superpotential should also be invariant under the Weyl group of $SU(2)$ which acts as $X \rightarrow -X$. As $\mathcal{W}(X)$ is a non-constant holomorphic function on a compact domain it must have a singularity somewhere. As usual such a singularity should correspond to a point at which our low-energy description of the theory breaks down due to the presence of extra light degrees of freedom. There is only one such point on the classical moduli space: the origin $X = 0$ where non-Abelian gauge symmetry is restored. Thus we will assume that $\mathcal{W}$ has an isolated singularity at this point and no other singularities. Finally, as there is only one elliptic function of order two with a double pole at the origin we have the result [144],

$$\mathcal{W}(X) = m_1 m_2 m_3 (\mathcal{P}(X) + C(\tau)) \quad (4.145)$$

Importantly, we have obtained a superpotential which is independent of R .

Starting from an $\mathcal{N} = 2$ theory in four dimensions, the superpotential of the $\mathcal{N} = 1$ theory on $R^3 \times S^1$ obtained by introducing a mass for the adjoint scalar in the $\mathcal{N} = 2$ vector multiplet should coincide with the complexified potential of the corresponding integrable system. This connection should be a general one which applies to all $\mathcal{N} = 1$ theories obtained by soft breaking of $\mathcal{N} = 2$ supersymmetry. The key point is that the Coulomb branch of an $\mathcal{N} = 2$ theory on $R^3 \times S^1$ coincides with the complexified phase space of the integrable Hamiltonian system associated with the corresponding theory on R^4 . The dynamical variables of the integrable system, $\{X^a, P^a\}$ yield holomorphic coordinates on the Coulomb branch of the compactified theory. The dimensional transmutation provides then the corresponding superpotential for the Toda case too. The superpotential for the Toda case has been also derived from the brane picture for instantons as wrapped branes [146].

4.3 $\mathcal{N} = 1$ theories and integrability

We complete this section with general comments concerning the relation between integrable dynamics and $\mathcal{N} = 1$ SYM theories. First of all, one should expect that superpotentials in $\mathcal{N} = 1$ theories are associated with Hamiltonians of integrable systems²⁸. Therefore, the vacuum in $\mathcal{N} = 1$ theory correspond to a stationary state of the dynamical system

$$\partial\mathcal{W}/\partial\phi_i = \partial H/\partial x_i = 0 \quad (4.146)$$

and the number of vacua in $\mathcal{N} = 1$ theory coincides with the number of stationary states. Note that the locus corresponding to the stationary states of the elliptic Calogero system plays the essential role in the relation between the Calogero system and the KdV hierarchy. Namely, the initial data in the KdV hierarchy upon this correspondence should follow from the equilibrium locus in the Calogero system.

Let us now say how to come from the $\mathcal{N} = 2$ theory to the $\mathcal{N} = 1$ one in integrable terms. First, the $\mathcal{N} = 2$ system has to be placed at the singularities of the Coulomb branch. This means that some integrals of motion in the integrable system have to be fixed. If we are sitting at the point of maximal degeneracy, all the integrals of motion have to be fixed and the finite-gap solution degenerates to the solitonic one. This has been carefully done in the paper [148]. At the next step, one has to give a mass to the adjoint scalar that breaks $\mathcal{N} = 2$ SUSY down to $\mathcal{N} = 1$. The spectral curve is then “rotated” within $\mathbb{C}P^3$ and is given by two polynomial equations

$$P(v, u, x) = 0, \quad Q(v, u, x) = 0 \quad (4.147)$$

Peculiar expressions for these polynomials for different $\mathcal{N} = 1$ theories are presented in [53]. These equations definitely mean that the integrable system is to be modified. However, the question of possible interpretation of the spectral curve of in $\mathcal{N} = 1$ theory as a spectral curve of some integrable system still remains open.

We have discussed above the (very restricted) role played by the $\mathcal{N} = 2$ integrable system in the context of $\mathcal{N} = 1$ theory. However, it seems that the Whitham integrability (see section 6 below) is much more prominent candidate for describing the $\mathcal{N} = 1$ theories, since it actually governs the

²⁸The idea of this identification comes from the fact that supersymmetry is basically broken by the $\text{tr}\Phi^2$ -term whose vacuum value is the Hamiltonian of the corresponding $\mathcal{N} = 2$ integrable system (we discuss the $SU(2)$ case), see also below.

dependence of the effective action on $\mathcal{N} = 2$ SUSY breaking parameters. In a sense, the Whitham theory which is nothing but RG flows predicts a very peculiar “time”-dependence consistent with RG flows [149]. All the “times” but the discrete one are SUSY breaking parameters and some predictions concerning even $\mathcal{N} = 0$ theories can be made²⁹.

5 Perturbative prepotentials

After a short discussion of $\mathcal{N} = 1$ SUSY theories, we return to the main subject of this survey, $\mathcal{N} = 2$ theories.

In previous sections we established connections between Seiberg-Witten theories and gauge theories. However, so far we studied only a kind of general landscape in the theory, since the real, numerical results enter the game only with the prepotential calculated. Note that even the correspondence between gauge and integrable theories is supported by comparing the perturbative prepotentials. We discuss this issue in this section.

In fact, as we noted already, calculating the prepotential is quite a challenging task. It can be done, mostly in the framework of integrable systems, order by order. However, technically one needs to use different specific properties of prepotentials to simplify these calculations.

The other essential point is that the prepotential celebrates many properties that allow one to push forward the theory in many directions. This makes the prepotential an important object from both mathematical and physical standing points.

5.1 Perturbative prepotentials in gauge theories

We start with the perturbative prepotentials, since, on one hand, one is able to compare the prepotentials calculated in physical theories and in integrable systems only at perturbative level and, on the other hand, many important properties of the prepotential are observed already at this level. Therefore, now we are going to give an explicit description of the perturbative prepotentials for the Seiberg-Witten theories discussed in section 3.

The technical tool that allows one to proceed with effective perturbative expansion of the prepotential is “the residue formula”, the variation of the period matrix, i.e. the third derivatives of $\mathcal{F}(a)$ (see, e.g. [39]):

$$\frac{\partial^3 \mathcal{F}}{\partial a_i \partial a_j \partial a_k} = \frac{\partial T_{ij}}{\partial a_k} = \text{res}_{d\xi=0} \frac{d\omega_i d\omega_j d\omega_k}{\delta dS}, \quad (5.1)$$

where $\delta dS \equiv d\left(\frac{dS}{d\xi}\right) d\xi$, or, explicitly

$$\begin{aligned} \delta dS &= d\lambda d\xi && \text{for } 4d \text{ models} \\ \delta dS &= R_5^{-1} d\log \lambda d\xi && \text{for } 5d \text{ models} \\ \delta dS &= d\zeta d\xi && \text{for } 6d \text{ models} \end{aligned} \quad (5.2)$$

²⁹Within the Whitham treatment, one can re-sort the argument identifying superpotentials and Hamiltonians of integrable systems, at least, in the $SU(2)$ case. Indeed, one just needs to note that $\frac{\partial^2 \mathcal{F}}{\partial (\log \Lambda)^2} = \frac{\partial h_2}{\partial \log \Lambda}$ (see (6.46), (6.47)) is proportional to the gluino condensate (as a consequence of low-energy theorems). On the other hand, this condensate is obtained by taking the derivative of the superpotential w.r.t. Λ as it follows from Ward identities.

Here the differential $d\xi$ given on the bare torus degenerates into dw/w when torus degenerates into the punctured sphere (Toda and spin chains). We remark that although $d\xi$ does not have zeroes on the *bare* spectral curve when it is a torus or doubly punctured sphere, it does in general however possess them on the covering $\mathcal{C}$. Hereafter, we denote via ξ the variable living on the bare spectral curve (torus), while ζ is the variable living on the compactification torus (in 6 dimensions) or on the compactification circle (in 5 dimensions).

In order to construct the perturbative prepotentials, one merely can note that, at the leading order, the Riemann surface (spectral curve of the integrable system) becomes rational. Therefore, the residue formula allows one to obtain immediately the third derivatives of the prepotential as simple residues on the sphere. In this way, one can check that the prepotential has actually the form (3.1) [39]. We discuss these calculations later, but now see what the perturbative prepotentials are expected from the gauge theory point of view.

As a concrete example, let us consider the $SU(N_c)$ gauge group. Then, say, the perturbative prepotential for the pure gauge theory acquires the form

$$\mathcal{F}_{pert}^V = \frac{1}{8\pi i} \sum_{ij} (a_i - a_j)^2 \log(a_i - a_j) \quad (5.3)$$

This formula establishes that when v.e.v.'s of the scalar fields in the gauge supermultiplet are non-vanishing (perturbatively a_r are eigenvalues of the vacuum expectation matrix $\langle\phi\rangle$), the fields in the gauge multiplet acquire masses $m_{rr'} = a_r - a_{r'}$ (the pair of indices (r, r') label a field in the adjoint representation of $SU(N_c)$). The eigenvalues are subject to the condition $\sum_i a_i = 0$. Analogous formula for the adjoint matter contribution to the prepotential is

$$\mathcal{F}_{pert}^A = -\frac{1}{8\pi i} \sum_{ij} (a_i - a_j + M)^2 \log(a_i - a_j + M) \quad (5.4)$$

while the contribution of one fundamental matter hypermultiplet reads as

$$\mathcal{F}_{pert}^F = -\frac{1}{8\pi i} \sum_i (a_i + m)^2 \log(a_i + m) \quad (5.5)$$

Similar formulas can be obtained for the other groups. The eigenvalues of $\langle\phi\rangle$ in the first fundamental representation of the classical series of the Lie groups are

$$\begin{aligned} B_n (SO(2n+1)) : & \quad \{a_1, \dots, a_n, 0, -a_1, \dots, -a_n\}; \\ C_n (Sp(n)) : & \quad \{a_1, \dots, a_n, -a_1, \dots, -a_n\}; \\ D_n (SO(2n)) : & \quad \{a_1, \dots, a_n, -a_1, \dots, -a_n\} \end{aligned} \quad (5.6)$$

while the eigenvalues in the adjoint representation have the form

$$\begin{aligned} B_n : & \quad \{\pm a_j; \pm a_j \pm a_k\}; \quad j < k \leq n \\ C_n : & \quad \{\pm 2a_j; \pm a_j \pm a_k\}; \quad j < k \leq n \\ D_n : & \quad \{\pm a_j \pm a_k\}, \quad j < k \leq n \end{aligned} \quad (5.7)$$

Analogous formulas can be written for the exceptional groups too. The prepotential in the pure gauge theory can be read off from the formula (5.7) and has the form

$$\begin{aligned}
B_n : \quad \mathcal{F}_{pert} &= \frac{1}{8\pi i} \sum_{i,i} \left((a_i - a_j)^2 \log(a_i - a_j) + (a_i + a_j)^2 \log(a_i + a_j) \right) + \frac{1}{2} \sum_i a_i^2 \log a_i; \\
C_n : \quad \mathcal{F}_{pert} &= \frac{1}{8\pi i} \sum_{i,i} \left((a_i - a_j)^2 \log(a_i - a_j) + (a_i + a_j)^2 \log(a_i + a_j) \right) + 2 \sum_i a_i^2 \log a_i; \quad (5.8) \\
D_n : \quad \mathcal{F}_{pert} &= \frac{1}{8\pi i} \sum_{i,i} \left((a_i - a_j)^2 \log(a_i - a_j) + (a_i + a_j)^2 \log(a_i + a_j) \right)
\end{aligned}$$

The perturbative prepotentials in 4d theories with fundamental matter are discussed in detail in [39] (see also [150]).

All these formulas can be almost immediately extended to the “relativistic” 5d $\mathcal{N} = 2$ SUSY gauge models with one compactified dimension. One can understand the reason for this “relativization” in the following way. Considering four plus one compact dimensional theory one should take into account the contribution of all Kaluza-Klein modes to each 4-dimensional field. Roughly speaking it leads to the 1-loop contributions to the effective charge of the form ($a_{ij} \equiv a_i - a_j$)

$$T_{ij} \sim \sum_{\text{masses}} \log \text{ masses} \sim \sum_m \log \left(a_{ij} + \frac{m}{R_5} \right) \sim \log \prod_m (R_5 a_{ij} + m) \sim \log \sinh R_5 a_{ij} \quad (5.9)$$

i.e. coming from 4d to 5d one should make a substitution $a_{ij} \rightarrow \sinh R_5 a_{ij}$, at least, in the formula for perturbative prepotentials.

The same general argument can be equally applied to the 6d case, or to the theory with *two* extra compactified dimensions, of radii R_5 and R_6 . Indeed, the account of the Kaluza-Klein modes allows one to predict the perturbative form of charges in the 6d case as well. Namely, one should expect them to have the form

$$\begin{aligned}
T_{ij} &\sim \sum_{\text{masses}} \log \text{ masses} \sim \sum_{m,n} \log \left(a_{ij} + \frac{m}{R_5} + \frac{n}{R_6} \right) \sim \\
&\sim \log \prod_{m,n} \left(R_5 a_{ij} + m + n \frac{R_5}{R_6} \right) \sim \log \theta_* \left(R_5 a_{ij} \middle| i \frac{R_5}{R_6} \right)
\end{aligned} \quad (5.10)$$

i.e. coming from 4d (5d) to 6d one should replace the rational (trigonometric) expressions by the elliptic ones, at least, in the formulas for perturbative prepotentials.

5.2 Perturbative prepotentials in integrable systems

Now we are going to compare these results with calculations done for integrable models. We start with the integrable chains [39, 98, 99].

As we saw in the previous section (ss.3.2,3.4,3.5,3.6,3.8), the spectral curve in all cases of spin chains (fundamental matter) can be put in the form

$$w + \frac{Q^{(d)}(\zeta)}{w} = 2P^{(d)}(\zeta) \quad (5.11)$$

or

$$W + \frac{1}{W} = \frac{2P^{(d)}(\zeta)}{\sqrt{Q^{(d)}(\zeta)}}, \quad W \equiv \frac{w}{\sqrt{Q^{(d)}(\zeta)}} \quad (5.12)$$

In the perturbative limit, only the first term in the r.h.s. of these formulas survives so that the spectral curve becomes rational.

The generating differential dS is always of the form (to unify the notations, we use the same letter ζ for λ in $4d$ and $\frac{1}{2}\log\lambda$ in $5d$)

$$dS = \zeta d\log W \quad (5.13)$$

The concrete forms of the functions introduced here are:

$$Q^{(4)}(\zeta) \sim \prod_{\alpha}^{N_f} (\zeta - m_{\alpha}), \quad Q^{(5)}(\zeta) \sim \prod_{\alpha}^{N_f} \sinh(\zeta - m_{\alpha}), \quad Q^{(6)}(\zeta) \sim \prod_{\alpha}^{N_f} \frac{\theta_*(\zeta - m_{\alpha})}{\theta_*^2(\zeta - \zeta_i)} \quad (5.14)$$

$$P^{(4)} \sim \prod_i^N (\zeta - a_i), \quad P^{(5)} \sim \prod_i^N \sinh(\zeta - a_i), \quad P^{(6)} \sim \prod_i^N \frac{\theta_*(\zeta - a_i)}{\theta_*(\zeta - \zeta_i)} \quad (5.15)$$

(in $P^{(5)}(\zeta)$, there is also some exponential of ζ unless $N_f = 2N$, see (3.69)).

In order to come to the pure gauge theory in $4d$ and $5d$, one should just put $Q^{(d)}(\zeta)$ equal to unity (it corresponds to sending hypermultiplet masses to infinity), while in $6d$ it is still unclear how to remove the restriction $N_f = 2N_c$ which prevents constructing the pure gauge theory.

The perturbative part of the prepotential can be calculated using these manifest expressions and the residue formula.

5.3 $4d$ theory: a simple example of calculation

Let us start from the simplest example of the Toda chain. In the perturbative limit ($\frac{\Lambda_{QCD}}{\langle\phi\rangle} \rightarrow 0$) the second term in (5.11) vanishes, the curve acquires the form

$$W = 2P_{N_c}(\lambda), \quad P_{N_c}(\lambda) = \prod (\lambda - \lambda_i) \quad (5.16)$$

(a rational curve with punctures³⁰) and the generating differential (3.19) turns into

$$dS_{\text{pert}}^{(4)} = \lambda d\log P_{N_c}(\lambda) \quad (5.17)$$

Now the set of the $N_c - 1$ independent canonical holomorphic differentials is

$$d\omega_i = \left(\frac{1}{\lambda - \lambda_i} - \frac{1}{\lambda - \lambda_{N_c}} \right) d\lambda = \frac{\lambda_{iN_c} d\lambda}{(\lambda - \lambda_i)(\lambda - \lambda_{N_c})}, \quad i = 1, \dots, N_c - 1, \quad \lambda_{ij} \equiv \lambda_i - \lambda_j \quad (5.18)$$

and the A -periods $a_i = \lambda_i$ (the independent ones are, say, with $i = 1, \dots, N_c - 1$) coincide with the roots of polynomial $P_{N_c}(\lambda)$, λ_i . The prepotential can be now computed via the residue formula

$$\mathcal{F}_{\text{pert},ijk} = \text{res}_{d\log P_{N_c}(\lambda)=0} \frac{d\omega_i d\omega_j d\omega_k}{d\log P_{N_c}(\lambda) d\lambda} \quad (5.19)$$

³⁰These punctures emerge as a degeneration of the handles of the hyperelliptic surface so that the A -cycles encircle the punctures.

which can be explicitly done. The only technical trick is that it is easier to compute the residues at the poles of $d\omega$'s instead of the zeroes of $d\log P$. This can be done immediately since there are no contributions from the infinity $\lambda = \infty$. The final results have the following form

$$\begin{aligned}
2\pi i \mathcal{F}_{pert,iii} &= \sum_{k \neq i} \frac{1}{\lambda_{ik}} + \frac{6}{\lambda_{iN_c}} + \sum_{k \neq N_c} \frac{1}{\lambda_{kN_c}}, \\
2\pi i \mathcal{F}_{pert,iij} &= \frac{3}{\lambda_{iN_c}} + \frac{2}{\lambda_{jN_c}} + \sum_{k \neq i,j,N_c} \frac{1}{\lambda_{kN_c}} - \frac{\lambda_{jN_c}}{\lambda_{iN_c} \lambda_{ij}}, \quad i \neq j, \\
2\pi i \mathcal{F}_{pert,ijk} &= 2 \sum_{l \neq N_c} \frac{1}{\lambda_{lN_c}} - \sum_{l \neq i,j,k,N_c} \frac{1}{\lambda_{lN_c}}, \quad i \neq j \neq k;
\end{aligned} \tag{5.20}$$

giving rise to the prepotential formula

$$\begin{aligned}
\mathcal{F}_{pert} &= \frac{1}{8\pi i} \sum_{ij} f^{(4)}(\lambda_{ij}) \\
f^{(4)}(x) &= x^2 \log x^2
\end{aligned} \tag{5.21}$$

Now let us turn to the case of $4d \mathcal{N} = 2$ QCD with massive hypermultiplets. Then, $Q^{(4)}(\zeta) \neq 1$ and there arise additional differentials corresponding to the derivatives of dS with respect to masses. They are of the form

$$d\omega_\alpha = -\frac{1}{2} \frac{d\lambda}{\lambda - m_\alpha} \tag{5.22}$$

It is again straightforward to use the residue formula³¹

$$\mathcal{F}_{pert,IJK} = \operatorname{res}_{d\log \frac{P}{\sqrt{Q}}=0} \frac{d\omega_I d\omega_J d\omega_K}{d\log \frac{P(\lambda)}{\sqrt{Q(\lambda)}} d\lambda}, \quad \{I, J, K, \dots\} = \{i, j, k, \dots | \alpha, \beta, \gamma, \dots\} \tag{5.23}$$

and obtain for the prepotential³²

$$\mathcal{F}_{pert} = \frac{1}{8\pi i} \sum_{i,j} f^{(4)}(a_{ij}) - \frac{1}{8\pi i} \sum_{i,\alpha} f^{(4)}(a_i - m_\alpha) + \frac{1}{32\pi i} \sum_{\alpha,\beta} f^{(4)}(m_\alpha - m_\beta) \tag{5.24}$$

upon the identification $\lambda_i = a_i$.

5.4 From 4d to higher dimensions

Now let us turn to the $5d$ case, to the **XXZ** chain. In the perturbative limit, the hyperelliptic curve (3.18) turns into the rational curve

$$W = 2 \frac{P_{N_c}(\lambda)}{\sqrt{Q_{N_f}(\lambda)}} \equiv 2 \frac{\prod_i^{N_c} (\lambda - \lambda_i)}{\sqrt{\prod_\alpha^{N_f} (\lambda - \lambda_\alpha)}} \tag{5.25}$$

³¹Let us note that, despite the differentials $d\omega_\alpha$ have the pole at infinity, this does not contribute into the residue formula because of the quadratic pole of $d\lambda$ in the denominator.

³²The term in the prepotential that depends only on masses is not essential for the standard Seiberg-Witten theory but is crucial for the prepotential to enjoy its main properties, similar to the WDVV equations. This term is unambiguously restored from the residue formula.

with $\lambda_i = e^{2a_i}$, $\sum_i^{N_c} a_i = 0$, $\lambda_\alpha = e^{2m_\alpha}$. First, let us construct the set of holomorphic differentials $d\omega_I$.

The perturbative differential dS is

$$dS = \frac{1}{2} \log \lambda \, d \log \left(\frac{P(\lambda)}{\sqrt{Q(\lambda)}} \right) = \frac{1}{2} \log \lambda d\lambda \left(\sum_{i=1}^{N_c} \frac{1}{\lambda - \lambda_i} - \frac{1}{2} \sum_{\alpha=1}^{N_f} \frac{1}{\lambda - \lambda_\alpha} \right) \quad (5.26)$$

or, equivalently,

$$dS = \zeta d\zeta \left(\sum_{i=1}^{N_c} \coth(\zeta - a_i) - \frac{1}{2} \sum_{\alpha=1}^{N_f} \coth(\zeta - m_\alpha) + N_c - \frac{1}{2} N_f \right) \quad (5.27)$$

The variation of (5.26) gives

$$\begin{aligned} 2\delta dS &= -\log \lambda d \left(\sum_{i=1}^{N_c} \frac{\delta \lambda_i}{\lambda - \lambda_i} - \frac{1}{2} \sum_{\alpha=1}^{N_f} \frac{\delta \lambda_\alpha}{\lambda - \lambda_\alpha} \right) \cong \\ &\cong \frac{d\lambda}{\lambda} \sum_{i=1}^{N_c-1} \frac{\delta \lambda_i}{\lambda_i} \left(\frac{\lambda_i}{\lambda - \lambda_i} - \frac{\lambda_{N_c}}{\lambda - \lambda_{N_c}} \right) - \frac{1}{2} \frac{d\lambda}{\lambda} \sum_{\alpha=1}^{N_f} \frac{\delta \lambda_\alpha}{\lambda_\alpha} \left(\frac{\lambda_\alpha}{\lambda - \lambda_\alpha} \right) \end{aligned} \quad (5.28)$$

where it is implied that only $N_c - 1$ parameters λ_i are independent so that $\frac{\delta \lambda_{N_c}}{\lambda_{N_c}} = -\sum_{i=1}^{N_c-1} \frac{\delta \lambda_i}{\lambda_i}$. Now the set of the $N_c - 1$ independent canonical holomorphic differentials is

$$d\omega_i = \frac{d\lambda}{\lambda} \left(\frac{\lambda_i}{\lambda - \lambda_i} - \frac{\lambda_{N_c}}{\lambda - \lambda_{N_c}} \right) = \frac{\lambda_{iN_c} d\lambda}{(\lambda - \lambda_i)(\lambda - \lambda_{N_c})} \quad i = 1, \dots, N_c - 1, \quad \lambda_{ij} \equiv \lambda_i - \lambda_j \quad (5.29)$$

and one can check that the A -periods of dS are $\log \lambda_i = 2a_i$, while the condition of vanishing the sum of all residues gives the relation (3.66) $\sum_i a_i = \frac{1}{2} \sum m_\alpha$. In s.3.5, we discussed how this strange constraint should be removed off by using the twisted **XXZ** spin chain. The analytical explanation of this is that one can easily reproduce any lack of residues by putting some additional poles at infinities in ζ or at $\lambda = 0, \infty$. It is analogous to the procedure in $4d$. We saw, however, that the same procedure does not work in $d = 6$.

The additional differentials corresponding to the derivatives of dS (5.26) with respect to masses have the form ³³

$$d\omega_\alpha = -\frac{d\lambda}{2\lambda} \frac{\lambda_\alpha}{\lambda - \lambda_\alpha} \quad (5.32)$$

³³It can be instructive to describe in detail why (5.18) arises not only in $4d$, but also in $5d$ models. Strictly speaking, in the $5d$ case, one should consider differentials on annulus – on sphere with 2 marked points. Therefore, instead of $\frac{d\lambda}{\lambda - \lambda_i}$, one rather needs to take

$$\sum_{m=-\infty}^{+\infty} \frac{d\zeta}{\zeta - a_i + m} \sim \coth(\zeta - a_i) d\zeta = \frac{\lambda + \lambda_i}{\lambda - \lambda_i} \frac{d\lambda}{2\lambda} \quad (5.30)$$

Taking now, instead of (5.18), the differentials (5.30)

$$d\omega_i = \frac{\lambda + \lambda_i}{\lambda - \lambda_i} \frac{d\lambda}{2\lambda} - \frac{\lambda + \lambda_{N_c}}{\lambda - \lambda_{N_c}} \frac{d\lambda}{2\lambda} = \frac{\lambda_{iN_c} d\lambda}{(\lambda - \lambda_i)(\lambda - \lambda_{N_c})} \quad (5.31)$$

we obtain again the formula (5.18). On the other hand, the mass differentials (5.32) look different from (5.30). It turns out, however, that the difference does not contribute into the results and, therefore, one is free to choose any of these two possible mass differentials.

The better choice for these differentials (see footnote 31) is, however, [99]

$$d\omega_\alpha = -\frac{d\lambda}{4\lambda} \frac{\lambda + \lambda_\alpha}{\lambda - \lambda_\alpha} = -\frac{d\xi}{2} \coth(\xi - m_\alpha) \quad (5.33)$$

Now one can insert these manifest expressions for differentials into the residue formula in order to calculate prepotential.

Similar formulas are also correct for the **XYZ** chain. A generic **XYZ** curve (3.127) in the perturbative limit turns into

$$W = h_0 \frac{\prod_{i=1}^{N_c} \theta_*(\zeta - a_i)}{\sqrt{\prod_{\alpha=1}^{N_f} \theta_*(\zeta - m_\alpha)}} \quad (5.34)$$

where $h_0 = \frac{2P_0}{\sqrt{Q_0}}$. Then, the rescaled generating differential (3.128) is

$$\begin{aligned} dS &= \left(\sum \log \theta_*(\zeta - a_i) - \frac{1}{2} \sum \log \theta_*(\zeta - m_\alpha) \right) d\zeta + \log h_0 d\zeta \cong \\ &\cong \zeta d\zeta \left(\sum_{i=1}^{N_c} \mathfrak{z}(\zeta - a_i) - \frac{1}{2} \sum_{\alpha=1}^{N_f} \mathfrak{z}(\zeta - m_\alpha) \right) \end{aligned} \quad (5.35)$$

where the linear piece disappears due to $\sum a_i = \frac{1}{2} \sum m_\alpha$ and $\mathfrak{z}$ denotes the Weierstrass ζ -function [109]. This expression should be compared with (5.27) in the $5d$ case.

Note that the sum of residues vanishes only provided the constraint (3.126) is fulfilled. It can not be removed by adding an additional pole at infinity, since the variable ζ , in contrast to the $4d$ and $5d$ cases, lives on the compact surface with out boundaries (torus) and, therefore, is always finite.

Now one can take the total variation of dS (we neglect the trivial variation δh_0 giving rise to the holomorphic differential $d\zeta$ on the bare torus)

$$\delta dS^{XYZ} \cong \sum_{i=1}^{N_c} \bar{\mathfrak{z}}(\zeta - a_i) \delta a_i d\zeta - \frac{1}{2} \sum_{\alpha=1}^{N_f} \bar{\mathfrak{z}}(\zeta - m_\alpha) \delta m_\alpha d\zeta \quad (5.36)$$

where $\bar{\mathfrak{z}}(\zeta|\tau) \equiv \mathfrak{z}(\zeta|\tau) - \eta(\tau)\zeta$. Therefore, the differentials related to gauge moduli are

$$d\omega_i = (\bar{\mathfrak{z}}(\zeta - a_i) - \bar{\mathfrak{z}}(\zeta - a_{N_c})) d\zeta \quad (5.37)$$

while those related to masses are

$$d\omega_\alpha = -\frac{1}{2} (\bar{\mathfrak{z}}(\zeta - m_\alpha) - \bar{\mathfrak{z}}(\zeta - a_{N_c})) d\zeta \quad (5.38)$$

The second term in this expression is due to the condition (3.126). Thus, one can see that, in the $6d$ case, mass moduli are practically identical to the gauge ones.

5.5 Perturbative prepotentials: fundamental matter

The result of calculation for the prepotentials using all these manifest expressions for the differentials and the residue formula is always of the form

$$\begin{aligned}
\mathcal{F}_{pert} = & \frac{1}{8\pi i} \sum_{i,j} f^{(d)}(a_{ij}) - \frac{1}{8\pi i} \sum_{i,\alpha} f^{(d)}(a_i - m_\alpha) + \frac{1}{32\pi i} \sum_{\alpha,\beta} f^{(d)}(m_\alpha - m_\beta) + \\
& + \frac{\delta_{d,5}}{24\pi i} (2N_c - N_f) \left(\sum_i a_i^3 + \frac{1}{4} \sum_\alpha m_\alpha^3 \right)
\end{aligned} \tag{5.39}$$

The explicit form of these functions is

$$\begin{aligned}
f^{(4)}(x) &= x^2 \log x, \quad f^{(5)}(x) = \sum_n f^{(4)}\left(x + \frac{n}{R_5}\right) = \frac{1}{3} |x^3| - \frac{1}{2} \text{Li}_3\left(e^{-2|x|}\right), \\
f^{(6)}(x) &= \sum_{m,n} f^{(4)}\left(x + \frac{n}{R_5} + \frac{m}{R_6}\right) = \sum_n f^{(5)}\left(x + n \frac{R_5}{R_6}\right) = \\
&= \left(\frac{1}{3} |x^3| - \frac{1}{2} \text{Li}_{3,q}\left(e^{-2|x|}\right) + \text{quadratic terms} \right)
\end{aligned} \tag{5.40}$$

so that

$$f^{(4)''} = \log x, \quad f^{(5)''}(x) = \log \sinh x, \quad f^{(6)''}(x) = \log \theta_*(x) \tag{5.41}$$

Note that, in the $6d$ case, N_f is always equal to $2N$, and, in $d = 5, 6$, there is a restriction $\sum a_i = \sum \zeta_i = \frac{1}{2} \sum m_\alpha$ which implies that the gauge moduli would be rather associated with a_i shifted by the constant $\frac{1}{2N} \sum m_\alpha$. In these formulas, $\text{Li}_3(x)$ is the tri-logarithm, while $\text{Li}_{3,q}(x)$ is the elliptic tri-logarithm [151]. In order to get the prepotential for the pure gauge theories, one just need to omit all the terms containing mass moduli m_α .

These formulas perfectly match the perturbative prepotentials obtained in s.5.1 in supersymmetric gauge theories. This gives another support for their identification with integrable systems.

Let us discuss now the cubic (in moduli) terms appearing in the perturbative prepotential only in higher-dimensional theories. First, take the $5d$ case, where these terms correspond to the Chern-Simons term in the field theory Lagrangian $\text{Tr}(A \wedge F \wedge F)$. In our computation they appear due to the constant piece $(N_c - \frac{1}{2}N_f)$ in the brackets in (5.27).

In fact, let us choose the special values of the second Casimir functions at m sites so that the polynomial $Q(\lambda)$ acquires the form $\lambda^m \overline{Q}_{N_f}(\lambda)$, i.e. m of $2N_c$ factors in (3.63) turn into λ^m . In this case one gets the constant piece $N_c - m - N_f/2$ in $dS^{(5)}$, with an integer $m \leq 2N_c - N_f$. Therefore, the coefficient in front of cubic term for $d = 5$ in (5.39) can be made equal to

$$c = \frac{1}{6} (N_c - m - N_f/2) \tag{5.42}$$

Restoring the dependence on R_5 in $d = 5$ in (5.39) ($a_i \rightarrow a_i R_5$ and $m_\alpha \rightarrow m_\alpha R_5$) one can study the different limits of the system. The simplest limit corresponds to the $4d$ case and is given by $R_5 \rightarrow 0$. In this limit, $f^{(5)}(x) \xrightarrow{x \sim 0} f^{(4)}(x)$, the cubic terms vanish and we reproduce the perturbative $4d$ prepotential (5.24). At the level of integrable system it is enough to replace $S_i \rightarrow R_5 S_i$, $\mu \rightarrow e^{R_5 \mu}$ in the Lax operators (3.57), (3.67) in order to reproduce the Lax operators of the **XXX** chain.

Another interesting limit is the limit of flat $5d$ space-time, i.e. $R_5 \rightarrow \infty$. In this limit, only cubic terms survive in the prepotential (5.39) for $d = 5$ (one should carefully fix the branch of $f^{(6)}(x)$ which leading to appearing of the absolute value in (5.43)):

$$\begin{aligned}
2\pi i \mathcal{F}_{pert} = & \frac{1}{12} \sum_{i,j} |a_{ij}|^3 - \frac{1}{12} \sum_{i,\alpha} |a_i + m_\alpha|^3 + \frac{1}{48} \sum_{\alpha,\beta} |m_\alpha - m_\beta| + \\
& + \frac{1}{12} (2N_c - N_f) \left(\sum_i a_i^3 + \frac{1}{4} \sum_\alpha m_\alpha^3 \right)
\end{aligned} \tag{5.43}$$

In fact, there are two different sources of cubic terms [152]. The first one comes from the function $f^{(5)}(x)$. Since this function can be obtained as the sum over the $4d$ Kaluza-Klein perturbative contributions, these cubic terms have perturbative origin and come from the 1-loop (due to the well-known effect of generation of the CS terms in odd-dimensional gauge theories).

The second source of the cubic terms is due to the bare CS Lagrangian. As it was shown in [152], one can consider these terms with some coefficient c_{cl} :

$$\frac{c_{cl}}{6} \sum_i a_i^3 \tag{5.44}$$

restricted only to satisfy the quantization condition $c_{cl} + \frac{N_f}{2} \in \mathbb{Z}$ and the inequality $|c_{cl}| \geq N_c - \frac{N_f}{2}$. In (5.42), we easily reproduce these conditions so that (5.43) coincides with [152], provided the prepotential is defined in a fixed Weyl chamber. We leave, however, the wall crossing jumps out of the discussion.

In the $6d$ case the calculations are quite similar, and one would expect the similar cubic terms correspond to $\text{Tr}(F \wedge F \wedge F)$. However, although all the difference in calculations is due to slightly different mass differentials (5.33) and (5.38), the bare cubic terms in $6d$ are absent. This occurs due to the $N_f = 2N_c$ condition (since the coefficient in front of cubic terms is proportional to $N_c - \frac{1}{2}N_f$) and, therefore, due to the cancellation of the constant term in the brackets of (5.35) as compared with (5.27). This is one of the problems with $6d$ theories: one can not deform (twist) the theory so that the constraints $N_f = 2N_c$ and (3.126) look unavoidable. As for the “quantum generated” terms, i.e. those coming from the function $f^{(6)}$, they are certainly presented, with the coefficient equal to the $5d$ coefficient.

5.6 $SU(2)$ adjoint matter in $5d$

Now let us consider the adjoint matter case. We discuss here only the $5d$ theory, i.e. the Ruijsenaars-Schneider system, since the limit to the $4d$ case is done straightforwardly, while many concrete calculations in the Calogero-Moser system look even more tedious.

Suppose we parameterize a bare curve that is elliptic by the algebraic equation,

$$\widehat{y}^2 = \widehat{x}^3 - \alpha \widehat{x}^2 - \beta \widehat{x} - \gamma. \tag{5.45}$$

Then we can use the standard parameterization via Weierstrass $\wp$ -function and its first derivative: $\widehat{x} = \wp(\xi) + \alpha/3$ and $\widehat{y} = -\frac{1}{2}\wp'(\xi)$ so that $d\xi = -2\frac{d\widehat{x}}{\widehat{y}}$.

The perturbative (weak-coupling) limit is then given by the imaginary period $2\omega'$ becoming infinite, $\tau \rightarrow i\infty$. In this weak coupling limit (without any double-scaling which we did considering the degeneration to the periodic Toda chain) the bare curve (5.45) becomes

$$y^2 = x^2(x - 1). \tag{5.46}$$

Then $\hat{x} \rightarrow x - 1/3$ with $x = \frac{1}{\sin^2 \xi}$, $\hat{y} \rightarrow y = -\frac{\cos \xi}{\sin^3 \xi}$, $\alpha = 1$, $\beta = \gamma = 0$ and $d\xi \rightarrow \frac{dx}{x\sqrt{x-1}}$, i.e. $dS = R_5^{-1} \log \lambda d\xi \rightarrow R_5^{-1} \log \lambda \frac{dx}{x\sqrt{x-1}}$. At the same limit,

$$c_0(\xi|\epsilon) \rightarrow \frac{1}{\sqrt{\sin^{-2} \xi - \sin^{-2} \epsilon}} = \frac{\sin \xi \sin \epsilon}{\sqrt{\sin(\epsilon - \xi) \sin(\epsilon + \xi)}}, \quad c_{\pm}(\xi|\epsilon) \rightarrow \frac{\sin \xi \sin \epsilon}{\sin(\epsilon \pm \xi)} e^{\mp \epsilon \xi / 3} \equiv \bar{c}_{\pm} e^{\mp \epsilon \xi / 3}. \quad (5.47)$$

The corresponding integrable system is then nothing but the trigonometric Ruijsenaars-Schneider model we already met in s.2.2. Finally the special point $\epsilon = \omega = \frac{\pi}{2}$ sees

$$\wp(\omega) = e_1 \rightarrow \frac{2}{3} \quad (5.48)$$

and the particular combination (3.90) becomes

$$c_0(\xi|\epsilon = \pi/2) \rightarrow \tan \xi. \quad (5.49)$$

Now let us look at the period matrix in the perturbative limit in the simplest $SU(2)$ case, the corresponding spectral curve being (3.95). As usual, the derivatives of the 1-form dS with respect to the moduli are holomorphic differentials. A particular choice of coordinates for the moduli will lead to the canonical holomorphic 1-differentials. For the example at hand there is only one modulus and

$$dv = \frac{\partial dS}{\partial u} = R_5^{-1} \frac{1}{\lambda} \frac{\partial \lambda}{\partial u} d\xi = R_5^{-1} \frac{c_0 d\xi}{2\lambda - c_0 u} = \frac{1}{R_5} \frac{d\xi}{\sqrt{u^2 - 4/c_0^2}} = \frac{1}{2R_5} \frac{d\xi}{\sqrt{\frac{u^2}{4} + \wp(\epsilon) - \wp(\xi)}}. \quad (5.50)$$

If we order the roots $e_3 \leq e_2 \leq e_1$, we may take for the A -integral

$$\oint_A dv = \frac{1}{i\pi} \int_{e_3}^{e_2} dv = \frac{1}{i\pi R_5 \sqrt{(\frac{u^2}{4} + \wp(\epsilon) - e_2)(e_1 - e_3)}} K \left(\sqrt{\frac{(\frac{u^2}{4} + \wp(\epsilon) - e_1)(e_2 - e_3)}{(\frac{u^2}{4} + \wp(\epsilon) - e_2)(e_1 - e_3)}} \right). \quad (5.51)$$

Here $K(q)$ is the complete elliptic integral of the first kind. Dividing dv by the right hand side of this expression would then give us the canonical holomorphic 1-differential $d\Omega$. Similarly for the B -integral we have

$$\oint_B dv = \frac{1}{i\pi} \int_{e_2}^{e_1} dv = -\frac{1}{\pi R_5 \sqrt{(\frac{u^2}{4} + \wp(\epsilon) - e_2)(e_1 - e_3)}} K \left(\sqrt{\frac{(\frac{u^2}{4} + \wp(\epsilon) - e_3)(e_1 - e_2)}{(\frac{u^2}{4} + \wp(\epsilon) - e_2)(e_1 - e_3)}} \right), \quad (5.52)$$

and so the period matrix is

$$T = \oint_B d\Omega = -iK \left(\sqrt{\frac{(\frac{u^2}{4} + \wp(\epsilon) - e_3)(e_1 - e_2)}{(\frac{u^2}{4} + \wp(\epsilon) - e_2)(e_1 - e_3)}} \right) \Bigg/ K \left(\sqrt{\frac{(\frac{u^2}{4} + \wp(\epsilon) - e_1)(e_2 - e_3)}{(\frac{u^2}{4} + \wp(\epsilon) - e_2)(e_1 - e_3)}} \right). \quad (5.53)$$

One can take the limit $\epsilon \rightarrow 0$ in this expression to obtain $T = -\tau$. This agrees perfectly with formula $\mathcal{F}_{class} \sim \tau a^2$ for the classical part of the prepotential and justifies our identification of the gauge theory coupling constant and the modulus of the bare spectral curve.

Let us now calculate the same quantities in the perturbative limit:

$$dv \rightarrow \left(\frac{1}{2R_5} \right) \frac{dx}{x\sqrt{(x-1)(U^2-x)}}, \quad (5.54)$$

with $U^2 = \frac{1}{\sin^2 \epsilon} + \frac{u^2}{4}$. We have $e_2 = e_3 = -1/3$ and $e_1 = 2/3$ in this limit and so the A -period in this case shrinks to a contour around $x = 0$. Now $\oint_A dv = 1/(2iR_5U)$ and we may identify the canonical differential $d\Omega = 2iR_5U dv$. This result also follows from (5.51) upon using

$$K(q) \underset{q \rightarrow 0}{=} \frac{\pi}{2} \left(1 + \frac{q^2}{4} + \dots \right)$$

The B -period of $d\Omega$ again gives the period matrix. The corresponding integral now goes (twice) between $x = 0$ and $x = 1$, and the integral $\oint_B dv$ diverges logarithmically in the vicinity of $x = 0$. This divergence was to be expected because the period matrix contains a term τ on the right hand side (coming from the classical part of the prepotential $\mathcal{F}_{class} = -\tau a^2$) and the perturbative limit is given by $\tau \rightarrow i\infty$. Upon making the rational substitution $x = \frac{v}{1+v}$ we obtain

$$\begin{aligned} T_{pert} &= \oint_B d\Omega = \frac{U}{\pi} \int_0^1 \frac{dx}{x\sqrt{(x-1)(U^2-x)}} = \lim_{\varepsilon \rightarrow 0} \frac{U}{i\pi\sqrt{U^2-1}} \int_\varepsilon^\infty \frac{dv}{v\sqrt{v + \frac{U^2}{U^2-1}}} \\ &= -\frac{1}{i\pi} \lim_{\varepsilon \rightarrow 0} \left(\log \frac{\varepsilon}{4} \right) + \frac{1}{i\pi} \log \frac{U^2}{1-U^2} \end{aligned} \quad (5.55)$$

where ε is a small- x cut-off. Thus, the U dependent part of this integral is finite and can be considered as the “true” perturbative correction, while the divergent part just renormalizes the bare “classical” coupling constant τ . Again the same result follows from (5.53) upon using

$$\lim_{q \rightarrow 1} \left(K(q) - \frac{1}{2} \ln \left(\frac{16}{1-q^2} \right) \right) = 0.$$

The final ingredient we wish are the a -variables, i.e. the A -period of dS itself. This will correspond to the integral of (5.51) with respect to u , which is a rather complicated integral. For our purposes the perturbative limit will suffice when there are several simplifications. From the definition of (1.3) and that of dS we find

$$a = \frac{1}{R_5} \oint \frac{\log \lambda dx}{x\sqrt{x-1}} = \frac{\log \lambda|_{x=0}}{iR_5}$$

Now at $x = 0$ we have that $c_0 = i \sin \epsilon$ while at the same time $\lambda + \lambda^{-1} = c_0 u$. Together these yield

$$2 \cos aR_5 = iu \sin \epsilon$$

and

$$U^2 = \frac{u^2}{4} + \frac{1}{\sin^2 \epsilon} = \frac{1}{\sin^2 \epsilon} (1 - \cos^2 aR_5) = \frac{\sin^2 aR_5}{\sin^2 \epsilon} \quad (5.56)$$

Substituting this expression for U^2 into (5.55) we obtain for the finite part of the period matrix

$$T_{pert}^{finite} = \frac{1}{i\pi} \log \frac{\sin^2 aR_5}{\sin^2 \epsilon - \sin^2 aR_5} = \frac{1}{i\pi} \log \frac{\sin^2 aR_5}{\sin(\epsilon + aR_5) \sin(\epsilon - aR_5)}. \quad (5.57)$$

5.7 Perturbative prepotentials: adjoint matter

It is quite clear that to repeat this calculation for more general curve is a hard job. Instead, let us see what is happening with the full spectral curves under the perturbative degeneration³⁴. The results turns out to be so strikingly simple, despite quite involved form of the full spectral curve that this allows us to immediately calculate the perturbative prepotential for arbitrary $SU(N_c)$.

We again start with a warm-up example of $SU(2)$, with the curve (3.95) at the special point $\epsilon = \pi/2$. Then, upon using (5.49) we see that (3.95) reduces in the perturbative limit to

$$\cot \xi = \frac{u\lambda}{\lambda^2 + 1} \quad (5.58)$$

The rational spectral curve for this situation can be put in the form

$$w = \frac{(\lambda - \lambda_1)(\lambda - \lambda_1^{-1})}{(\lambda + \lambda_1)(\lambda + \lambda_1^{-1})}. \quad (5.59)$$

upon setting $w = -e^{2i\xi}$ and $iu = (\lambda_1 + \lambda_1^{-1})$.

Similarly in the $SU(3)$ case, in perturbative limit and at the special point $\epsilon = \pi/2$ (3.102) turns into

$$\lambda^3 - v\lambda = i(1 - u\lambda^2) \tan \xi \quad (5.60)$$

for the choice $c = -ic_0$. Equivalently,

$$i \tan \xi = \frac{\lambda(\lambda^2 + v)}{u\lambda^2 + 1}. \quad (5.61)$$

This curve can be put in the form

$$w = -\frac{(\lambda - \lambda_1)(\lambda - \lambda_2)(\lambda - \lambda_3)}{(\lambda + \lambda_1)(\lambda + \lambda_2)(\lambda + \lambda_3)}. \quad (5.62)$$

upon the identification $w = -e^{2i\xi}$ and with $u = \lambda_1 + \lambda_2 + \lambda_3$ and $v = \lambda_1\lambda_2 + \lambda_2\lambda_3 + \lambda_3\lambda_1 = \lambda_1\lambda_2\lambda_3(\lambda_1^{-1} + \lambda_2^{-1} + \lambda_3^{-1})$, $\lambda_1\lambda_2\lambda_3 = 1$.

Now let us see what is the curve at the special point where $\epsilon = \pi/2$ for the general $SU(N_c)$ case [122].

We saw already that substantial simplifications occur at the special point $\epsilon = \omega$ before even considering the perturbative limit. Now in the perturbative limit $\sigma(z) \rightarrow \sin(z)e^{z^2/6}$. Using this we easily obtain that

$$i \cot \xi = \frac{P(\lambda) - (-)^{N_c} P(-\lambda)}{P(\lambda) + (-)^{N_c} P(-\lambda)}, \quad \text{and} \quad w \equiv -e^{2i\xi} = (-)^{N_c} \frac{P(\lambda)}{P(-\lambda)}. \quad (5.63)$$

with $dS \sim \log \lambda \frac{dw}{w}$.

We finally turn to the case of generic ϵ in the perturbative limit. The system in this limit is described by the trigonometric Ruijsenaars-Schneider model. We have already seen that in this limit σ -functions are proportional to sines with exponential factors. However, the periodicity of our spectral

³⁴Similar degeneration for the Calogero curves was studied in [153].

curve means that these exponential factors must cancel amongst themselves. For example we find that in this limit

$$D_k(\xi|\epsilon) = \frac{1}{\bar{c}_-^k} \cdot \frac{\sin(\xi + (k-1)\epsilon)}{\sin(\xi - \epsilon)} \cdot \frac{(-1)^{\frac{k(k-1)}{2}}}{\sin^{k(k-2)}(\epsilon)}, \quad (5.64)$$

where $\bar{c}_-$ is given by (5.47). Now the ratio

$$\frac{\sin(\xi + (k-1)\epsilon)}{\sin(\xi - \epsilon)} = \cos k\epsilon + \sin k\epsilon \cot(\xi - \epsilon) \quad (5.65)$$

is expressible in terms of the single function $\cot(\xi - \epsilon)$. This simple observation enables us to separate the variables ξ and λ in the equation for the spectral curve upon choosing³⁵ $c = -ie^{i\epsilon}\bar{c}_-$. With $\bar{h}_k = H_k/\sin^{k(k-2)}(\epsilon)$ we may simplify the spectral curve to give

$$i \cot(\xi - \epsilon) = -\frac{\sum_{\text{odd } k}^{N_c} \bar{h}_k(-\lambda)^{N_c-k} (e^{2ik\epsilon} + 1)}{\sum_{\text{even } k}^{N_c} \bar{h}_k(-\lambda)^{N_c-k} (e^{2ik\epsilon} - 1)} = \frac{P(\lambda) + e^{2iN_c\epsilon} P(\lambda e^{-2i\epsilon})}{P(\lambda) - e^{2iN_c\epsilon} P(\lambda e^{-2i\epsilon})} \quad (5.66)$$

where $P(\lambda) = \sum_{\text{all } k}^{N_c} \bar{h}_k(-\lambda)^{N_c-k} = \prod_i^{N_c} (\lambda - e^{2ia_i})$ with some constants a_i , $\sum_i^{N_c} a_i = 0$. Introducing the variable $w = e^{2i(\xi-\epsilon)}$, one finally arrives the spectral curve in the form

$$w = e^{-2i\epsilon N_c} \frac{P(\lambda)}{P(\lambda e^{-2i\epsilon})} \quad (5.67)$$

with

$$dS \cong \log \lambda \frac{dw}{w}. \quad (5.68)$$

Thus, we have shown that our system leads in the perturbative limit to the rational spectral curve (5.67) and the generating differential (5.68). One may now calculate the corresponding prepotential using the residue formula. In fact, this calculation has been already done for the spin chains in the previous subsections, where the residue formula is applied to a general rational function of the form

$$w \sim \frac{\prod_i^{N_c} (\lambda - \lambda_i)}{\sqrt{\prod_\alpha^{N_f} (\lambda - \lambda_\alpha)}},$$

with $\lambda_i = e^{2a_i}$, $\sum_i a_i = 0$, $\lambda_\alpha = e^{2m_\alpha}$. By choosing $N_f = 2N_c$, with the hypermultiplets masses pairwise coinciding and equal to $a_i + \epsilon$ we obtain our curve (5.67). One then finds the prepotential from (5.39) gives the prepotential for the Ruijsenaars-Schneider system

$$\mathcal{F}_{\text{pert}}^R = \frac{1}{8\pi i} \sum_{i,j} f^{(5)}(R_5 a_{ij}) - \frac{1}{8\pi i} \sum_{i,j} f^{(5)}(R_5 a_{ij} + \epsilon) \quad (5.69)$$

and we are done. Further, upon setting $\epsilon = \pi/2$ we reproduce the results above.

Now in order to obtain the prepotential in the 4d theory with adjoint matter described by the Calogero-Moser system, one can just take the limit we described above, $R_5 \rightarrow 0$ and $\epsilon = MR_5$. Therefore, there is no special point $\epsilon = \frac{\pi}{2}$ in the Calogero-Moser system! The perturbative prepotential for the Calogero-Moser system [39, 154, 69] is obtain by degenerating (5.69) so that $f^{(5)}(a) \rightarrow f^{(4)}(a)$:

³⁵ Note that this separation is not in general possible non-perturbatively where there is no analogue of eq.(5.65). (This is because the ratios $\frac{\sigma(\xi+(k-1)\epsilon)}{\sigma(\xi-\epsilon)}$ transform differently under $\xi - \epsilon \rightarrow \xi - \epsilon + 1$ and $\xi - \epsilon \rightarrow \xi - \epsilon + \tau$ for different k .) The two notable exceptions are:

- (i) The case of $SU(2)$, when the variables separated for the standard choice $c = c_0$, see eq.(3.95);
- (ii) The case of $\epsilon = \pi/2$ (for any $SU(N_c)$), when separation of variables survives non-perturbatively, see eq.(3.106).

$$\mathcal{F}_{pert}^C = \frac{1}{8\pi i} \sum_{i,j} f^{(4)}(a_{ij}) - \frac{1}{8\pi i} \sum_{i,j} f^{(4)}(a_{ij} + M) \quad (5.70)$$

which can be also calculated from the spectral curve obtained by the degeneration of (5.67)

$$w = \frac{P(\lambda)}{P(\lambda - M)} \quad (5.71)$$

with the same differential (5.68).

In section 7, we shall see that this form of the perturbative prepotential persists in the Dell systems.

6 Theory of prepotential

So far we discussed only the properties of Seiberg-Witten theory related to underlying integrable systems. However, the whole construction turns out to give far more. In fact, in previous section we just discussed how *to construct* different physical data in terms of integrable systems, but almost missed *meaning* of several key notions. Maybe the main object that is to be discussed in far more details is the prepotential, the quantity not properly interpreted yet in previous sections.

Indeed, looking from the integrable standing point, one should find natural both emerging the Riemann surfaces with their moduli space as the corresponding spectral curves, as well as the proper coordinates on the moduli space given by the action integrals (1.3). However, the notion of the prepotential which allows one to make quantitative predictions and defines the low-energy effective action was so far introduced by hands, just via formula (1.3). This formula implies that not only A - but also B -periods of the generating differential dS should get a proper interpretation. This means that one should better understand meaning of dS and its properties. One has particularly to recover the reason for dS to give rise to holomorphic differentials, or, to put it differently, the reason for the second derivative of the prepotential to be a period matrix.

In this section we are going to recover another kind of integrability, Whitham integrability which stands behind the notion of the prepotential and generating differential dS and discuss several very essential its features, in particular, the WDVV equations (1.8).

6.1 From Seiberg-Witten theory to Whitham hierarchies: RG dynamics

Let us consider a generic recipe how the generating differential is introduced in integrable systems [4, 5, 155]. It is essentially based on the analogue of the renormalization group flows in field theory, where one performs the integration over heavy modes giving rise to the effective action for light modes. Within the framework of integrable many-body systems, an analogous procedure looks as follows. First, one should start with a solution to the equations of motion with moduli. At the next step, one assumes that the moduli adiabatically depend on some new time variables. In this way, one obtains *the Whitham dynamics* on the moduli space, with the generating differential dS playing the role of action differential and the prepotential being logarithm of the τ -function. For instance, for the finite-gap solutions that are in charge of supersymmetric gauge theories without adjoint matter, i.e. those described by hyperelliptic curves, it is natural to choose the corresponding branching points as these slowly evolving moduli.

Therefore, we come to the conclusion [37] that Seiberg-Witten theory has much to do with *two* different structures: with finite-gap solutions of integrable systems of the Hitchin type or of spin chains

(so that the integrals of motion are fixed and related to the vacuum expectation values (vev's) of scalar fields in supersymmetric theories) *and* with the corresponding *Whitham* hierarchy constructed in the vicinity of these solutions and describing the evolution of integrals of motion.

Since the vev's of scalar fields physically depend on the scale Λ_{QCD} , one associates one of the Whitham evolutions with the RG dynamics w.r.t. this scale (in fact, Λ_{QCD} is identified with the first Whitham time). Other slow (Whitham) times describes vev-dependence on perturbations of the original SYM action by different operators (from topological sector). To put it differently, while Seiberg-Witten theory describes the low-energy effective action, the Whitham dynamics governs the behaviour of correlators at low energies. This correlators are from the topological subsector of the theory, since only such correlators survive at low energies. Note that this subsector is manifestly described by the Donaldson theory, [156, 157, 158].

In fact, this is the prepotential that is the generating function of these (connected) topological correlators, i.e. logarithm of the corresponding partition function. Therefore, now the prepotential depends on both the integrals of motion and the Whitham times as on *independent* variables [82]. Before defining such an object, let us briefly review the very framework of Whitham hierarchies, from the very general point of view.

Following I.Krichever [4, 155], we consider initially a *local* system of holomorphic functions Ω_I on complex curve, i.e. functions of some local parameter ξ in a neighborhood of a point P . One can introduce a set of parameters t_I and define on the space (ξ, t_I) a 1-form

$$\omega = \sum_I \Omega_I(P, \mathbf{t}) \delta t_I \quad (6.1)$$

Considering its total external derivative $\delta\omega = \sum_I \delta\Omega_I(P, \mathbf{t}) \wedge \delta t_I$, where $\delta\omega = \partial_\xi \Omega_I \delta\xi \wedge \delta t_I + \partial_J \Omega_I \delta t_I \wedge \delta t_J$, one can define general *Whitham equations* as

$$\delta\omega \wedge \delta\omega = 0 \quad (6.2)$$

so that it is necessary to check the independent vanishing of two different terms: δt^4 and $\delta t^3 \delta\xi$. The second term gives

$$\sum \partial_\xi \Omega_{[I} \partial_J \Omega_{C]} = 0 \quad (6.3)$$

where [...] means antisymmetrization. Fixing now some $I = I_0$ and introducing the “Darboux” coordinates

$$t_{I_0} \equiv x \quad \Omega_{I_0}(P, \mathbf{t}) \equiv p \quad (6.4)$$

one can get from (6.3) the Whitham equations in their standard form [155]

$$\begin{aligned} \partial_I \Omega_J - \partial_J \Omega_I + \{\Omega_I, \Omega_J\} &= 0 \\ \{\Omega_I, \Omega_J\} &\equiv \frac{\partial \Omega_I}{\partial x} \frac{\partial \Omega_J}{\partial p} - \frac{\partial \Omega_J}{\partial x} \frac{\partial \Omega_I}{\partial p} \end{aligned} \quad (6.5)$$

The explicit form of the equations (6.5) strongly depends on choice of the local co-ordinate p . Equations (6.2), (6.3) and (6.5) are defined only locally and have a huge amount of solutions.

Now let us note that the Whitham equations (6.5) can be considered as compatibility conditions of the system

$$\frac{\partial \lambda}{\partial t_I} = \{\lambda, \Omega_I\} \quad (6.6)$$

The function λ can be used itself in order to define a new local parameter. In this co-ordinate, the system (6.5) turns into [155]

$$\left. \frac{\partial \Omega_I(\lambda, t)}{\partial t_J} \right|_{\lambda=const} = \left. \frac{\partial \Omega_J(\lambda, t)}{\partial t_I} \right|_{\lambda=const} \quad (6.7)$$

or for the differentials $d\Omega_I \equiv d_\Sigma \Omega_I = \frac{\partial \Omega_I}{\partial \lambda} d\lambda$

$$\left. \frac{\partial d\Omega_I(\lambda, t)}{\partial t_J} \right|_{\lambda=const} = \left. \frac{\partial d\Omega_J(\lambda, t)}{\partial t_I} \right|_{\lambda=const} \quad (6.8)$$

The Whitham equations in the form (6.7) imply the existence of a potential S such that

$$\Omega_I(\lambda, t)|_{\lambda=const} = \left. \frac{\partial S(\lambda, t)}{\partial t_I} \right|_{\lambda=const} \quad (6.9)$$

With this potential, the 1-form ω (6.1) can be rewritten as

$$\omega = \delta S - \frac{\partial S(\lambda, t)}{\partial \lambda} d\lambda \quad (6.10)$$

To fill all these formulas with a real content, one can consider interesting and important examples of solutions to the Whitham hierarchy arising when one takes the basis functions Ω_I to be the “quasi-energies” of the finite-gap solution; these are the (globally multi-valued) functions whose periods determine the “phase” of the quasi-periodic Baker-Akhiezer function corresponding to this finite-gap solution. The potential should be then identified with

$$S = \int^P dS \quad (6.11)$$

$$\frac{\partial S}{\partial t_I} = \int^P \frac{\partial dS}{\partial t_I} = \int^P d\Omega_I = \Omega_I$$

where dS is, as usual, the generating differential for the finite gap system. Usually for KP/Toda hierarchy these functions are taken to be “half”-multi-valued, i.e. their $\mathbf{A}$ -periods are fixed to be zero: $\oint_{\mathbf{A}} d\Omega_I = 0$. The equations (6.11) are obviously satisfied in a trivial way in the case of the (finite-gap solutions to) KP/Toda theory, when $\{t_I\}$ are taken to be *external* parameters and $\{\Omega_I\}$ do not depend themselves on $\{t_I\}$. The idea, however, is to deform this trivial solution to (6.2), (6.3) and (6.5) into a nontrivial one by a *flow* in the moduli space. Practically it means that the formulas (6.11) should be preserved even when $\{\Omega_I\}$ *depend* on $\{t_I\}$, but in a special way – determined by the equations of Whitham hierarchy. In other words, the Whitham equations correspond to the dynamics on moduli space of complex curves and the moduli become depending on the Whitham times.

To realize all this technically, one choose $d\Omega_I$ to be a set of (normalized) differentials on the complex curve holomorphic outside the marked points where they have some fixed behavior. These differentials depend on Whitham times only through moduli dependence.

In fact, for this whole construction to work (i.e. for the Whitham hierarchy to have some non-trivial solutions), one needs to choose both differentials (with their normalization) and local parameter in a clever way. This is the thing to be added to Seiberg-Witten theory. Indeed, Seiberg-Witten theory is given by the set of data that includes Riemann surfaces, their moduli space and the differential dS ,

while the Whitham set-up additionally includes a (non-invariant geometrically) proper choice of the local parameter.

An important example of the λ co-ordinate is the *hyperelliptic* co-ordinate on the hyperelliptic curve (corresponding to the KdV equation)

$$y^2 = R(\lambda) = \prod_{\beta}^{2g+2} (\lambda - r_{\beta}) \quad (6.12)$$

Properly choosing the differentials $d\Omega_I$ and expanding the equations (6.8) near the point $\lambda = r_{\alpha}$ one gets the Whitham equations over the finite-gap solutions of the KdV equation

$$\frac{\partial r_{\alpha}}{\partial t_I} = \frac{d\Omega_J}{d\Omega_I} \Big|_{\lambda=r_{\alpha}} \frac{\partial r_{\alpha}}{\partial t_J} \equiv v_{IJ}^{(\alpha)}(\mathbf{r}) \frac{\partial r_{\alpha}}{\partial t_J} \quad (6.13)$$

The set of ramification points $\{\mathbf{r}\} = \{r_{\beta}\}$ is usually called Riemann invariants for the Whitham equations.

The particular case of Whitham hierarchy associated with the finite-gap solution of the KdV hierarchy described by the hyperelliptic curve

$$y^2 = R(\lambda) = \prod_{\beta}^{2g+1} (\lambda - r_{\beta}) \quad (6.14)$$

was considered first in [159].

Note also that in the framework of Whitham hierarchy it becomes possible to introduce a counterpart of the tau-function, again a generating function for the solutions, which can be symbolically defined as [155]

$$\log \mathcal{T}_{\text{Whitham}} = \int_{\Sigma} \bar{d}S \wedge dS \quad (6.15)$$

where the two-dimensional integral over Σ is actually “localized” on the non-analyticities of S . In fact, this $\log \mathcal{T}_{\text{Whitham}}$ is nothing but the prepotential.

Recently an analogue of Whitham dynamics for deformations of the analytic curves instead of the complex surfaces has been found [160, 161]. At the language of dynamical systems it means that the dynamics in the space of real Hamiltonians instead of holomorphic ones is considered.

6.2 Whitham hierarchy for the Toda chain

Now let us discuss how one can choose the differentials $d\Omega_I$ in order to get non-trivial solutions to Whitham hierarchy. They can be defined for the following set of data:

- complex curve (Riemann surface) of genus g ;
- a set of punctures P_i (marked points);
- co-ordinates ξ_i in the vicinities of the punctures P_i ;
- pair of differentials, say, $(d\lambda, dz)$ with fixed periods.

We start with considering the simplest case of a single puncture, say at $\xi = \xi_0 = 0$. This situation is typical, e.g., for the Whitham hierarchy in the vicinity of the finite-gap solutions to the KP or KdV hierarchies. For instance, in the KdV case the spectral curve has the form (6.14) so that the marked point is at $\lambda = \infty$ but, in contrast to the Toda curve (3.8), there is only one infinite point (since it is the ramification point).

Given this set of data, one can introduce meromorphic differentials with the poles only at some point P_0 such that in some local co-ordinate ξ : $\xi(P_0) = \xi_0 = 0$

$$d\Omega_n \underset{P \rightarrow P_0}{=} (\xi^{-n-1} + O(1)) d\xi, \quad n \geq 1 \quad (6.16)$$

This condition defines $d\Omega_n$ up to arbitrary linear combination of g holomorphic differentials $d\omega_i$, $i = 1, \dots, g$ and there are two different natural ways to fix this ambiguity. The first way already mentioned above is to require that $d\Omega_n$ have vanishing A -periods,

$$\oint_{A_i} d\Omega_n = 0 \quad \forall i, n \quad (6.17)$$

Their generating functional ($\zeta \equiv \xi(P')$, $\xi \equiv \xi(P)$)

$$W(\xi, \zeta) = \sum_{n=1}^{\infty} n \zeta^{n-1} d\zeta d\Omega_n(\xi) + \dots \quad (6.18)$$

is well known in the theory of Riemann surfaces. It can be expressed through the Prime form $E(P, P')$ (see Appendix A of [82] for details):³⁶

$$W(P, P') = \partial_P \partial_{P'} \log E(P, P') \quad (6.19)$$

Such $W(P, P')$ has a second order pole on diagonal $P \rightarrow P'$,

$$W(\xi, \zeta) \sim \frac{d\xi d\zeta}{(\xi - \zeta)^2} + O(1) = \sum_{n=1}^{\infty} n \frac{d\xi}{\xi^{n+1}} \zeta^{n-1} d\zeta + O(1) \quad (6.20)$$

It is the differentials (6.16) that should be related with the potential differential dS by

$$d\Omega_n = \frac{\partial dS}{\partial t_n} \quad (6.21)$$

The second way to normalize differentials is to impose the condition

$$\frac{\partial d\widehat{\Omega}_n}{\partial \text{moduli}} = \text{holomorphic} \quad (6.22)$$

so that these differentials becomes similar to the generating differential dS_{SW} of integrable system.

³⁶ For example, for genus $g = 1$ in the co-ordinate $\xi \sim \xi + 1 \sim \xi + \tau$ the formulas acquire the form

$$d\Omega_1 = (\wp(\xi) - \text{const}) d\xi, \quad d\Omega_2 = -\frac{1}{2} \wp'(\xi) d\xi, \quad \dots, \quad d\Omega_n = \frac{(-)^{n+1}}{n!} \partial^{n-1} \wp(\xi) d\xi$$

and

$$W(\xi, \xi') = \sum_{n=1}^{\infty} \frac{(-)^{n+1} \xi'^{n-1}}{(n-1)!} \frac{\partial^{n-1}}{\partial \xi^{n-1}} \wp(\xi) d\xi d\xi' - \text{const} \cdot d\xi d\xi' = \wp(\xi - \xi') d\xi d\xi' = \partial_\xi \partial_{\xi'} \log \theta_*(\xi - \xi')$$

where $*$ denotes the (on torus the only one) odd theta-characteristic. For $g = 1$ $\nu_*^2(\xi) = \theta_{*,i}(0) d\omega_i$ is just $d\xi$. Let us also point out that chosen in this way co-ordinate ξ is not convenient from the point of view of Whitham hierarchy since its “periods” ($\tau = \oint_B d\xi$) depend on moduli of the curve.

Now let us specialize this description to the simplest case of Seiberg-Witten theory corresponding to the $4d$ pure gauge theory, i.e. to the Toda chain. Therefore, we deal with the family of spectral curves (3.6). The first problem is that naively there are no solutions to eq.(6.22) because the curves (3.6) are spectral curves of the *Toda-chain* hierarchy (not of KP/KdV type). The difference is that the adequate description in the Toda case requires *two* punctures instead of one. As already mentioned in sect.3.3, the curves (3.6) have two marked points and there exists a function w with the N_c -degree pole and zero at two (marked) points $\lambda = \infty_{\pm}$, where $\pm$ labels two sheets of the hyperelliptic representation (3.6), $w(\lambda = \infty_+) = \infty$, $w(\lambda = \infty_-) = 0$. Accordingly, there are two families of the differentials $d\Omega_n$: $d\Omega_n^+$ with the poles at ∞_+ and $d\Omega_n^-$ with the poles at ∞_- .

However, there are *no* differentials $d\hat{\Omega}^{\pm}$, only $d\hat{\Omega}_n = d\hat{\Omega}_n^+ + d\hat{\Omega}_n^-$, i.e. condition (6.22) requires $d\hat{\Omega}_n$ to have the poles at both punctures. Moreover, the coefficients in front of w^{n/N_c} at ∞_+ and w^{-n/N_c} at ∞_- (6.16) coincide (in Toda-hierarchy language, this is the Toda-chain case with the same dependence upon negative and positive times).

The differentials $d\hat{\Omega}_n$ for the family (3.6) have the form [82]:

$$d\hat{\Omega}_n = R_n(\lambda) \frac{dw}{w} = P_+^{n/N_c}(\lambda) \frac{dw}{w} \quad (6.23)$$

The polynomials $R_n(\lambda)$ of degree n in λ are defined by the property that $P'\delta R_n - R_n'\delta P$ is a polynomial of degree less than $N_c - 1$. Thus, $R_n(\lambda) = P_+^{n/N_c}(\lambda)$, where $\left(\sum_{k=-\infty}^{+\infty} c_k \lambda^k\right)_+ = \sum_{k=0}^{+\infty} c_k \lambda^k$. For example³⁷:

$$\begin{aligned} R_0 &= 1, \\ R_1 &= \lambda, \\ R_2 &= \lambda^2 - \frac{2}{N_c} u_2, \\ R_3 &= \lambda^3 - \frac{3}{N_c} u_2 \lambda - \frac{3}{N_c} u_3, \\ R_4 &= \lambda^4 - \frac{4}{N_c} u_2 \lambda^2 - \frac{4}{N_c} u_3 \lambda - \left(\frac{4}{N_c} u_4 + \frac{2(N_c - 4)}{N_c^2} u_2^2 \right), \\ &\dots \end{aligned} \quad (6.24)$$

These differentials satisfy (6.22) provided the moduli-derivatives are taken at constant w (not λ !). Thus, the formalism of the previous section is applicable for the local parameter $\xi = w^{\mp 1/N_c}$.³⁸

Thus, among the data one needs for the definition of solution to the Whitham hierarchy associated with the Toda chain, there are the punctures at $\lambda = \infty_{\pm}$ and the relevant co-ordinates in the vicinities of these punctures $\xi \equiv w^{-1/N_c} \sim \lambda^{-1}$ at ∞_+ and $\xi \equiv w^{+1/N_c} \sim \lambda^{-1}$ at ∞_- . The parameterization in terms of w , however, does not allow one to use the advantages of hyperelliptic form (3.8).

³⁷In the case of the Toda chain, since there are two punctures, one can consider the meromorphic differential with the simple pole at each of them.

³⁸Rule of thumb generally is to choose as the local co-ordinate for the Whitham hierarchy the parameter living on the bare spectral curve. For the Toda case, one may choose as a bare curve the w -cylinder with the corresponding generating differential $dS = \lambda d \log w$.

6.3 Solution to Whitham hierarchy and prepotential

Now we are going to extend the system of differentials $d\Omega_i$ introduced in the previous section to include the holomorphic differentials. This will allow us to construct the function that, as a function of one set of variables, is the prepotential of the Seiberg-Witten theory (as defined in sect.2) and in the other set of *independent* variables is the (logarithm of) τ -function $\mathcal{T}_{\text{Whitham}}$ of the associated solution to the Whitham hierarchy. On this way, we also get a generic solution to the associated Whitham equations.

Thus, we extend the set of Whitham equations by the equations involving the holomorphic differentials $d\omega_i$ and moduli α_i :

$$\frac{\partial d\Omega_n}{\partial \alpha_i} = \frac{\partial d\omega_i}{\partial t_n}, \quad \frac{\partial d\omega_i}{\partial \alpha_j} = \frac{\partial d\omega_j}{\partial \alpha_i} \quad (6.25)$$

This system is solved by the differential dS that satisfies

$$\frac{\partial dS}{\partial \alpha_i} = d\omega_i, \quad \frac{\partial dS}{\partial t_n} = d\Omega_n \quad (6.26)$$

Then, the first equation in (6.26) implies that dS is to be looked for as a linear combination of the differentials $d\hat{\Omega}_n$, satisfying (6.22). Let us, following [90, 91, 82] introduce a generating functional for $d\hat{\Omega}_n$ with infinitely many auxiliary parameters t_n :

$$dS = \sum_{n \geq 1} t_n d\hat{\Omega}_n = \sum_{i=1}^g \alpha_i d\omega_i + \sum_{n \geq 1} t_n d\Omega_n \quad (6.27)$$

The periods

$$\alpha_i = \oint_{A_i} dS \quad (6.28)$$

can be considered as particular co-ordinates on the moduli space. Note that these periods do not exactly coincide with (1.3), eq.(6.28) defines α_i as functions of h_k and t_n , or, alternatively, h_k as functions of α_i and t_n so that derivatives $\partial h_k / \partial t_n$ are non-trivial. In what follows we shall consider α_i and t_n ,

$$t_n = \text{res}_{\xi=0} \xi^n dS(\xi) \quad (6.29)$$

as independent variables so that partial derivatives w.r.t. α_i are taken at constant t_n and partial derivatives w.r.t. t_n are taken at constant α_i .

The differential dS (6.27) determines a generic form of the solution associated to Seiberg-Witten type Whitham hierarchy. The Whitham dynamics itself for given dS can be formulated in terms of equations (6.26). Note that, if one restricts himself to the Whitham hierarchy with several first times (generally, for the genus g complex curve there are $g + n - 1$ independent times, with n being the number of punctures), all higher times in (6.27) play the role of constants (parameters) of generic Whitham solution (see example of $SU(2)$ below).

Note that the Seiberg-Witten differential dS_{SW} is $dS_{SW} = d\hat{\Omega}_2$, i.e.

$$dS|_{t_n=\delta_{n,1}} = dS_{SW}, \quad \alpha^i|_{t_n=\delta_{n,1}} = a^i, \quad \alpha_i^D|_{t_n=\delta_{n,1}} = a_i^D \quad (6.30)$$

and α -variables are naturally associated with the Seiberg-Witten moduli, while t -variables – with the corresponding Whitham times (although α_i 's can be also considered as variables of Whitham dynamics, cf. with (6.25)).

Now one can introduce the Whitham tau-function (6.15) whose logarithm is a *prepotential* $\mathcal{F}(\alpha_i, t_n) \equiv \log \mathcal{T}_{\text{Whitham}}$ by an analog of conditions (1.3):

$$\frac{\partial \mathcal{F}}{\partial \alpha_i} = \oint_{B_i} dS, \quad \frac{\partial \mathcal{F}}{\partial t_n} = \frac{1}{2\pi i n} \text{res}_0 \xi^{-n} dS \quad (6.31)$$

Their consistency follows from (6.26) and Riemann identities. In particular,

$$\frac{\partial^2 \mathcal{F}}{\partial t_m \partial t_n} = \frac{1}{2\pi i n} \text{res}_0 \xi^{-n} \frac{\partial dS}{\partial t_m} = \frac{1}{2\pi i n} \text{res}_0 \xi^{-n} d\Omega_m = \frac{1}{2\pi i m} \text{res}_0 \xi^{-m} d\Omega_n \quad (6.32)$$

This calculation implies that the definition (6.31) assumes the co-ordinate ξ is not changed under the variation of moduli. It means that this provides a moduli-independent parameterization of the entire family – like w in the case of (3.6). Since moduli-independence of ξ should be also consistent with (6.22), the choice of ξ is strongly restricted: to $w^{\pm 1/N_c}$ in the case of (3.6) (see the discussion in the previous section).

The last relation in (6.32) (symmetricity) is just an example of the Riemann relations and it is proved by the standard argument:

$$\begin{aligned} 0 &= \int d\Omega_m \wedge d\Omega_n = \oint_{A_i} d\Omega_m \oint_{B_i} d\Omega_n - \oint_{A_i} d\Omega_n \oint_{B_i} d\Omega_m + \frac{1}{2\pi i} \text{res} (d\Omega_m d^{-1} d\Omega_n) = \\ &= 0 + 0 + \frac{1}{2\pi i n} \text{res}_0 \xi^{-n} d\Omega_m - \frac{1}{2\pi i m} \text{res}_0 \xi^{-m} d\Omega_n \end{aligned} \quad (6.33)$$

where (6.17) and (6.16) are used at the final stage. The factors like n^{-1} arise since $\xi^{-n-1} d\xi = -d(\xi^{-n}/n)$. We shall also use a slightly different normalization $d\Omega_n \sim w^{n/N_c} \frac{dw}{w} = \frac{N_c}{n} dw^{n/N_c}$, accordingly the residues in (6.33) and (6.31) and (6.32) will be multiplied by N_c/n instead of $1/n$.

By definition, the prepotential is a homogeneous function of its arguments a_i and t_n of degree 2,

$$2\mathcal{F} = \alpha_i \frac{\partial \mathcal{F}}{\partial \alpha_i} + t_n \frac{\partial \mathcal{F}}{\partial t_n} = \alpha_i \alpha_j \frac{\partial^2 \mathcal{F}}{\partial \alpha_i \partial \alpha_j} + 2\alpha_i t_n \frac{\partial^2 \mathcal{F}}{\partial \alpha_i \partial t_n} + t_m t_n \frac{\partial^2 \mathcal{F}}{\partial t_m \partial t_n} \quad (6.34)$$

Again, this condition can be proved with the help of Riemann identities, starting from (6.31), (6.28) and (6.29). At the same time, $\mathcal{F}$ is not just a quadratic function of a_i and t_n , a non-trivial dependence on these variables arise through the dependence of $d\omega_i$ and $d\Omega_n$ on moduli (like u_k or h_k) which in their turn depend on a_i and t_n . This dependence is obtained, for example, by substitution of (6.27) into (6.26):

$$d\hat{\Omega}_n + t_m \frac{\partial d\hat{\Omega}_m}{\partial u_l} \frac{\partial u_l}{\partial t_n} = d\Omega_n, \quad (6.35)$$

i.e.

$$\left(\sum_{m,l} t_m \frac{\partial u_l}{\partial t_n} \right) \oint_{A_i} \frac{\partial d\hat{\Omega}_m}{\partial u_l} = - \oint_{A_i} d\hat{\Omega}_n \quad (6.36)$$

The integral in the l.h.s. is expressed, according to (6.22), through the integrals of holomorphic 1-differentials, while the integral in the r.h.s. – through the periods of dS . If

$$\oint_{A_i} \frac{\partial dS}{\partial u_l} = \oint_{A_i} dV_l = \Sigma_{il} \quad (6.37)$$

then

$$t_m \frac{\partial u_k}{\partial t_m} = \Sigma_{ki}^{-1} \alpha_i \quad (6.38)$$

The relations (6.38) can be thought of as the other form of the Whitham hierarchy.

Thus, we have constructed the prepotential for the Whitham hierarchy associated with Seiberg-Witten theory. The Whitham equations themselves describe the dependence of effective coupling constants given by the period matrix of the Seiberg-Witten curve on bare coupling constants t_n . The prepotential also depends on characteristics of the vacuum parametrized by vev's u_k . Using this interpretation, one expects [82] that the constructed Whitham prepotential has to be associated with the generating function of topological correlators, the Whitham times being bare coupling constants of different local operators. It issues the problem of calculating derivatives of the prepotential with their further comparison with the topological correlators. Indeed, the first two derivatives have been calculated in [82] and turned out to be

$$\frac{\partial \mathcal{F}}{\partial t^n} = \frac{\beta}{2\pi i n} \sum_m m t_m \mathcal{H}_{m+1, n+1} = \frac{\beta}{2\pi i n} t_1 \mathcal{H}_{n+1} + O(t_2, t_3, \dots) \quad (6.39)$$

$$\frac{\partial^2 \mathcal{F}}{\partial \alpha^i \partial t^n} = \frac{\beta}{2\pi i n} \frac{\partial \mathcal{H}_{n+1}}{\partial a^i} \quad (6.40)$$

$$\frac{\partial^2 \mathcal{F}}{\partial t^m \partial t^n} = -\frac{\beta}{2\pi i} \left(\mathcal{H}_{m+1, n+1} + \frac{\beta}{m n} \frac{\partial \mathcal{H}_{m+1}}{\partial a^i} \frac{\partial \mathcal{H}_{n+1}}{\partial a^j} \partial_{ij}^2 \log \theta_E(\vec{0}|T) \right) \quad (6.41)$$

etc. In these formulas parameter $\beta = 2N_c$, $m, n = 1, \dots, N_c - 1$ and $\mathcal{H}_{m, n}$ are certain homogeneous combinations of u_k , defined in terms of $P_N(\lambda)$:

$$\mathcal{H}_{m+1, n+1} = -\frac{N_c}{m n} \text{res}_\infty \left(P^{n/N_c}(\lambda) dP_+^{m/N_c}(\lambda) \right) = \mathcal{H}_{n+1, m+1} \quad (6.42)$$

and

$$\mathcal{H}_{n+1} \equiv \mathcal{H}_{n+1, 2} = -\frac{N_c}{n} \text{res}_\infty P^{n/N_c}(\lambda) d\lambda \quad (6.43)$$

The Seiberg-Witten theory itself allows one to evaluate derivatives (6.39), (6.41) at $n = 1$, since after appropriate rescaling

$$h_k \rightarrow t_1^k h_k, \quad \mathcal{H}_k \rightarrow t_1^k \mathcal{H}_k \quad (6.44)$$

T_1 can be identified with Λ_{QCD} . Note that, for doing this, one needs to use the Toda curve in the form (3.7). Eqs. (6.39), (6.41) at $n = 1$ are naturally interpreted in terms of the stress-tensor anomaly,

$$\dots + \vartheta^4 \Theta_\mu^\mu = \beta \text{tr} \Phi^2 = \dots + \vartheta^4 \beta \text{tr} \left(G_{\mu\nu} G^{\mu\nu} + i G_{\mu\nu} \tilde{G}^{\mu\nu} \right), \quad (6.45)$$

since for any operator $\mathcal{O}$

$$\frac{\partial}{\partial \log \Lambda} \langle \mathcal{O} \rangle = \langle \beta \text{Tr} \Phi^2, \mathcal{O} \rangle \quad (6.46)$$

Analogous interpretation for $n \geq 2$ involves anomalies of W_{n+1} -structures.

For $n = 1$, eq.(6.39), i.e. eq.(6.46) for $\mathcal{O} = I$, has been derived in [162] in the form

$$\frac{\partial \mathcal{F}_{SW}}{\partial \log \Lambda_{QCD}} = \frac{\beta}{2\pi i} (t_1^2 h_2) \quad (6.47)$$

Eq.(6.46) for $\mathcal{O} = h_m$, the analogue of (6.41) for $n = 1$ and any $m = 1, \dots, N_c - 1$, is

$$\frac{\partial h_m}{\partial \log \Lambda_{QCD}} = -\beta \frac{\partial h_2}{\partial a^i} \frac{\partial h_m}{\partial a^j} \partial_{ij}^2 \log \theta_E(\vec{0}|T) \quad (6.48)$$

Formulas of such a type first appeared for the $SU(2)$ case in [156], and literally in this form for the $SU(2)$ case in [157, 158] and for the general $SU(N_c)$ case in [157]. It was deduced from its modular properties and also from sophisticated reasoning in the framework of the Donaldson theory (the topological correlator is an object from the Donaldson theory, i.e. topological subsector of the $\mathcal{N} = 2$ SUSY Yang-Mills theory, which can be investigated with the help of the twisting procedure). In (6.48), one can substitute h_m by any homogeneous function of h 's, e.g. by $\mathcal{H}_m$.

Eq. (6.41) can be also rewritten as:

$$\frac{\partial^2}{\partial t^m \partial t^n} \left(\mathcal{F}(\alpha, t) - \frac{\beta^2}{4\pi i} \mathcal{F}^{GKM}(\alpha, t) \right) = -\frac{\beta^2}{2\pi i m n} \frac{\partial \mathcal{H}_{m+1}}{\partial a^i} \frac{\partial \mathcal{H}_{n+1}}{\partial a^j} \partial_{ij}^2 \log \theta_E(\vec{0}|T) \quad (6.49)$$

where the prepotential of the Generalized Kontsevich Model [163],

$$\mathcal{F}^{GKM}(\alpha|t) \equiv \frac{1}{2N} \sum_{m,n} t_m t_n \mathcal{H}_{m+1, n+1} \quad (6.50)$$

implicitly depends on the moduli α^i through the coefficients of the polynomial $P(\lambda)$.

Here we considered only the simplest example of the 4d pure gauge theory. This approach was generalized to the theories with fundamental matter and with other gauge groups in [149, 164, 165]. Further development on the connections between Whitham hierarchies, τ -functions of integrable hierarchies of the KP/Toda type and Donaldson topological theories can be found in [166]. These latter, very interesting results are still waiting to be re-interpreted in spirit of the approach of [163] to the Generalized Kontsevich Model.

As far as we know, the adjoint matter case has not been done yet.

6.4 Whitham hierarchy for $SU(2)$: an example

To conclude our discussion of Whitham hierarchies, let us illustrate the scheme developed above in the simplest case of 2-site periodic Toda chain³⁹. One can restrict himself to the first two differentials $d\Omega_n$ and the first two times t_0, t_1 .⁴⁰ Then, the Whitham equations

$$\frac{\partial d\Omega_i}{\partial t_j} = \frac{\partial d\Omega_j}{\partial t_i} \quad (6.51)$$

are reduced to the only equation

³⁹We do not discuss here how to calculate the second derivative of the prepotential in this case, since it is not too illuminating. The details can be found in Appendix B of [82].

⁴⁰It can be done since the higher flows are dependent on the two first flows (or, which is the same, u depends only on two independent combinations of times). Generically, for a complex curve of genus g , there are $g + 1$ independent flows.

$$\frac{\partial d\Omega_0}{\partial t_1} = \frac{\partial d\Omega_1}{\partial t_0} \quad (6.52)$$

To write it down explicitly one should remember that

- There are two independent differentials (see (6.23) and (6.24))

$$\begin{aligned} d\Omega_0 &= dz + \gamma_0 \frac{d\lambda}{y} = \left(1 + \frac{\gamma_0}{2\lambda}\right) dz \\ d\Omega_1 &= \lambda dz + \gamma_1 \frac{d\lambda}{y} = \left(\lambda + \frac{\gamma_1}{2\lambda}\right) dz \end{aligned} \quad (6.53)$$

and two corresponding Whitham times t_0 and t_1 . Here we introduced the notation $dz \equiv dw/w$.

- The coefficients γ_i ($i = 0, 1$) are fixed, as usual, by vanishing of the correspondent A -periods

$$\oint_A d\Omega_i = 0 \quad (6.54)$$

i.e.

$$\gamma_1 = -\frac{a}{\sigma}, \quad \gamma_0 = -\frac{1}{\sigma} \quad (6.55)$$

where

$$a = \oint_A dS = \oint_A \lambda dz, \quad \sigma = \oint_A \frac{d\lambda}{y} = \frac{\partial a}{\partial u} \quad (6.56)$$

- The derivatives over moduli in (6.51) are taken at fixed values of z -variable, while the λ -variable depends on moduli and this dependence is given by (3.18). This rule leads to the relations

$$\frac{\partial \lambda}{\partial u} = \frac{1}{2\lambda}, \quad \frac{\partial a}{\partial u} = \oint_A \frac{\partial \lambda}{\partial u} dz = \oint_A \frac{dz}{2\lambda} = \sigma \quad (6.57)$$

Using these relations one easily gets from (6.51) the following Whitham equation on moduli

$$\frac{\partial u}{\partial t_1} = a(u) \frac{\partial u}{\partial t_0} \quad (6.58)$$

The function $a(u)$ is an elliptic integral. Its explicit expression depends on choice of the cycles. With the choice used in [82], i.e. when the $\mathbf{A}$ -cycles encircle the points $\lambda = \sqrt{u - 2\Lambda^2} \equiv r^-$ and $\lambda = \sqrt{u + 2\Lambda^2} \equiv r^+$,

$$a = \frac{2}{\pi} r^+ E(k), \quad \sigma = \frac{1}{\pi r^+} K(k) \quad (6.59)$$

where $K(k)$ and $E(k)$ are complete elliptic integrals of the first and the second kinds respectively and the elliptic modulus is $k = \frac{2}{r^+}$.

The Whitham equation (6.58) has the general solution:

$$u = F(t_0 + a(u)t_1) \quad (6.60)$$

where $F(x)$ is an arbitrary function. This solution can be also rewritten in the form

$$\Phi(u) = t_0 + a(u)t_1 \quad (6.61)$$

where $\Phi(u)$ is the function inverse to $F(x)$.

Thus, we constructed the solution to the Whitham hierarchy just straightforwardly. Now we do the same, using the solution for the differential dS (6.27) found in the previous subsection. We start first with only two first non-zero times so that dS has a particular form

$$dS = t_0 d\widehat{\Omega}_0 + t_1 d\widehat{\Omega}_1 = t_0 dz + t_1 \lambda dz \quad (6.62)$$

Then, using formulas of the end of the previous section for the differentials $d\Omega_{1,2}$, one can easily obtain from (6.35) for $n = 0, 1$ the following two equations:

$$\begin{aligned} t_1 \frac{\partial u}{\partial t_1} &= -\frac{a}{\sigma}, \\ t_1 \frac{\partial u}{\partial t_0} &= -\frac{1}{\sigma} \end{aligned} \quad (6.63)$$

Any solution to the equations (6.63) evidently solves simultaneously (6.58), but not *vice versa*.

Let us use the manifest form of functions $a(u)$ and $\sigma(u)$ from (6.59):

$$\begin{aligned} t_1 \frac{\partial u}{\partial t_1} &= -\frac{8}{k^2} \frac{E(k)}{K(k)}, \\ t_1 \frac{\partial u}{\partial t_0} &= -\frac{2\pi}{k} \frac{1}{K(k)} \end{aligned} \quad (6.64)$$

Then, the first equation of (6.63) has the solution [167]

$$t_1 = c(t_0) \frac{k}{E(k)} \quad (6.65)$$

with arbitrary function $c(t_0)$. The solution to the both equations (6.63) takes the form

$$a(u) = \frac{\text{const} - t_0}{t_1} \quad (6.66)$$

This particular solution has to be compared with (6.61) with the constant function $\Phi(u)$ (since it has no inverse, the second form (6.60) does not exist for this concrete solution).

In order to get the general solution (6.61) to the Whitham hierarchy, one has to require for all higher odd times in (6.27) (i.e. with $n > 1$) to be non-zero constants. Then, these higher times parameterize the general function $\Phi(u)$. Indeed, how it can be easily obtained from (6.35), the solution to the Whitham equations with general dS (6.27) is given by formula (6.61) with

$$\Phi'(u) = \sum_{k=1} t_{2k+1} (-)^k \frac{(2k-1)!!}{2^k} u^k \quad (6.67)$$

Therefore, the differential dS (6.27) actually leads to the general solution to the Whitham hierarchy, the higher times being just constants which parameterize the solution.

To complete our consideration of particular system (6.63), we discuss what happens when one considers u as a function of *three* independent variables: t_0 , t_1 and α . In this case, one still gets the same solution depending on *two* independent variables.

Indeed, since

$$\alpha \equiv \oint_A dS = t_1 a + t_0 \quad (6.68)$$

(cf. with eq.(6.66)) and using (6.56), (6.63) turns into

$$\begin{aligned} t_1 \frac{\partial u}{\partial t_1} &= -\alpha \frac{\partial u}{\partial \alpha} + t_0 \frac{\partial u}{\partial \alpha}, \\ \frac{\partial u}{\partial t_0} &= -\frac{\partial u}{\partial \alpha} \end{aligned} \quad (6.69)$$

with the solution

$$u = \Psi \left(\frac{\alpha - t_0}{t_1} \right) \quad (6.70)$$

where $\Psi(x)$ is an arbitrary function. Note that (6.66) now can be understood just as the condition of constant α .

Let us now note that $a(u)$ for the given data is some known function of u . Then, the inverse (not arbitrary!) function $u = \Psi(a)$ can be rewritten in the form (6.70) using (6.68) with independent variables α , t_0 and t_1 . It means that one can not consider arbitrary function $\Psi(x)$ with the particular ansatz of zero higher times in (6.27), i.e. solutions to the equations (6.69) should be additionally constrained.

If now one forgets about the specific ansatz for dS with zero higher times and considers unconstrained dS (6.27), one should also consider the additional Whitham equation

$$\frac{\partial d\Omega_n}{\partial \alpha} = \frac{\partial d\omega}{\partial t_n} \quad (6.71)$$

that exactly gives rise to the equations (6.69)

$$\begin{aligned} \frac{\partial u}{\partial t_1} &= -a \frac{\partial u}{\partial \alpha}, \\ \frac{\partial u}{\partial t_0} &= -\frac{\partial u}{\partial \alpha} \end{aligned} \quad (6.72)$$

These equations are equivalent to (6.58) with u being a function of $t_0 - \alpha$. It means that we again obtain the same solution (6.61) that describes u as a function of *two* independent variables t_1 and $t_0 - \alpha$. This perfectly matches the number $g + n - 1 = 2$ of independent variables.

6.5 WDVV equations

We started this review with discussing integrability of Seiberg-Witten theory and continued with Whitham hierarchies associated both with integrable and topological properties of Seiberg-Witten theory. Now we are going to discuss briefly a feature of the prepotential that has much to do more with topological theories, while is not that straightforwardly (to our present knowledge) related to integrability. This property briefly mentioned in the Introduction is the WDVV equations (1.8), which are solved by the prepotentials of theories with the spherical bare spectral curve.

Note that we still do not understand any special reason for the WDVV equations to emerge in topological and Seiberg-Witten theories, although, first, the latter two are connected through the

Whitham hierarchies (see the previous subsection) and there are also tight connections of the Whitham hierarchies with the WDVV equations [40].

To get some insight of general structure of the WDVV equations, we start with examples not related to Seiberg-Witten theory but covering the main fields where the WDVV equations have been met so far (that is, type A and type B $2d$ topological theories).

The simplest non-trivial example of $N_c = 3$ WDVV (the equations at $N_c = 1, 2$ are trivially satisfied) comes from the type B topological theory, namely, $\mathcal{N} = 2$ SUSY Ginzburg-Landau theory with the superpotential $W'(\lambda) = \lambda^3 - q$ [40]. In this case, the prepotential reads as

$$F = \frac{1}{2}a_1a_2^2 + \frac{1}{2}a_1^2a_3 + \frac{q}{2}a_2a_3^2 \quad (6.73)$$

and the matrices F_i (the third derivatives of the prepotential) are

$$F_1 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad F_2 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & q \end{pmatrix}, \quad F_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & q \\ 0 & q & 0 \end{pmatrix}. \quad (6.74)$$

One can easily check that these matrices do really satisfy the WDVV equations (1.8).

The second example is the quantum cohomologies of $\mathbb{C}P^2$ (type A topological σ -model). In this case, the prepotential is given by the formula [42]

$$F = \frac{1}{2}a_1a_2^2 + \frac{1}{2}a_1^2a_3 + \sum_{k=1}^{\infty} \frac{N_k a_3^{3k-1}}{(3k-1)!} e^{ka_2} \quad (6.75)$$

where the coefficients N_k (describing the rational Gromov-Witten classes) counts the number of the rational curves in $\mathbb{C}P^2$ and are to be calculated. Since the matrices F have the form

$$F_1 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad F_2 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & F_{222} & F_{223} \\ 0 & F_{223} & F_{233} \end{pmatrix}, \quad F_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & F_{223} & F_{233} \\ 0 & F_{233} & F_{333} \end{pmatrix} \quad (6.76)$$

the WDVV equations are equivalent to the identity

$$F_{333} = F_{223}^2 - F_{222}F_{233} \quad (6.77)$$

which, in turn, results into the recurrent relation defining the coefficients N_k :

$$\frac{N_k}{(3k-4)!} = \sum_{a+b=k} \frac{a^2b(3b-1)b(2a-b)}{(3a-1)!(3b-1)!} N_a N_b. \quad (6.78)$$

The crucial feature of the presented examples is that, in both cases, there exists a constant matrix F_1 . Following [40], one can consider it as a flat metric on the moduli space. In fact, in its original version, the WDVV equations have been written in a slightly different form, that is, as the associativity condition of some algebra. We shall discuss this later, and now just remark that, having distinguished (constant) metric $\eta \equiv F_1$, one can naturally rewrite (1.8) as the equations

$$C_i C_j = C_j C_i \quad (6.79)$$

for the matrices $C_i \equiv \eta^{-1} F_i$, i.e. $(C_i)_k^j = \eta^{jl} F_{ilk}$. Formula (6.79) is equivalent to (1.8) with $j = 1$. Moreover, this particular relation is already sufficient [39, 168] to reproduce the whole set of the WDVV equations (1.8). Indeed, since $F_i = F_1 C_i$, we obtain

$$F_i F_j^{-1} F_k = F_1 (C_i C_k^{-1} C_j) \quad (6.80)$$

which is obviously symmetric under the permutation $i \leftrightarrow j$. Let us also note that, although the WDVV equations can be fulfilled only for some specific choices of the coordinates a_i on the moduli space, they still admit any linear transformation. This defines the flat structures on the moduli space, and we often call a_i flat coordinates.

In fact, the existence of the flat metric is not necessary for (1.8) to be true, how we explain below. Moreover, the Seiberg-Witten theories give exactly an example of such a case, where there is no distinguished constant matrix. This matrix can be found in topological theories because of existence their field theory interpretation where the unity operator is always presented.

Now note that in the context of the two-dimensional Ginzburg-landau topological theories, the WDVV equations arose as associativity condition of some polynomial algebra. We will prove below that the equations in Seiberg-Witten theory have the same origin. Now we briefly remind the main ingredients of this approach in the standard case of the Ginzburg-Landau theories.

In this case, one deals with the chiral ring formed by a set of polynomials $\{\Phi_i(\lambda)\}$ and two co-prime (i.e. without common zeroes) fixed polynomials $\mathcal{Q}(\lambda)$ and $P(\lambda)$. The polynomials Φ form the associative algebra with the structure constants C_{ij}^k given with respect to the product defined by modulo P' :

$$\Phi_i \Phi_j = C_{ij}^k \Phi_k \mathcal{Q}' + (*) P' \longrightarrow C_{ij}^k \Phi_k \mathcal{Q}' \quad (6.81)$$

the associativity condition being

$$(\Phi_i \Phi_j) \Phi_k = \Phi_i (\Phi_j \Phi_k), \quad (6.82)$$

$$\text{i.e.} \quad C_i C_j = C_j C_i, \quad (C_i)_k^j = C_{ik}^j \quad (6.83)$$

Now, in order to get from these conditions the WDVV equations, one needs to choose properly the flat moduli [40]:

$$a_i = -\frac{n}{i(n-i)} \text{res} \left(P^{i/n} d\mathcal{Q} \right), \quad n = \text{ord}(P) \quad (6.84)$$

Then, there exists the prepotential whose third derivatives are given by the residue formula

$$F_{ijk} = \text{res}_{P'=0} \frac{\Phi_i \Phi_j \Phi_k}{P'} \quad (6.85)$$

On the other hand, from the associativity condition (6.83) and the residue formula (6.85), one obtains that

$$F_{ijk} = (C_i)_j^l F_{\mathcal{Q}'lk}, \quad \text{i.e.} \quad C_i = F_i F_{\mathcal{Q}'}^{-1} \quad (6.86)$$

Substituting this formula for C_i into (6.83), one finally reaches the WDVV equations in the form

$$F_i G^{-1} F_j = F_j G^{-1} F_i, \quad (6.87)$$

$$G \equiv F_{Q'}$$

The choice $Q' = \Phi_l$ gives the standard equations (1.8). As we already mentioned, in two-dimensional topological theories, there is always the unity operator that corresponds to $Q' = 1$ and leads to the constant metric $F_{Q'}$.

Thus, from this short study of the WDVV equations in the Ginzburg-Landau theories, we can get three main ingredients necessary for these equations to hold. These are:

- associative algebra
- flat moduli (coordinates)
- residue formula

As we already know, in Seiberg-Witten theory only the first ingredient requires a non-trivial check, while the other two are automatically presented due to the proper integrable structures.

6.6 WDVV equations in Seiberg-Witten theory

Now we are ready to study the most interesting for us example of Seiberg-Witten theory and determine when the prepotential satisfies the WDVV equations. First, let us note that if the prepotential satisfies the WDVV equations, its leading perturbative part satisfies them too (since the classical quadratic piece does not contribute into the third derivatives). In each case it can be checked by straightforward calculations.

We already discussed the perturbative prepotentials in detail in the previous section, see also [39] which also contains a proof of the WDVV equations for the perturbative prepotentials. Here we just list some results of [39] adding some newer statements.

i) The WDVV equations always hold for the pure gauge theories $\mathcal{F}_{pert} = \mathcal{F}_{pert}^V$ (including the exceptional groups)⁴¹. In fact, in [39] it has been proved that, if one starts with the general prepotential of the form

$$\mathcal{F} = \frac{1}{4} \sum_{i,j} \left(\alpha_- (a_i - a_j)^2 \log(a_i - a_j) + \alpha_+ (a_i + a_j)^2 \log(a_i + a_j) \right) + \frac{\eta}{2} \sum_i a_i^2 \log a_i \quad (6.88)$$

the WDVV hold iff $\alpha_+ = \alpha_-$ or $\alpha_+ = 0$, η being arbitrary. A new result due to A.Veselov [170] shows that this form can be further generalized by adding boundary terms. For this latter case, however, we do not know the non-perturbative extension.

ii) If one considers the gauge supermultiplets interacting with the n_f matter hypermultiplets in the first fundamental representation with masses m_α

$$\mathcal{F}_{pert} = \mathcal{F}_{pert}^V + r \mathcal{F}_{pert}^F + K f_F(m) \quad (6.89)$$

⁴¹The rank of the group should be bigger than 2 for the WDVV equations not to be empty, thus for example in the pure gauge G_2 -model they are satisfied trivially. It have been checked in [39, 169] (using MAPLE package) that the WDVV equations hold for the perturbative prepotentials of the pure gauge F_4 , E_6 and E_7 models. Note that the corresponding non-perturbative Seiberg-Witten curves are not obviously hyperelliptic.

(where r and K are some undetermined coefficients), the WDVV equations do not hold unless $K = r^2/4$, the masses are regarded as moduli (i.e. the equations (1.8) contain the derivatives with respect to masses) and

$$f_F(m) = \frac{1}{4} \sum_{\alpha, \beta} \left((m_\alpha - m_\beta)^2 \log(m_\alpha - m_\beta) \right) \quad (6.90)$$

for the $SU(N_c)$ gauge group and

$$\begin{aligned} f_F(m) = & \frac{1}{4} \sum_{\alpha, \beta} \left((m_\alpha - m_\beta)^2 \log(m_\alpha - m_\beta) + (m_\alpha + m_\beta)^2 \log(m_\alpha + m_\beta) \right) + \\ & + \frac{r(r+s)}{4} \sum_{\alpha} m_\alpha^2 \log m_\alpha \end{aligned} \quad (6.91)$$

for other classical groups, $s = 2$ for the orthogonal groups and $s = -2$ for the symplectic ones.

Note that at value $r = -2$ the prepotential (6.89) can be considered as that in the pure gauge theory with the gauge group of the higher rank, $\text{rank}(\mathbf{G}) + n_f$. At the same time, at value $r = 2$, like a_i 's lying in irrep of $\mathbf{G}$, masses m_α 's can be regarded as lying in irrep of some $\tilde{\mathbf{G}}$ so that if $\mathbf{G} = A_n, C_n, D_n$, $\tilde{\mathbf{G}} = A_n, D_n, C_n$ accordingly (this is nothing but the notorious (gauge group $\longleftrightarrow$ flavor group) duality, see, e.g. [114]). These correspondences "explain" the form of the mass term in the prepotential $f(m)$.

iii) The set of the perturbative prepotentials satisfying the WDVV equations can be further extended. Namely, one can consider higher dimensional SUSY gauge theories, in particular, $5d$ theories. Then, the perturbative prepotentials in the $SU(N_c)$ theory (without adjoint matter) is given by formula (5.39) with $d = 5$. One can check that these perturbative prepotentials satisfy the WDVV equations. Note that the presence of the cubic terms in (5.39) turns to be absolutely crucial for the WDVV equations to hold. This means that the coefficient c_c in (5.43) is fixed by the equations. This is not surprising since that the WDVV equations are not necessarily satisfied when $c_c \neq N_c - N_f/2$. Since it corresponds to degeneration of an integrable system, one fixes some moduli (masses) which are the parameters to vary in the WDVV equations (see [39] for details).

One can also consider other classical groups. Then, the perturbative prepotentials acquire the form (5.8) with all $x^2 \log x$ substituted by $\frac{i}{3}x^3 + \frac{1}{2}\text{Li}_3(e^{-2iRx})$. One can easily check, along the line of [39] that these prepotentials satisfy the WDVV equations.

iv) If in the $4d$ theory the adjoint matter hypermultiplets are presented, i.e. $\mathcal{F}_{pert} = \mathcal{F}_{pert}^V + \mathcal{F}_{pert}^A + f_A(m)$, the WDVV equations never hold. At the same time, the WDVV equations are fulfilled for the theory with matter hypermultiplets in the symmetric/antisymmetric square of the fundamental representation⁴² iff the masses of these hypermultiplets are equal to zero.

v) Our last example [122, 69] has the most unclear status, at the moment. It corresponds to the $5d$ theory with adjoint matter. It is described by the perturbative prepotential (5.69). This prepotential

⁴²These hypermultiplets contribute to the prepotential

$$\begin{aligned} \mathcal{F}_{pert}^S &= -\frac{1}{4} \sum_{i \leq j} (a_i + a_j + m)^2 \log(a_i + a_j + m) \\ \mathcal{F}_{pert}^{AS} &= -\frac{1}{4} \sum_{i < j} (a_i + a_j + m)^2 \log(a_i + a_j + m) \end{aligned} \quad (6.92)$$

satisfies the WDVV equations iff $\epsilon = \pi/2$ [122]. Moreover, it gives the most general solution for a general class of perturbative prepotentials $\mathcal{F}_{pert}$ assuming the functional form

$$\mathcal{F} = \sum_{\alpha \in \Phi} f(\alpha \cdot a), \quad (6.93)$$

where the sum is over the root system Φ of a Lie algebra. The mystery about this prepotential is that, on one hand, it never satisfies the WDVV equations unless $\epsilon = \pi$ and we do not know if the corresponding complete non-perturbative prepotential satisfies the WDVV even for $\epsilon = \pi/2$. On the other hand, in the limit $R_5 \rightarrow 0$ and $\epsilon \sim MR$ for finite M , (when the mass spectrum (3.84) reduces to the two points $M_n = 0$ and $M_n = M$), the theory is the four dimensional SYM model with $\mathcal{N} = 4$ SUSY softly broken to $\mathcal{N} = 2$, i.e. includes the adjoint matter hypermultiplet. It corresponds to item **iv)** when the WDVV always do *not* hold. By all these reasons, this exceptional case deserves further investigation.

From the above consideration of the WDVV equations for the perturbative prepotentials, one can learn the following lessons:

- masses are to be regarded as moduli
- as an empiric rule, one may say that the WDVV equations are satisfied by perturbative prepotentials which depend only on the pairwise sums of the type $(a_i \pm b_j)$, where moduli a_i and b_j are either periods or masses⁴³. This is the case for the models that contain either massive matter hypermultiplets in the first fundamental representation (or its dual), or massless matter in the square product of those. Troubles arise in all other situations because of the terms with $a_i \pm b_j \pm c_k \pm \dots$ (The inverse statement is wrong – there are some exceptions when the WDVV equations hold despite the presence of such terms – e.g., for the exceptional groups.)

Note that, for the non-UV-finite theories with the perturbative prepotential satisfying the WDVV equations, one can add one more parameter to the set of moduli – the parameter Λ_{QCD} that enters all the logarithmic terms as $x^2 \log x / \Lambda_{QCD}$. Then, some properly defined WDVV equations still remain correct despite the matrices $\mathcal{F}_i^{-1}$ no longer exist (one just needs to consider instead of them the matrices of the proper minors) [171], see footnote 45.

Now let us check the WDVV equations for full Seiberg-Witten prepotentials. As we already discussed, in order to derive the WDVV equations along the line used in the context of the Ginzburg-Landau theories, we need three crucial ingredients: flat moduli, residue formula and associative algebra. However, the first two of these are always contained in the SW construction provided the underlying integrable system is known. Thus, the only point to be checked is the existence of the associative algebra. The residue formula (5.1) hints that this algebra is to be the algebra Ω^1 of the holomorphic differentials $d\omega_i$. In the forthcoming discussion we restrict ourselves to the case of pure gauge theory, the general case being treated in complete analogy.

Let us consider the algebra Ω^1 and fix three differentials $dQ, d\omega, d\lambda \in \Omega^1$. The product in this algebra is given by the expansion

$$d\omega_i d\omega_j = C_{ij}^k d\omega_k dQ + (*)d\omega + (*)d\lambda \quad (6.94)$$

⁴³This general rule can be easily interpreted in D-brane terms, since the interaction of branes is caused by strings between them. The pairwise structure $(a_i \pm b_j)$ exactly reflects this fact, a_i and b_j should be identified with the ends of string.

that should be factorized over the ideal spanned by the differentials $d\omega$ and $d\lambda$. This product belongs to the space of quadratic holomorphic differentials:

$$\Omega^1 \cdot \Omega^1 \in \Omega^2 \cong \Omega^1 \cdot (d\mathcal{Q} \oplus d\omega \oplus d\lambda) \quad (6.95)$$

Since the dimension of the space of quadratic holomorphic differentials is equal to $3g - 3$, the l.h.s. of (6.94) with arbitrary $d\omega_i$'s is the vector space of dimension $3g - 3$. At the same time, at the r.h.s. of (6.94) there are g arbitrary coefficients C_{ij}^k in the first term (since there are exactly so many holomorphic 1-differentials that span the arbitrary holomorphic 1-differential $C_{ij}^k d\omega_k$), $g - 1$ arbitrary holomorphic differentials in the second term (one differential should be subtracted to avoid the double counting) and $g - 2$ holomorphic 1-differentials in the third one. Thus, totally we get that the r.h.s. of (6.94) is spanned also by the basis of dimension $g + (g - 1) + (g - 2) = 3g - 3$.

This means that the algebra exists in the general case of the SW construction. However, generally this algebra is not associative. This is because, unlike the LG case, when it was the algebra of polynomials and, therefore, the product of the two belonged to the same space (of polynomials), product in the algebra of holomorphic 1-differentials no longer belongs to the same space but to the space of quadratic holomorphic differentials. Indeed, to check associativity, one needs to consider the triple product of Ω^1 :

$$\Omega^1 \cdot \Omega^1 \cdot \Omega^1 \in \Omega^3 = \Omega^1 \cdot (d\mathcal{Q})^2 \oplus \Omega^2 \cdot d\omega \oplus \Omega^2 \cdot d\lambda \quad (6.96)$$

Now let us repeat our calculation: the dimension of the l.h.s. of this expression is $5g - 5$ that is the dimension of the space of holomorphic 3-differentials. The dimension of the first space in expansion of the r.h.s. is g , the second one is $3g - 4$ and the third one is $2g - 4$. Since $g + (3g - 4) + (2g - 4) = 6g - 8$ is greater than $5g - 5$ (unless $g \leq 3$), formula (6.96) **does not** define the unique expansion of the triple product of Ω^1 and, therefore, the associativity spoils.

The situation can be improved if one considers the curves with additional involutions. As an example, let us consider the family of hyperelliptic curves: $y^2 = Pol_{2g+2}(\lambda)$. In this case, there is the involution, $\sigma : y \rightarrow -y$ and Ω^1 is spanned by the σ -odd holomorphic 1-differentials $\frac{x^{i-1}dx}{y}$, $i = 1, \dots, g$. Let us also note that both dQ and $d\omega$ are σ -odd, while $d\lambda$ is σ -even. This means that $d\lambda$ can be only meromorphic on the surface without punctures (which is, indeed, the case in the absence of mass hypermultiplets). Thus,

$d\lambda$ omits from formula (6.94) that acquires the form

$$\Omega_+^2 = \Omega_-^1 \cdot dQ \oplus \Omega_-^1 \cdot d\omega \quad (6.97)$$

where we expanded the space of holomorphic 2-differentials into the parts with definite σ -parity: $\Omega^2 = \Omega_+^2 \oplus \Omega_-^2$, which are manifestly given by the differentials $\frac{x^{i-1}(dx)^2}{y^2}$, $i = 1, \dots, 2g - 1$ and $\frac{x^{i-1}(dx)^2}{y}$, $i = 1, \dots, g - 2$ respectively. Now it is easy to understand that the dimensions of the l.h.s. and r.h.s. of (6.97) coincide and are equal to $2g - 1$.

Analogously, in this case, one can check the associativity. It is given by the expansion

$$\Omega_-^3 = \Omega_-^1 \cdot (dQ)^2 \oplus \Omega_+^2 \cdot d\omega \quad (6.98)$$

where both the l.h.s. and r.h.s. have the same dimension: $3g - 2 = g + (2g - 2)$. Thus, the algebra of holomorphic 1-differentials on hyperelliptic curve is really associative. This completes the proof of the WDVV equations in this case.

Now let us briefly look at cases when there exist the associative algebras basing on the spectral curves discussed in the previous section. First of all, it exists in the theories with the gauge group A_n , both in the pure gauge $4d$ and $5d$ theories and in the theories with fundamental matter, since, in accordance with the previous section, the corresponding spectral curves are hyperelliptic ones of genus n .

The theories with the gauge groups $SO(n)$ or $Sp(n)$ are also described by the hyperelliptic curves. The curves, however, are of higher genus $2n - 1$. This would naively destroy all the reasoning of this section. The arguments, however, can be restored by noting that the corresponding curves (see (3.55)) have yet **another** involution, $\rho : \lambda \rightarrow -\lambda$. This allows one to expand further the space of holomorphic differentials into the pieces with definite ρ -parity: $\Omega_-^1 = \Omega_{--}^1 \oplus \Omega_{-+}^1$ etc. so that the proper algebra is generated by the differentials from Ω_{--}^1 . One can easily check that it leads again to the associative algebra.

Consideration is even more tricky for the exceptional groups, when the corresponding curves are looking non-hyperelliptic. However, additional symmetries should allow one to get associative algebras in these cases too.

There are more cases when the associative algebra exists. First of all, these are $5d$ theories, with and without fundamental matter [39]. One can also consider the SYM theories with gauge groups being the product of several factors, with matter in the bi-fundamental representation [51]. As we discussed earlier, these theories are described by $sl(p)$ spin chains [38] and the existence of the associative algebra in this case has been checked in [172].

The situation is completely different in the adjoint matter case. In four dimensions, the theory is described by the Calogero-Moser integrable system. Since, in this case, the curve is non-hyperelliptic and has no enough symmetries, one needs to include into consideration both the differentials $d\omega$ and $d\lambda$ for algebra to exist. However, under these circumstances, the algebra is no longer to be associative how it was demonstrated above. This can be done also by direct calculation for several first values of N_c (see [39]). This also explains the lack of the perturbative WDVV equations in this case.

6.7 Covariance of the WDVV equations

After we have discussed the role of the (generalized) WDVV equations in SYM gauge theories of the Seiberg-Witten type, let us briefly describe the general structure of the equations themselves. We look at them now just as at some over-defined set of non-linear equations for a function (prepotential) of r variables⁴⁴ (times), $F(t^i)$, $i = 1, \dots, r$, which can be written in the form (6.87)

$$F_i G^{-1} F_j = F_j G^{-1} F_i, \quad (6.99)$$

$$G = \sum_{k=1}^r \eta^k F_k, \quad \forall i, j = 1, \dots, r \quad \text{and} \quad \forall \eta^k(t)$$

F_i being $r \times r$ matrices $(F_i)_{jk} = F_{,ijk} = \frac{\partial^3 F}{\partial t^i \partial t^j \partial t^k}$ and the "metric" matrix G is an arbitrary linear combination of F_k 's, with coefficients $\eta^k(t)$ that can be time-dependent.⁴⁵

The WDVV equations imply consistency of the following system of differential equations [173]:

⁴⁴We deliberately choose different notations for these variables, t instead of a in the gauge theories, in order to point out more general status of the discussion.

⁴⁵We already remarked in the previous subsection that one can add to the set of times (moduli) in the WDVV equations the parameter Λ_{QCD} [171]. In this case, the prepotential that depends on one extra variable $t^0 \equiv \Lambda$ can be

$$\left(F_{,ijk} \frac{\partial}{\partial t^l} - F_{,ijl} \frac{\partial}{\partial t^k} \right) \psi^j(t) = 0, \quad \forall i, j, k \quad (6.100)$$

Contracting with the vector $\eta^l(t)$, one can also rewrite it as

$$\frac{\partial \psi^i}{\partial t^k} = C_{jk}^i D \psi^j, \quad \forall i, j \quad (6.101)$$

where

$$C_k = G^{-1} F_k, \quad G = \eta^l F_l, \quad D = \eta^l \partial_l \quad (6.102)$$

(note that the matrices C_k and the differential D depend on choice of $\{\eta^l(t)\}$, i.e. on choice of the metric G) and (6.99) can be rewritten as

$$[C_i, C_j] = 0, \quad \forall i, j \quad (6.103)$$

As we already discussed, the set of the WDVV equations (6.99) is invariant under *linear* change of the time variables with the prepotential unchanged [39]. According to the second paper of [40] and especially to [174], there can exist also *non-linear* transformations which preserve the WDVV structure, but they generically change the prepotential. In [175], it is shown that such transformations are naturally induced by solutions of the linear system (6.100):

$$\begin{aligned} t^i &\longrightarrow \tilde{t}^i = \psi^i(t), \\ F(t) &\longrightarrow \tilde{F}(\tilde{t}), \end{aligned} \quad (6.104)$$

so that the period matrix remains intact:

$$F_{,ij} = \frac{\partial^2 F}{\partial t^i \partial t^j} = \frac{\partial^2 \tilde{F}}{\partial \tilde{t}^i \partial \tilde{t}^j} \equiv \tilde{F}_{,\tilde{i}\tilde{j}} \quad (6.105)$$

Now let us make some comments.

1. As explained in [173], the linear system (6.100) has infinitely many solutions. The "original" time-variables are among them: $\psi^i(t) = t^i$.

naturally considered as a homogeneous function of degree 2:

$$\mathcal{F}(t^0, t^1, \dots, t^r) = (t^0)^2 F(t^i/t^0),$$

see [155, 82] for the general theory. As explained in [171], the WDVV equations (6.99) for $F(t^i)$ can be also rewritten in terms of $\mathcal{F}(t^I)$:

$$\mathcal{F}_I \hat{\mathcal{G}}^{-1} \mathcal{F}_J = \mathcal{F}_J \hat{\mathcal{G}}^{-1} \mathcal{F}_I, \quad \forall I, J = 0, 1, \dots, r; \{\eta^K(t)\}$$

where this time $\mathcal{F}_I$ are $(r+1) \times (r+1)$ matrices of the third derivatives of $\mathcal{F}$ and

$$\mathcal{G} = \sum_{k=0}^r \eta^K \mathcal{F}_K, \quad \hat{\mathcal{G}}^{-1} = (\det \mathcal{G}) \mathcal{G}^{-1}$$

Note that the homogeneity of $\mathcal{F}$ implies that t^0 -derivatives are expressed through those w.r.t. t^i , e.g.

$$t^0 \mathcal{F}_{,0ij} = -\mathcal{F}_{,ijk} t^k, \quad t^0 \mathcal{F}_{,00i} = \mathcal{F}_{,ikl} t^k t^l, \quad t^0 \mathcal{F}_{,000} = -\mathcal{F}_{,klm} t^k t^l t^m \quad \text{etc.}$$

Thus, all the "metrics" $\mathcal{G}$ are degenerate, but $\hat{\mathcal{G}}^{-1}$ are non-degenerate. One can easily reformulate the entire present section in terms of $\mathcal{F}$. Then, e.g., the Baker-Akhiezer vector-function $\psi(t)$ should be just substituted by the manifestly homogeneous (of degree 0) function $\psi(t^i/t^0)$. The extra variable t^0 should not be mixed with the distinguished "zero-time" associated with the constant metric in the $2d$ topological theories which generically does not exist (when it does, see comment 3 below, we identify it with t^r).

2. Condition (6.105) guarantees that the transformation (6.104) changes the linear system (6.100) only by a (matrix) multiplicative factor, i.e. the set of solutions $\{\psi^i(t)\}$ is invariant of (6.104). Among other things this implies that successively applying (6.104) one does not produce new sets of time-variables.

3. We already discussed that, in the case of $2d$ topological models [42, 40, 174], there is a distinguished time-variable, say, t^r , such that all F_{ijk} are independent of t^r :

$$\frac{\partial}{\partial t^r} F_{ijk} = 0 \quad \forall i, j, k = 1, \dots, r \quad (6.106)$$

(equivalently, $\frac{\partial}{\partial t^i} F_{rjk} = 0 \quad \forall i, j, k$). Then, one can make the Fourier transform of (6.100) with respect to t^r and substitute it by the system

$$\frac{\partial}{\partial t^j} \widehat{\psi}_z^i = z C_{jk}^i \widehat{\psi}_z^k, \quad \forall i, j \quad (6.107)$$

where $\widehat{\psi}_z^k(t^1, \dots, t^{r-1}) = \int \psi_z^k(t^1, \dots, t^{r-1}, t^r) e^{zt^r} dt^r$. In this case, the set of transformations (6.104) can be substituted by a family, labeled by a single variable z :

$$t^i \longrightarrow \widetilde{t}_z^i = \widehat{\psi}_z^i(t) \quad (6.108)$$

In the limit $z \rightarrow 0$ and for the particular choice of the metric, $\check{G} = F_r$, one obtains the particular transformation

$$\frac{\partial \widetilde{t}^i}{\partial t^j} = \check{C}_{jk}^i h^k, \quad h^k = \text{const}, \quad (6.109)$$

discovered in [174]. (Since $\check{C}_i = \partial_j \check{C}$, one can also write $\check{C}_i = \check{C}_k^i h^k$, $\check{C}_k^i = (F_r^{-1})^{il} F_{lk}$.)

4. Parameterization like (6.109) can be used in the generic situation (6.104) as well (i.e. without distinguished t^r -variable and for the whole family (6.104)), the only change is that h^k is no longer a constant, but a solution to

$$(\partial_j - DC_j)_k^i h^k = 0 \quad (6.110)$$

($h^k = D\psi^k$ is always a solution, provided ψ^k satisfies (6.101)).

Note also that, although we have described a set of non-trivial non-linear transformations which preserve the structure of the WDVV equations (6.99), the consideration above does not *prove* that *all* such transformations are of the form (6.104), (6.105). Still, (6.104) is already unexpectedly large, because (6.99) is an *over-defined* system and it could seem to be very *restrictive*, if to have any solutions at all.

To conclude this short survey of the prepotentials and WDVV equations, let us emphasize that there are a lot of problems to be solved. We already mentioned the problem of lack of the WDVV equations for the Calogero-Moser system. The way to resolve this problem might be to construct higher associativity conditions like it has been done by E. Getzler in the elliptic case [176], that is to say, for the elliptic Gromov-Witten classes. Even if this will succeed, it is not an *explanation* why at all generalized WDVV equations should be satisfied for the Seiberg-Witten prepotentials.

7 Dualities in the many-body systems and gauge theories

In this section, we are going to discuss duality, a property of integrable systems which has much to do with the possibility of exact solving SUSY gauge theories at low energies and with the possibility of re-interpreting them in terms of strings/branes (see also the Introduction). This property also will allow us to fill in the right bottom corner of table 1, i.e. to construct a double elliptic system.

In fact, as in string theory there are several dualities (S-, T- and U-dualities), as in integrable systems there are several kinds of duality [177]. We briefly survey all of them here, and then concentrate on the most interesting, important and still remaining mysterious one which can be described via canonical transformations [178, 92]. Note that all the dualities at the moment are attributed only with the Hitchin type systems, i.e. the Seiberg-Witten theories with adjoint matter, while spin chains are still not embedded into this treatment.

Let us point out that the dualities under discussion interchanges different theories, while there is also a duality that interchanges descriptions of *the same* integrable system. This is the duality between $N_c \times N_c$ and 2×2 Lax representations, we discuss it in the next section.

7.1 D-branes, gauge theory and separation of variables

Thus, similarly to branes/string theories, the Hitchin type systems are involved into three different dualities. Let us start with considering a counterpart of the T-duality which can be identified with the separation of variables in dynamical systems [177]. The separation of variables (SoV) in integrable system allows one to reduce the system with N degrees of freedom to N identical systems with one degree of freedom each.

Recently, E.Sklyanin formulated a recipe for the SoV in a large class of quantum integrable models [179]. The method is to study the pole dynamics of the Baker-Akhiezer function $\Psi(z, \lambda)$:

$$L(\xi)\Psi(\xi, \lambda) = \lambda(\xi)\Psi(\xi, \lambda) \quad (7.111)$$

with some clever choice of normalization which is the main, difficult and artistic part of the method. The poles ξ_i of $\Psi(\xi, \lambda)$ together with the eigenvalues $\lambda_i = \lambda(\xi_i)$ are then identified with the (canonically conjugated) separated variables.

In [177] there was constructed a geometrical picture behind the recipe (see also [180, 181]). Their construction of the separation of variables in integrable systems on moduli spaces of holomorphic bundles with some additional structures that can be described as a symplectomorphism between the moduli spaces of bundles (more precisely, torsion free sheaves) having different Chern classes. In [177], there were discussed three equivalent realizations of the Hitchin integrable systems.

The first one uses non-Abelian gauge fields on the base curve Σ embedded into a symplectic surface S . The phase space of the integrable system is the moduli space of the stable pairs $(\mathcal{E}, \phi)$, where $\mathcal{E}$ is the rank r vector bundle over Σ of degree l , while ϕ is the holomorphic section of $\omega_\Sigma^1 \otimes \text{End}(\mathcal{E})$. In the second realization, one deals with the moduli space of pairs $(\mathcal{E}, \phi)$, where C is the curve (divisor) in S which realizes the homology class $r[\Sigma]$ and $\mathcal{L}$ is the line bundle on C . The third realization is the Hilbert scheme of points on S with the number of points determined by the rank of the gauge group⁴⁶.

⁴⁶ The role of the Hilbert schemes in the context of Hitchin systems was first pointed out in [182] (see also the original context of [124]) for the surfaces without marked points and then generalized to the Calogero systems in [183, 177].

The equivalence of these three realizations can be interpreted in string theory terms. To this end, let us specify the moduli space $\mathcal{M}_{\vec{v}}$ of stable torsion free coherent sheaves $\mathcal{E}$ on S . Let $\widehat{A}_S = 1 - [\text{pt}] \in H^*(S, \mathbb{Z})$ be the A -roof genus of S . The vector $\vec{v} = Ch(\mathcal{E})\sqrt{\widehat{A}_S} = (r; \vec{w}; d - r) \in H^*(S, \mathbb{Z})$, $\vec{w} \in \Gamma^{3,19}$ corresponds to the sheaves with the Chern numbers:

$$\begin{aligned} ch_0(\mathcal{E}) &= r \in H^0(S; \mathbb{Z}) \\ ch_1(\mathcal{E}) &= \vec{w} \in H^2(S; \mathbb{Z}) \\ ch_2(\mathcal{E}) &= d \in H^4(S; \mathbb{Z}) \end{aligned} \tag{7.112}$$

Now let us describe geometrically string theory objects. In type IIA string theory compactified onto S , there are Dp-branes with even p which wrap different supersymmetric cycles in S labelled by $\vec{v} \in H^*(S, \mathbb{Z})$. The BPS states correspond to the cohomology classes of the moduli spaces, $\mathcal{M}_{\vec{v}}$ of brane configurations. These moduli spaces can be identified with the moduli spaces of appropriate sheaves, $\mathcal{M}_{\vec{v}}$.

The string theory compactified on S , has the following vacuum moduli space

$$\mathcal{M}_A = O(\Gamma^{4,20}) \backslash O(4, 20; R) / O(4; R) \times O(20; R)$$

where the arithmetic group $O(\Gamma^{4,20})$ is the group of discrete automorphisms that maps between the states with the different $\vec{v}$. The $\vec{v}^2$ is the only invariant of the group action.

Now, the equivalence of the first and the second realizations of Hitchin systems corresponds to the physical statement that the bound states of N $D2$ -branes wrapped around Σ are represented by a single $D2$ -brane which wraps a holomorphic curve C which is an N -sheet covering of the base curve Σ . The equivalence of the second and the third descriptions is naturally attributed to T-duality.

Let us mention that the separation of variables above provides some insights on the Langlands duality which involves the spectrum of the Hitchin Hamiltonians. An attempt to reformulate Langlands duality as a quantum separation of variables has been presented in [185], while in [178] the authors proposed to identify the proper classical version of the Langlands correspondence with the map of the phase space of the Hitchin system to the Hilbert scheme of points on four-dimensional manifold.

7.2 S-duality

Now let us see that S-duality well-established in field theory also has a clear counterpart in the holomorphic dynamical system. In field theory, S-duality just maps one theory onto another so that the weak coupling regime of one theory maps onto the strong coupling regime of the other one and vice versa. Therefore, a hint of what is S-duality in integrable systems can come from Seiberg-Witten theory where this duality just maps the weak coupling, perturbative limit of the gauge theory (due to the asymptotic freedom at large energies, where the gauge bosons are almost massless, with their masses proportional to a_i) onto the strong coupling limit (where the BPS monopole is almost massless, with the mass proportional to a_i^D). This means that S-duality interchanges theories with small a and small a_D . Note now that, in the integrable set-up, formula (1.3) it just corresponds to interchanging A - and B -cycles. Therefore, the duality transform is just the $SP(2g, \mathbb{Z})$ group acting on the $2g$ cycles and the whole set of variables $\{a_i, a_i^D\}$ is just local coordinates on the $SP(2g, \mathbb{Z})$ bundle.

It is an easy exercise on the properties of the period matrix of Abelian varieties to check that the two form

$$da^i \wedge da_i^D \quad (7.113)$$

vanishes. Therefore one can always locally find a function $\mathcal{F}$, *prepotential* such that

$$a_i^D = \frac{\partial \mathcal{F}}{\partial a^i} \quad (7.114)$$

To illustrate this scheme with some formulas, let us look at the two-body system, relevant for the $SU(2)$ $\mathcal{N} = 2$ supersymmetric gauge theory. It is the periodic Toda chain with the Hamiltonian (see s.3.2)

$$H = \frac{p^2}{2} + \cos(q) \quad (7.115)$$

The action variable is given by one of the periods of the differential pdq (3.19) on the curve (3.18). Then, the two periods a and a^D obey the Picard-Fuchs equation

$$\left(\frac{d^2}{du^2} + \frac{1}{4(u^2 - 1)} \right) \begin{pmatrix} a \\ a^D \end{pmatrix} = 0$$

which can be used to obtain the asymptotic expansions of the action variable and the prepotential in the vicinity of $u = \infty$ or $u = \pm 1$. The S-duality manifests itself in the fact that near $u = \infty$ (which corresponds to high energies in the two-body problem and also to the perturbative regime of the $SU(2)$ gauge theory) the appropriate action variable is a , while near $u = 1$ (which corresponds to the dynamics of the two-body system near the top of the potential and to the strongly coupled $SU(2)$ gauge theory) it is a^D . The monodromy invariant combination of these periods [162]

$$aa^D - 2\mathcal{F} = u \quad (7.116)$$

(which originates from the periods of Calabi-Yau manifolds, on one hand, and from the properties of anomaly in field theory, on the other) can be chosen as a global coordinate on the space of integrals of motion. At $u \rightarrow \infty$ the prepotential has an expansion of the form

$$\mathcal{F} \sim \frac{1}{2}u \log u + \dots \sim a^2 \log a + \sum_n f_n a^{2-4n}$$

Let us emphasize that S-duality maps the dynamical system to itself, just at a different energy.

7.3 “Mirror” symmetry in dynamical systems

Now we come to the most interesting type of duality, a duality between pairs of dynamical systems (which is self-duality for some systems) [62, 124, 178, 92]. This duality plays a role similar to the mirror symmetry.

To start with, let us see how this symmetry is formulated in field theory. The initial motivation comes from the 3d theory example [186], where the mirror symmetry interchanges the Coulomb and Higgs branches of the moduli space. The specifics of three dimensions is that both the Coulomb and Higgs branches are hyperkahler manifolds and the mirror symmetry can be formulated as a kind of hyperkahler rotation.

Taking into account the relation between dynamical systems and low-energy effective actions, one can formulate a counterpart of this mirror symmetry for gauge theories in different dimensions (see further details in [178]).

Let us consider a system with the phase space being a hyperkahler four dimensional manifold. Let this manifold at most involve two tori, i.e. be the elliptically fibered K3 manifold. One torus is where momenta live, and the second one is where coordinates do. The duality at hands actually interchanges momentum and coordinate tori. However due to the nontrivial fibration this interchange is far from being naive. Note that typically one deals with degenerations of these tori and only in the double elliptic case two elliptic moduli survive.

One can also consider a theory which is self-dual. This integrable system is called Dell (double elliptic) system⁴⁷ and describes the Seiberg-Witten theory for the six-dimensional theory with adjoint matter.

The other cases correspond to some degenerations of Dell system. Say, the degeneration of the momentum torus to C/Z_2 leads to the five-dimensional theory, while the degeneration to R^2 leads to the four-dimensional theory. Since the modulus of the coordinate torus has meaning of the complexified bare coupling constant in field theory, interpretation of the degenerations of the coordinate torus is different. In particular, the degeneration to the cylinder corresponds to switching off the instanton effects, i.e. to the perturbative regime.

We shall see below that the “mirror” symmetry maps theories in different dimensions to each other. Instanton effects in one theory “get mapped” into an additional compact dimension in the dual counterpart. We discuss here the pq -duality mostly for classical integrable systems only. Since the wave functions in the Hitchin like systems can be identified with some solutions to the KZ or qKZ equations, the quantum duality would imply some relations between solutions to the rational, trigonometric and elliptic KZ equations. Recently the proper symmetries for the KZ equations were discussed in [75].

Now let us formulate the same mirror symmetry in terms of dynamical system. The idea of duality here expresses a relationship between two completely integrable systems S_1, S_2 on a fixed symplectic manifold with given symplectic structure (M, ω) and goes back to [62]. We say the Hamiltonian systems are *dual* when the conserved quantities of S_1 and S_2 together form a coordinate system for M . Consider for example free particles, $H_k^{(1)} = \sum_i p_i^k / k$. For this system the free particles momenta are identical to the conserved quantities or action variables. Now consider the Hamiltonian $H_k^{(2)} = \sum_i q_i^k / k$ with conserved quantities q_i . Together $\{p_i, q_i\}$ form a coordinate system for phase space, and so the two sets of Hamiltonians are dual. Duality then in this simplest example is a transformation which interchanges momenta and coordinates. For more complicated interacting integrable systems finding dual Hamiltonians is a nontrivial exercise. Note that this whole construction manifestly depends on the particular choice of conserved quantities. A clever choice may result in the dual system arising by simply interchanging the momentum and coordinate dependence, as in the free system.

Some years ago Ruijsenaars [62] observed such dualities between various members of the Calogero-Moser and Ruijsenaars families: the rational Calogero and trigonometric Ruijsenaars models were dual to themselves while trigonometric Calogero model was dual with the rational Ruijsenaars system. These dualities were shown by starting with a Lax pair $L = L(p, q)$ and an auxiliary diagonal matrix $A = A(q)$. When L was diagonalized the matrix A became the Lax matrix for the dual Hamiltonian,

⁴⁷In fact, this is an example of the Dell system with 2 degrees of freedom. One may equally consider the Dell system with arbitrary many degrees of freedom, see ss.6.3,6.4.

while L was a function of the coordinates of the dual system. Dual systems for a model possessing a Lax representation are then related to the eigenvalue motion of the Lax matrix. We discuss this approach in more details below.

Another approach to finding a dual system is to make a canonical transformation which substitutes the original set of Poisson-commuting coordinates q_i , $\{q_i, q_j\} = 0$, by another obvious set of the Poisson-commuting variables: the Hamiltonians $h_i(\vec{p}, \vec{q})$ or, better, the action variables $a_i(\vec{h}) = a_i(\vec{p}, \vec{q})$. It will be clear below that in practice really interesting transformations are a little more sophisticated: h_i are identified with certain functions of the new coordinates (these functions determine the Ruijsenaars matrix $A(q)$), which – in the most interesting cases – are just the same Hamiltonians with the interactions switched-off. Such free Hamiltonians are functions of momenta alone, and the dual coordinates substitute these momenta, just as one had for the system of free particles.

The most interesting question for our purposes is: *what are the duals of the elliptic Calogero and Ruijsenaars systems?* Since the elliptic Calogero (Ruijsenaars) is rational (trigonometric) in momenta and elliptic in the coordinates, the dual will be elliptic in momenta and rational (trigonometric) in coordinates. Having found such a model the final elliptization of the coordinate dependence is straightforward, providing us with the wanted double-elliptic systems.

7.4 Duality as the canonical transformation: two-body system

The way of constructing dual systems via a canonical transformation is technically quite tedious for many-particle systems, but perfectly fits the case of $SU(2)$ which, in the center-of-mass frame, has only one coordinate and one momentum. In this case the duality transformation can be described explicitly since the equations of motion can be integrated in a straightforward way. Technically, given two Hamiltonian systems, one with the momentum p , coordinate q and Hamiltonian $h(p, q)$ and another with the momentum P , coordinate Q and Hamiltonian $H(P, Q)$ we may describe duality by the relation

$$\begin{aligned} h(p, q) &= f(Q), \\ H(P, Q) &= F(q). \end{aligned} \tag{7.117}$$

Here the functions $f(Q)$ and $F(q)$ are such that

$$dP \wedge dQ = -dp \wedge dq, \tag{7.118}$$

which expresses the fact we have an (anti-) canonical transformation. This relation entails that

$$F'(q) \frac{\partial h(p, q)}{\partial p} = f'(Q) \frac{\partial H(P, Q)}{\partial P}. \tag{7.119}$$

At this stage the functions $f(Q)$ and $F(q)$ are arbitrary. However, when the Hamiltonians depend on a coupling constant ν^2 and are such that their “free” part can be separated and depends only on the momenta,⁴⁸ the free Hamiltonians provide a natural choice for these functions: $F(q) = h_0(q)$ and $f(Q) = H_0(Q)$ where

⁴⁸ Note that this kind of duality relates the weak coupling regime for $h(p, q)$ to the weak coupling regime for $H(P, Q)$.

$$\begin{aligned} h(p, q)|_{\nu^2=0} &= h_0(p), \\ H(P, Q)|_{\nu^2=0} &= H_0(P). \end{aligned} \tag{7.121}$$

With such choice the duality equations become

$$\begin{aligned} h_0(Q) &= h(p, q), \\ H_0(q) &= H(P, Q), \\ \frac{\partial h(p, q)}{\partial p} H'_0(q) &= h'_0(Q) \frac{\partial H(P, Q)}{\partial P}. \end{aligned} \tag{7.122}$$

Free rational, trigonometric and elliptic Hamiltonians are $h_0(p) = \frac{p^2}{2}$, $h_0(p) = \cosh p$ and $h_0(p) = \text{cn}(p|k)$ respectively.

Note that from the main duality relation,

$$H_0(q) = H(P, Q) \tag{7.123}$$

it follows that

$$\left. \frac{\partial q}{\partial P} \right|_Q = \frac{1}{H'_0(q)} \frac{\partial H(P, Q)}{\partial P} \tag{7.124}$$

which together with (7.122) implies:

$$\left. \frac{\partial q}{\partial P} \right|_Q = \frac{1}{h'_0(Q)} \frac{\partial h(p, q)}{\partial p} \tag{7.125}$$

When compared with the Hamiltonian equation for the original system,

$$\frac{\partial q}{\partial t} = \frac{\partial h(p, q)}{\partial p}, \tag{7.126}$$

we see that $P = h'_0(Q)t$ is proportional to the ordinary time-variable t , while $h_0(Q) = h(p, q) = E$ expresses Q as a function of the energy E . This is a usual feature of classical integrable systems, exploited in Seiberg-Witten theory [37]: in the $SU(2)$ case the spectral curve $q(t)$ can be described by

$$h\left(p\left(\frac{\partial q}{\partial t}, q\right), q\right) = E. \tag{7.127}$$

where p is expressed through $\partial q/\partial t$ and q from the Hamiltonian equation $\partial q/\partial t = \partial H/\partial p$. In other words, the spectral curve is essentially the solution of the equation of motion of integrable system, where the time t plays the role of the spectral parameter and the energy E that of the modulus.

Let us consider several simple examples. The simplest one is the rational Calogero. In this case, the duality transformation connects two identical Hamiltonians, (7.120). Therefore, this system is

For example, in the rational Calogero case

$$\begin{aligned} h(p, q) &= \frac{p^2}{2} + \frac{\nu^2}{q^2} = \frac{Q^2}{2} \\ H(P, Q) &= \frac{P^2}{2} + \frac{\nu^2}{Q^2} = \frac{q^2}{2} \end{aligned} \tag{7.120}$$

We recall that, on the field theory side, the coupling constant ν is related to the mass of adjoint hypermultiplet and thus remains unchanged under duality transformations.

self-dual. Somewhat less trivial example is the trigonometric Calogero-Sutherland model. It leads to the following dual pair

$$\begin{aligned} h(p, q) &= \frac{p^2}{2} + \frac{\nu^2}{\sin^2 q} = \frac{Q^2}{2} \\ H(P, Q) &= \sqrt{1 - \frac{2\nu^2}{Q^2}} \cos P = \cos q \end{aligned} \quad (7.128)$$

One may easily recognize in the second Hamiltonian the rational Ruijsenaars Hamiltonian. Thus, the Hamiltonian rational in coordinates and momenta maps onto itself under the duality, while that with rational momentum dependence and trigonometric coordinate dependence maps onto the Hamiltonian with inverse, rational coordinate and trigonometric momentum dependencies. Therefore, we can really see that the pq -duality exchanges types of the coordinate and momentum dependence. This implies that the Hamiltonian trigonometrically dependent both on coordinates and momenta is to be self-dual. Indeed, one can check that the trigonometric Ruijsenaars-Schneider system is self-dual:

$$\begin{aligned} h(p, q) &= \sqrt{1 - \frac{2\nu^2}{q^2}} \cos p = \cos Q \\ H(P, Q) &= \sqrt{1 - \frac{2\nu^2}{Q^2}} \cos P = \cos q \end{aligned} \quad (7.129)$$

Now we make the next step and introduce into the game elliptic dependencies.

7.5 Two-body systems dual to Calogero and Ruijsenaars

We begin with the elliptic Calogero Hamiltonian⁴⁹

$$h(p, q) = \frac{p^2}{2} + \frac{\nu^2}{\operatorname{sn}^2(q|k)}, \quad (7.134)$$

⁴⁹We use below the Jacobi functions. Recall the standard relations:

$$\begin{aligned} \operatorname{sn}(q) &= \sqrt{\frac{e_{13}}{\wp(\tilde{q}) - e_3}} = \frac{1}{\sqrt{k}} \frac{\vartheta_1(\widehat{q})}{\vartheta_4(\widehat{q})} = \frac{1}{\sqrt{k}} \frac{\theta_{11}(\widehat{q}|\tau)}{\theta_{01}(\widehat{q}|\tau)}, \\ \operatorname{cn}(q) &= \sqrt{\frac{\wp(\tilde{q}) - e_1}{\wp(\tilde{q}) - e_3}} = \sqrt{\frac{k'}{k}} \frac{\vartheta_2(\widehat{q})}{\vartheta_4(\widehat{q})} = \sqrt{\frac{k'}{k}} \frac{\theta_{10}(\widehat{q}|\tau)}{\theta_{01}(\widehat{q}|\tau)}, \\ \operatorname{dn}(q) &= \sqrt{\frac{\wp(\tilde{q}) - e_2}{\wp(\tilde{q}) - e_3}} = \sqrt{k'} \frac{\vartheta_3(\widehat{q})}{\vartheta_4(\widehat{q})} = \sqrt{k'} \frac{\theta_{00}(\widehat{q}|\tau)}{\theta_{01}(\widehat{q}|\tau)}. \end{aligned} \quad (7.130)$$

Here the Jacobi moduli k^2 and $k'^2 = 1 - k^2$ are the cross-ratios of the ramification points of the (hyper-) elliptic representation of the torus,

$$y^2 = \prod_{a=1}^3 (x - e_a(\tau)), \quad \sum_{a=1}^3 e_a = 0, \quad x = \wp(\tilde{q}), \quad y = \frac{1}{2} \wp'(\tilde{q}). \quad (7.131)$$

Then

$$k^2 = \frac{e_{23}}{e_{13}} = \frac{\vartheta_2^4(0)}{\vartheta_3^4(0)} = \frac{\theta_{10}^4(0|\tau)}{\theta_{00}^4(0|\tau)}, \quad k'^2 = 1 - k^2 = \frac{e_{12}}{e_{13}} = \frac{\vartheta_4^4(0)}{\vartheta_3^4(0)} = \frac{\theta_{01}^4(0|\tau)}{\theta_{00}^4(0|\tau)} \quad (7.132)$$

and

$$e_{ij} = e_i - e_j, \quad q = 2K(k)\widehat{q}, \quad \tilde{q} = 2\omega\widehat{q}, \quad e_{13} = \frac{K^2(k)}{\omega^2} \quad (7.133)$$

where $K(k)$ is the complete elliptic integral of the first kind.

and seek a dual Hamiltonian elliptic in the momentum. Thus $h_0(p) = \frac{p^2}{2}$ and we seek $H(P, Q) = H_0(q)$ such that $H_0(q) = \text{cn}(q|k)$. Eqs.(7.123) become

$$\frac{Q^2}{2} = \frac{p^2}{2} + \frac{\nu^2}{\text{sn}^2(q|k)}, \quad \text{cn}(q|k) = H(P, Q), \quad p \cdot \text{cn}'(q|k) = Q \frac{\partial H(P, Q)}{\partial P}. \quad (7.135)$$

Upon substituting

$$\text{cn}'(q|k) = -\text{sn}(q|k)\text{dn}(q|k) = -\sqrt{(1-H^2)(k'^2 + k^2H^2)}, \quad (7.136)$$

(this is because $\text{sn}^2 q = 1 - \text{cn}^2 q$, $\text{dn}^2 q = k'^2 + k^2 \text{cn}^2 q$, $k'^2 + k^2 = 1$ and $\text{cn} q = H$) we get for (7.135):

$$\left(\frac{\partial H}{\partial P} \right)^2 = \frac{p^2}{Q^2} (1 - H^2)(k'^2 + k^2 H^2). \quad (7.137)$$

Now from the first eqn.(7.135) p^2 can be expressed through Q and $\text{sn}^2(q|k) = 1 - \text{cn}^2(q|k) = 1 - H^2$ as

$$\frac{p^2}{Q^2} = 1 - \frac{2\nu^2}{Q^2(1 - H^2)}, \quad (7.138)$$

so that

$$\left(\frac{\partial H}{\partial P} \right)^2 = \left(1 - \frac{2\nu^2}{Q^2} - H^2 \right) (k'^2 + k^2 H^2). \quad (7.139)$$

Therefore H is an elliptic function of P , namely [178, 92]

$$H(P, Q) = \text{cn}(q|k) = \alpha(Q) \cdot \text{cn} \left(P \sqrt{k'^2 + k^2 \alpha^2(Q)} \left| \frac{k\alpha(Q)}{\sqrt{k'^2 + k^2 \alpha^2(Q)}} \right. \right) \quad (7.140)$$

with

$$\alpha^2(Q) = \alpha_{rat}^2(Q) = 1 - \frac{2\nu^2}{Q^2}. \quad (7.141)$$

In the limit $\nu^2 = 0$, when the interaction is switched off, $\alpha(q) = 1$ and $H(P, Q)$ reduces to $H_0(P) = \text{cn}(P|k)$, as assumed in (7.135).

We have therefore obtained a dual formulation of the elliptic Calogero model (in the simplest $SU(2)$ case). At first glance our dual Hamiltonian looks somewhat unusual. In particular, the relevant elliptic curve is “dressed”: it is described by an effective modulus

$$k_{eff} = \frac{k\alpha(Q)}{\sqrt{k'^2 + k^2 \alpha^2(Q)}} = \frac{k\alpha(Q)}{\sqrt{1 - k^2(1 - \alpha^2(Q))}}, \quad (7.142)$$

which differs from the “bare” one k in a Q -dependent way. In fact k_{eff} is nothing but the modulus of the “reduced” Calogero spectral curve [90, 91].

Now one can rewrite (7.140) in many different forms, in particular, in terms of θ -functions, in hyperelliptic parameterization etc. This latter is rather illuminating and allows one to picture mapping between the bare and dressed tori. Therefore, we refer the reader for the details to the paper [92].

Let us now construct the action-angle variables for the 2-particle Calogero system and see how they related with the dual variables P and Q .

Since this system possesses one degree of freedom, we proceed similarly to the 2-particle Toda case of s.3.2 and construct the action variable, a as the integral of differential $dS = pdq$ on the level submanifold of the phase space. Then, using formula (7.135), one obtains

$$a = \oint \sqrt{Q^2 - \frac{2\nu^2}{\text{sn}^2(q)}} dq \quad (7.143)$$

which expresses a through Q quite transcendently. Using this relation, one obtains

$$da = dQ \oint \frac{Q}{\sqrt{Q^2 - \frac{2\nu^2}{\text{sn}^2(q)}}} dq \quad (7.144)$$

We already met this integral in s.5.6, formula (5.51). It can be manifestly calculated giving

$$da = \frac{1}{K(k)} \frac{K(k_{eff}(Q))}{\sqrt{k'^2 + k^2 \alpha^2(Q)}} dQ \quad (7.145)$$

where we used the formulas of footnote 49 for the both bare and dressed tori as well as the definitions (7.141) and (7.142). Note that $da = dQ$ at large $Q = \sqrt{2E}$.

Since the coordinate on the Jacobian differs from the argument of the Jacobi function by a factor $2K(k)$ (see footnote 49), one expects for the angle variable p^{Jac} (looking at the Hamiltonian (7.140))

$$p^{Jac} = \frac{P \sqrt{k'^2 + k^2 \alpha^2(Q)}}{2K(k_{eff}(Q))} \quad (7.146)$$

This is, indeed, the case, up to inessential constant normalization factor $2K(k)$ (emerging here since the original Calogero symplectic structure is defined here as $dp \wedge dq$ with q being the argument of the Jacobi function, but not θ -function), since it follows from (7.145) and (7.146) that

$$dP \wedge dQ = \frac{1}{2K(k)} dp^{Jac} \wedge da \quad (7.147)$$

To conclude our discussion of the Calogero model, we make some comments on its solution. According to the argument of the previous subsection, our Hamiltonian (7.140) should be simply related to the solution $q(t)$ of the equations of motion of the Calogero Hamiltonian, which in the case of $SU(2)$ are immediately integrated to give

$$H^{Cal} \left(\frac{\partial q}{\partial P}, q \right) = E \quad (7.148)$$

More explicitly, the equation

$$\frac{dq}{dt} = \sqrt{2E - \frac{2\nu^2}{\text{sn}^2(q|k)}}, \quad (7.149)$$

has a solution [187]:

$$\text{cn}(q|k) = \sqrt{1 - \frac{\nu^2}{E}} \cdot \text{cn} \left(t \sqrt{2E - 2\nu^2 k^2} \middle| k \sqrt{\frac{E - \nu^2}{E - \nu^2 k^2}} \right). \quad (7.150)$$

This may be derived straightforwardly by differentiating both sides and applying (7.136). Note that the Calogero equation (7.149) and the *family* of Calogero spectral curves are essentially independent

of the value of coupling constant ν^2 : it can be absorbed into rescaling of moduli (like E) and the time-variables (like t).

In order to see that (7.150) is identical to (7.140) one needs to put $E = Q^2/2$ and make the rescaling $P = h'_0(Q)t = Qt$. With these substitutions we find that

$$\sqrt{1 - \frac{\nu^2}{E}} = \sqrt{1 - \frac{\nu^2}{Q^2}} = \alpha_{rat}(Q), \quad (7.151)$$

and

$$\begin{aligned} t\sqrt{2E - 2\nu^2 k^2} &= P\sqrt{1 - \frac{2\nu^2 k^2}{Q^2}} = P\sqrt{k'^2 + k^2\alpha_{rat}^2(Q)}, \\ k\sqrt{\frac{E - \nu^2}{E - \nu^2 k^2}} &= k\sqrt{\frac{1 - 2\nu^2/Q^2}{1 - 2\nu^2 k^2/Q^2}} = \frac{k\alpha_{rat}(Q)}{\sqrt{k'^2 + k^2\alpha_{rat}^2(Q)}}. \end{aligned} \quad (7.152)$$

We then see that (7.150) is identical to (7.140).

We remark that the relevant symplectic structure here is⁵⁰

$$dE \wedge dt = QdQ \wedge dt = -dP \wedge dQ. \quad (7.153)$$

All of the above formulae are straightforwardly generalized from the Calogero (rational-elliptic) system to the Ruijsenaars (trigonometric-elliptic) system. The only difference ensuing is that the q -dependence of the dual (elliptic-trigonometric) Hamiltonian is now trigonometric rather than rational [92] (see also [178] for a more involved formula in the action-angle variables; this, however, makes formula not convenient for the double elliptic generalization):

$$\alpha^2(q) = \alpha_{trig}^2(q) = 1 - \frac{2\nu^2}{\sinh^2 q} \quad (7.154)$$

Rather than giving further details we will proceed directly to a consideration of the double-elliptic model.

7.6 Double-elliptic two-body system

In order to get a double-elliptic system one needs to exchange the rational Q -dependence in (7.135) for an elliptic one, and so we substitute $\alpha_{rat}^2(Q)$ by the obvious elliptic analogue $\alpha_{ell}^2(Q) = 1 - \frac{2\nu^2}{\text{sn}^2(Q|\tilde{k})}$. Moreover, now the elliptic curves for q and Q need not in general be the same, i.e. $\tilde{k} \neq k$.

Instead of (7.135) the duality equations now become

⁵⁰ In what follows we prove that not only the elliptic-rational (the dual of the elliptic Calogero model) but also the elliptic-trigonometric (the dual of the elliptic Ruijsenaars model) and the elliptic-elliptic (our new double-elliptic) Hamiltonians have the same form (7.150), but the latter with the identifications $E = \frac{1}{2}\sinh^2 Q$ and $E = \frac{1}{2}\text{sn}^2(Q|\tilde{k})$. Thus they are also related to Calogero equation (7.149). However, the relevant symplectic structures – which are always given by $dP \wedge dQ = h'_0(Q)dt \wedge dQ = dh_0(Q) \wedge dt$ – are no longer equivalent to $dE \wedge dt$ (since $E \neq h_0(Q)$, i.e. E is no longer associated with the proper Hamiltonian).

$$\begin{aligned}
\text{cn}(q|k) &= H(P, Q|k, \tilde{k}), \\
\text{cn}(Q|\tilde{k}) &= H(p, q|\tilde{k}, k), \\
\text{cn}'(Q|\tilde{k}) \frac{\partial H(P, Q|k, \tilde{k})}{\partial P} &= \text{cn}'(q|k) \frac{\partial H(p, q|\tilde{k}, k)}{\partial p},
\end{aligned} \tag{7.155}$$

and the natural ansatz for the Hamiltonian (suggested by (7.140)) is

$$\begin{aligned}
H(p, q|\tilde{k}, k) &= \alpha(q|\tilde{k}, k) \cdot \text{cn} \left(p \beta(q|\tilde{k}, k) \mid \gamma(q|\tilde{k}, k) \right) = \alpha \text{cn}(p\beta|\gamma), \\
H(P, Q|k, \tilde{k}) &= \alpha(Q|k, \tilde{k}) \cdot \text{cn} \left(P \beta(Q|k, \tilde{k}) \mid \gamma(Q|k, \tilde{k}) \right) = \tilde{\alpha} \text{cn}(P\tilde{\beta}|\tilde{\gamma}).
\end{aligned} \tag{7.156}$$

Substituting these ansatz into (7.155) and making use of (7.136), one can arrive, after some calculations [92], at⁵¹

$$\alpha^2(q|\tilde{k}, k) = \alpha^2(q|k) = 1 - \frac{2\nu^2}{\text{sn}^2(q|k)}, \quad \beta^2(q|\tilde{k}, k) = \tilde{k}'^2 + \tilde{k}^2 \alpha^2(q|k), \quad \gamma^2(q|\tilde{k}, k) = \frac{\tilde{k}^2 \alpha^2(q|k)}{\tilde{k}'^2 + \tilde{k}^2 \alpha^2(q|k)}, \tag{7.157}$$

with some constant ν and finally the double-elliptic duality becomes [92]

$$H(P, Q|k, \tilde{k}) = \text{cn}(q|k) = \alpha(Q|\tilde{k}) \text{cn} \left(P \sqrt{k'^2 + k^2 \alpha^2(Q|\tilde{k})} \mid \frac{k \alpha(Q|\tilde{k})}{\sqrt{k'^2 + k^2 \alpha^2(Q|\tilde{k})}} \right), \tag{7.158}$$

$$H(p, q|\tilde{k}, k) = \text{cn}(Q|\tilde{k}) = \alpha(q|k) \text{cn} \left(p \sqrt{\tilde{k}'^2 + \tilde{k}^2 \alpha^2(q|k)} \mid \frac{\tilde{k} \alpha(q|k)}{\sqrt{\tilde{k}'^2 + \tilde{k}^2 \alpha^2(q|k)}} \right). \tag{7.159}$$

We shall now consider various limiting cases arising from these and show that the double-elliptic Hamiltonian (7.159) contains the entire Ruijsenaars-Calogero and Toda family as its limiting cases, as desired. (Of course we have restricted ourselves to the $SU(2)$ members of this family.)

In order to convert the elliptic dependence of the momentum p into the trigonometric one, the corresponding “bare” modulus $\tilde{k}$ should vanish: $\tilde{k} \rightarrow 0$, $\tilde{k}'^2 = 1 - \tilde{k}^2 \rightarrow 1$ (while k can be kept finite). Then, since $\text{cn}(x|\tilde{k} = 0) = \cosh x$,

$$H^{dell}(p, q) \longrightarrow \alpha(q) \cosh p = H^R(p, q) \tag{7.160}$$

with the same

$$\alpha^2(q|k) = 1 - \frac{2\nu^2}{\text{sn}^2(q|k)}. \tag{7.161}$$

Thus we obtain the $SU(2)$ elliptic Ruijsenaars Hamiltonian.⁵² The trigonometric and rational Ruijsenaars as well as all of the Calogero and Toda systems are obtained through further limiting procedures

⁵¹For ease of expression hereafter we suppress the dependence of α, β, γ on k and $\tilde{k}$ in what follows using $\alpha(q)$ for $\alpha(q|\tilde{k}, k)$ and $\tilde{\alpha}(Q)$ for $\alpha(Q|k, \tilde{k})$ etc.

⁵²Indeed,

$$F^2(q) = c^2(\tilde{\epsilon}|k) (\wp(\tilde{\epsilon}) - \wp(\tilde{q})) = c^2(\tilde{\epsilon}|k) \left(\frac{e_{13}}{\text{sn}^2(\sqrt{e_{13}}\tilde{\epsilon}|k)} - \frac{e_{13}}{\text{sn}^2(\sqrt{e_{13}}\tilde{q}|k)} \right) = \frac{c^2(\tilde{\epsilon}|k)e_{13}}{\text{sn}^2(q|k)} \left(1 - \frac{\text{sn}^2(\tilde{\epsilon}|k)}{\text{sn}^2(q|k)} \right)$$

where $q = 2\omega\hat{q}\sqrt{e_{13}}$ and $2\nu^2 = \text{sn}^2(\tilde{\epsilon}|k)$.

in the standard way.

The other limit $k \rightarrow 0$ (with $\tilde{k}$ finite) gives $\alpha(q|k) \rightarrow \alpha_{trig}(q) = 1 - \frac{2\nu^2}{\cosh q}$ and

$$H^{dell}(p, q) \longrightarrow \alpha_{trig}(q) \cdot \text{cn} \left(p \sqrt{\tilde{k}'^2 + \tilde{k}^2 \alpha_{trig}^2(q)} \left| \frac{\tilde{k} \alpha_{trig}(q)}{\sqrt{\tilde{k}'^2 + \tilde{k}^2 \alpha_{trig}^2(q)}} \right. \right) = \tilde{H}^R(p, q). \quad (7.162)$$

This is the elliptic-trigonometric model, dual to the conventional elliptic Ruijsenaars (i.e. the trigonometric-elliptic) system. In the further limit of small q this degenerates into the elliptic-rational model with $\alpha_{trig}(q) \rightarrow \alpha_{rat}(q) = 1 - \frac{2\nu^2}{q^2}$, which is dual to the conventional elliptic Calogero (i.e. the rational-elliptic) system, analyzed in some detail above.

Our approach has been based on choosing appropriate functions $f(q)$ and $F(Q)$ and implementing duality. Other choices of functions associated with alternative free Hamiltonians may be possible. Instead of the duality relations (7.155) one could consider those based on $h_0(p) = \text{sn}(p|\tilde{k})$ instead of $\text{cn}(p|\tilde{k})$. With this choice one gets somewhat simpler expressions for β_s and γ_s :

$$\beta_s = 1, \quad \gamma_s(q|\tilde{k}, k) = \tilde{k} \alpha_s(q|k), \quad \alpha_s(q|k) = 1 - \frac{2\nu^2}{\text{cn}^2(q|k)} \quad (7.163)$$

and the final Hamiltonian is now

$$H_s(p, q|\tilde{k}, k) = \alpha_s(q|k) \cdot \text{sn}(p|\tilde{k} \alpha_s(q|k)). \quad (7.164)$$

Although this Hamiltonian is somewhat simpler than our earlier choice, the limits involved in obtaining the Ruijsenaars-Calogero-Toda reductions are somewhat more involved, and that is why we chose to present the Hamiltonian (7.159) first.

One might further try other elliptic functions for $h_0(p)$. Every solution we have obtained by making a different ansatz has been related to our solution (7.159) via modular transformations of the four moduli $\tilde{k}$, k , $\tilde{k}_{eff} = \tilde{\gamma}$ and $k_{eff} = \gamma$.

Another approach to the 2-particle double elliptic systems was proposed in [178]. It is suggested there to start with the four dimensional manifold, namely, the elliptically fibered K3 manifold. This manifold is the space where the separated variables live. Since it is a phase space manifold, it is provided with nontrivial Poisson brackets. They follow from the natural Poisson structure on the K3 manifold. The Hamiltonian can be chosen as a linear function on the base manifold while the dual Hamiltonian as a linear function on the fiber. Using explicit formulas for the Poisson brackets of coordinates on the elliptically fibered K3 manifold, in principle, one may obtain explicit Hamiltonians. We are planning to report on these calculations as well as on equivalence of the two approaches in a separate publication [188].

7.7 Duality, τ -functions and Hamiltonian reduction

Now one should extend the results of the two-particle case to the generic number of particles. Unfortunately, it is still unknown how to get very manifest formulas. Here we describe a method that ultimately allows one to construct commuting Hamiltonians of the N -particle Dell system, but very explicit expressions will be obtained only for first several terms in a perturbation theory.

However, in order to explain the idea, we need some preliminary work. In fact, we start from the old observation [73, 189, 72, 190] that, if one study the elliptic (trigonometric, rational) solution

to the KP hierarchy, the dynamics (dependence on higher times of the hierarchy) of zeroes of the corresponding τ -function w.r.t. the first time is governed by the elliptic (trigonometric, rational) Calogero system. Moreover, if one starts with the two-dimensional Toda lattice hierarchy and studies the zeroes of the τ -function w.r.t. the zero (discrete) time, the corresponding system governing zeroes is the Ruijsenaars system [190].

Let us illustrate this with the simplest example of the trigonometric solution of the KP hierarchy. This is, in fact, the N -solitonic solution, the corresponding τ -function being

$$\tau(\{t_k\} | \{\nu_i, \mu_i, X_i\}) = \det_{N \times N} \left(\delta_{ij} - L_{ij} e^{\sum_k t_k (\mu_i^k - \nu_i^k)} \right), \quad L_{ij} \equiv \frac{X_i}{\nu_i - \mu_j} \quad (7.165)$$

Here ν_i , μ_i and X_i are the soliton parameters. Let us impose the constraint $\nu_i = \mu_i + \epsilon$. Then, the standard Hamiltonian structure of the hierarchy on the soliton solution gives rise to the Poisson brackets (in [113, 66] similar calculation was done for the sine-Gordon case)

$$\{X_i, X_j\} = -\frac{4X_i X_j \epsilon^2}{(\mu_i - \mu_j)(\mu_i - \mu_j - \epsilon)(\mu_i - \mu_j + \epsilon)}, \quad \{\mu_i, \mu_j\} = 0, \quad \{X_i, \mu_j\} = X_i \delta_{ij} \quad (7.166)$$

Note that, upon identification $\mu_i = q_i$, $X_i \equiv e^{p_i} \prod_{l \neq i} \left(\frac{1}{(q_i - q_l)^2} - \frac{1}{\epsilon^2} \right)$, the matrix L_{ij} in (7.165) becomes the rational Ruijsenaars Lax operator, with the proper symplectic structure $\{p_i, q_j\} = \delta_{ij}$.

At the same time, (7.165) can be rewritten in the form

$$\tau(\{t_k\} | \{\nu_i, \mu_i, X_i\}) \sim \det_{N \times N} \left(e^{\epsilon x} \delta_{ij} - L_{ij} e^{\sum_{k \neq 1} t_k (\mu_i^k - \nu_i^k)} \right) \quad (7.167)$$

where $x \equiv t_1$. This determinant is the generating function of the rational Ruijsenaars Hamiltonians. On the other hand, the N zeroes of the determinant as the function of ϵx are just logarithms of the eigenvalues of the Lax operator. These zeroes are governed, as we know from [73, 189] by the trigonometric Calogero system, and are nothing but the N particle coordinates in the Calogero system:

$$\tau(\{t_k\} | \{\nu_i, \mu_i, X_i\}) \sim \sum_k e^{k \epsilon x} H_k^R \sim \prod_i^N \sin(\epsilon x - q_i) \quad (7.168)$$

Therefore, the eigenvalues of the (rational Ruijsenaars) Lax operator are the exponentials e^{q_i} of coordinates in the (trigonometric Calogero) dual system, while the τ -function simultaneously is the generating function of the Hamiltonians in one integrable systems and the function of coordinates in the dual system. This is exactly the form of duality, as it was first realized by S.Ruijsenaars [62]. Its relation with τ -functions was first observed by D.Bernard and O.Babelon [66] (see also [191]).

Similarly, one can consider the N -solitonic solution in the two-dimensional Toda lattice hierarchy. Then,

$$\tau_n(\{t_k\} | \{\nu_i, \mu_i, X_i\}) = \det_{N \times N} \left(\delta_{ij} - L_{ij} \left(\frac{\mu_i}{\nu_i} \right)^n e^{\sum_k t_k (\mu_i^k - \nu_i^k) + \bar{t}_k (\mu_i^{-k} - \nu_i^{-k})} \right), \quad L_{ij} \equiv \frac{X_i}{\nu_i - \mu_j} \quad (7.169)$$

This time one should make a reduction $\nu_i = e^\epsilon \mu_i$. Then, L_{ij} becomes the Lax operator of the trigonometric Ruijsenaars, and the zeroes w.r.t. to ϵn are governed by the same, trigonometric Ruijsenaars system. This is another check that this system is really self-dual.

Thus constructed duality can be also interpreted in terms of the Hamiltonian reduction [124]. In doing so, one starts with the moment map which typically is a constraint for two matrices or two matrix-valued functions. In order to solve it, one diagonalizes one of the matrices, its diagonal elements ultimately being functions (exponentials) of the coordinates, while the other matrix gives the Lax operator (its traces are the Hamiltonians). Now what one needs in order to construct the dual system is just to diagonalize the second matrix in order to get coordinates in the dual system, while the first matrix will provide one with the new Lax operator (its traces are the Hamiltonians in the dual system).

As an illustrative example, let us consider a Hamiltonian reduction of the trigonometric Calogero-Sutherland system which differs from that discussed in s.2.1 [60]. Now one starts from the free Hamiltonian system $H = \text{tr} A^2$ given on the phase space (A, B) of two $N \times N$ matrices, with the symplectic form $\text{tr}(\delta A \wedge B^{-1} \delta B)$ (i.e. A lies in the $gl(N)$ algebra, while B lies in the $GL(N)$ group). Then, one can impose the constraint

$$A - BAB^{-1} = \nu(\mathbf{I} - P) \quad (7.170)$$

where ν is a constant, $\mathbf{I}$ is the unit matrix and P is a matrix of rank 1. Now one has to make the Hamiltonian reduction with this constraint. First of all, one solves the constraint diagonalizing matrix B and using the gauge freedom of the system to transform P into the matrix with all entries unit. It gives⁵³ $A_{ij} = p_i \delta_{ij} + (1 - \delta_{ij}) \frac{\nu b_j}{b_i - b_j}$, where b_i are the diagonal elements of B . This is nothing but the Lax operator of the trigonometric Calogero-Sutherland model. Traces of powers of this Lax operators give the Calogero Hamiltonians.

One can also diagonalize the other matrix, A . Then, (7.170) can be rewritten in the form

$$AB - BA = \nu(B - \overline{P}) \quad (7.171)$$

where $\overline{P}$ is another rank 1 matrix which is can be gauged out to the matrix with all entries depending only on the number of row (let denote them X_i/ν). Solving then this constraint, one obtains $B = \frac{X_i}{a_i - a_j - \nu}$, where a_i are the diagonal elements of the matrix A . This is nothing but the Lax operator of the rational Ruijsenaars-Schneider system, traces of its powers give a set of the dual Hamiltonians.

Certainly, this picture [124] nicely explains the Ruijsenaars observation of duality [62] and is just the τ -function approach told in different words.

The same scheme, certainly, works in for the trigonometric Ruijsenaars too, where one can use a similar procedure in order to show that diagonalizing either of the matrices A and B , one arrives at the same system, trigonometric Ruijsenaars model, checking again that it is self-dual.

In fact, similarly to the trigonometric Calogero system, there are several equivalent ways to obtain the trigonometric Ruijsenaars model. Say, instead of the Hamiltonian reduction described in section 2, one can make the Hamiltonian reduction of the gauge theory given on a two-torus $\mathbf{T}^2 = S^1 \times S^1$ [178]. The other way of doing is to consider the Poisson reduction [65]. Namely, one introduces a set of commuting functions on the space of graph connection on a graph corresponding to a moduli space of flat connections on a torus with one hole. Being reduced to a particular symplectic leaf of the moduli space of flat connections on the torus in the symplectic structure [192], this set of functions turns out to be a full set of commuting Hamiltonians. This picture is far more geometrical and the self-duality of trigonometric Ruijsenaars system gets within this framework rather transparent geometrical meaning.

⁵³Since (7.170) does not fix the diagonal elements of A , they are parametrized by arbitrary numbers p_i .

That is, this Poisson reduction physically is nothing but the three dimensional Chern-Simons theory on $\mathbf{T}^2 \times \mathbf{R}^1$ (the line gives the temporal variable) with the insertion of an appropriate temporal Wilson line and spatial Wilson loop. It is the freedom to choose the place of the latter one that is in charge of the duality. The group of (self-) dualities of this model is therefore generated by the (modular) group acting on the non-trivial torus cycles.

7.8 Dual Hamiltonians for many-body systems

Unfortunately, the scheme described in the previous subsection does not that immediately applied to elliptic models with arbitrary number of particles. The reason is that even having the dual Lax operator calculated, one typically meets in elliptic cases a problem of constructing the dual Hamiltonians, i.e. invariant combinations involving the Lax operator.

Instead, in order to obtain dual Hamiltonians, here we use the other approach discussed above, that dealing with zeroes of the τ -functions [92, 93]. This approach gives no very explicit form of the dual Hamiltonians. Therefore, we construct later a kind of perturbative procedure for the Hamiltonians which allows one to get them absolutely explicitly term by term [93].

Consider $SU(N)$ ($N = g + 1$) system. The whole construction is based on the fact that the spectral curve of the original integrable system (Toda chain, Calogero, Ruijsenaars or the most interesting double elliptic system) has a period matrix $T_{ij}(\vec{a})$, $i, j = 1, \dots, N$ with the special property:

$$\sum_{j=1}^N T_{ij}(a) = \tau, \quad \forall i \quad (7.172)$$

where τ does not depend on a . As a corollary, the genus- N theta-function is naturally decomposed into a linear combination of genus- g theta-functions:

$$\Theta^{(N)}(p_i | T_{ij}) = \sum_{n_i \in \mathbb{Z}} \exp \left(i\pi \sum_{i,j=1}^N T_{ij} n_i n_j + 2\pi i \sum_{i=1}^N n_i p_i \right) = \sum_{k=0}^{N-1} \theta_{[\frac{k}{N}, 0]}(N\zeta | N\tau) \cdot \check{\Theta}_k^{(g)}(\check{p}_i | \check{T}_{ij}) \quad (7.173)$$

where

$$\Theta_k \equiv \check{\Theta}_k^{(g)}(\check{p}_i | \check{T}_{ij}) \equiv \sum_{\substack{n_i \in \mathbb{Z} \\ \sum_{i=1}^N n_i = k}} \exp \left(-i\pi \sum_{i,j=1}^N \check{T}_{ij} n_{ij}^2 + 2\pi i \sum_{i=1}^N n_i \check{p}_i \right) \quad (7.174)$$

and $p_i = \zeta + \check{p}_i$, $\sum_{i=1}^N \check{p}_i = 0$; $T_{ij} = \check{T}_{ij} + \frac{\tau}{N}$, $\sum_{j=1}^N \check{T}_{ij} = 0$, $\forall i$.

Now we again use the argument that zeroes of the KP (Toda) τ -function (i.e. essentially the Riemannian theta-function), associated with the spectral curve (7.191) are nothing but the coordinates q_i of the original (Calogero, Ruijsenaars) integrable system. In more detail, due to the property (7.172), $\Theta^{(N)}(p|T)$ as a function of $\zeta = \frac{1}{N} \sum_{i=1}^N p_i$ is an elliptic function on the torus $(1, \tau)$ and, therefore, can be decomposed into an N -fold product of the genus-one theta-functions. Remarkably, their arguments are just $\zeta - q_i$:

$$\Theta^{(N)}(p|T) = c(p, T, \tau) \prod_{i=1}^N \theta(\zeta - q_i(p, T) | \tau) \quad (7.175)$$

(In the case of the Toda chain when $\tau \rightarrow i\infty$ this “sum rule” is implied by the standard expression for the individual e^{q_i} through the KP τ -function.) Since one can prove that q_i form a Poisson-commuting

set of variables with respect to the symplectic structure (7.176) [73], this observation indirectly justifies the claim [92, 93] that all the ratios Θ_k/Θ_l (in order to cancel the non-elliptic factor $c(p, T, \tau)$) are Poisson-commuting with respect to the Seiberg-Witten symplectic structure

$$\sum_{i=1}^N dp_i^{Jac} \wedge da_i \quad (7.176)$$

(where $a_N \equiv -a_1 - \dots - a_{N-1}$, i.e. $\sum_{i=1}^N a_i = 0$):

$$\left\{ \frac{\Theta_k}{\Theta_l}, \frac{\Theta_m}{\Theta_n} \right\} = 0 \quad \forall k, l, m, n \quad (7.177)$$

It can be equivalently written as

$$\Theta_i \{\Theta_j, \Theta_k\} + \Theta_j \{\Theta_k, \Theta_i\} + \Theta_k \{\Theta_i, \Theta_j\} = 0 \quad \forall i, j, k$$

or

$$\{\log \Theta_i, \log \Theta_j\} = \left\{ \log \frac{\Theta_i}{\Theta_j}, \log \Theta_k \right\} \quad \forall i, j, k$$

The Hamiltonians of the dual integrable system can be chosen in the form $H_k = \Theta_k/\Theta_0$, $k = 1, \dots, g$. However, these dual Hamiltonians are not quite manifest, since they depend on the period matrix of the original system expressed in terms of its action variables. Nevertheless, what is important, one can work with the θ -functions as with series and, using the instantonic expansion for the period matrix T_{ij} , construct Hamiltonians term by term in the series.

7.9 Dual Hamiltonians perturbatively

Let us see how it really works. We start with the simplest case of the Seiberg-Witten family, the periodic Toda chain.

Perturbative approximation. First we consider the perturbative approximation which is nothing but the open Toda chain, see (5.16) and (5.17). This is the case of the perturbative $4d$ pure $N = 2$ SYM theory with the prepotential

$$\mathcal{F}_{pert}(a) = \frac{1}{2i\pi} \sum_{i < j}^N a_{ij}^2 \log a_{ij} \quad (7.178)$$

Then, the period matrix is singular and only finite number of terms survives in the series for the theta-function:

$$\Theta^{(N)}(p|T) = \sum_{k=0}^{N-1} e^{2\pi i k \zeta} H_k^{(0)}(p, a), \quad (7.179)$$

$$H_k^{(0)}(p, a) = \sum_{I, [I]=k} \prod_{i \in I} e^{2\pi i p_i} \prod_{j \in \bar{I}} \mathcal{Z}_{ij}^{(0)}(a) \quad (7.180)$$

Here

$$\mathcal{Z}_{ij}^{(0)}(a) = e^{-i\pi T_{ij}^{(0)}} = \frac{\Lambda_{QCD}}{a_{ij}} \quad (7.181)$$

and I are all possible partitions of N indices into the sets of $k = [I]$ and $N - k = [\overline{I}]$ elements. Parameter Λ_{QCD} becomes significant only when the system is deformed: either non-perturbatively or to more complex systems of the Calogero–Ruijsenaars–double-elliptic family. The corresponding τ -function $\Theta^{(N)}(p|T)$, eq.(7.179), describes an N -soliton solution to the KP hierarchy and is equal to the determinant (7.165). The Hamiltonians $H_k^{(0)}$, eq.(7.180) dual to the open Toda chain, are those of the degenerated rational Ruijsenaars system, and they are well-known to Poisson-commute with respect to the relevant Seiberg-Witten symplectic structure (7.176). This is not surprising since the open Toda chain is obtained by a degeneration from the trigonometric Calogero system (s.3.3) which is dual to the rational Ruijsenaars.

The same construction for other perturbative Seiberg-Witten systems ends up with the Hamiltonians of more sophisticated systems.

For the spectral curves (5.71) of the trigonometric Calogero system (perturbative $4d \mathcal{N} = 4$ SYM with SUSY softly broken down to $\mathcal{N} = 2$ by the adjoint mass M) the Poisson-commuting (with respect to the same (7.176)), whose perturbative prepotential is given by (5.70), Hamiltonians $H_k^{(0)}$ are given by (7.180) with

$$\mathcal{Z}_{ij}^{(0)}(a) = \frac{\sqrt{a_{ij}^2 - M^2}}{a_{ij}} \quad (7.182)$$

i.e. are the Hamiltonians of the rational Ruijsenaars system which are really dual to the trigonometric Calogero model.

For the spectral curve (5.67) of the trigonometric Ruijsenaars system (perturbative $5d \mathcal{N} = 2$ SYM compactified on a circle with an ϵ twist as the boundary conditions) whose perturbative prepotential is given by (5.70), the Hamiltonians are given by (7.180) with

$$\mathcal{Z}_{ij}^{(0)}(a) = \frac{\sqrt{\sinh(a_{ij} + \epsilon) \sinh(a_{ij} - \epsilon)}}{\sinh a_{ij}} \quad (7.183)$$

i.e. are the Hamiltonians of the trigonometric Ruijsenaars system, which is, indeed, self-dual.

Finally, for the perturbative limit of the most interesting self-dual double-elliptic system (the explicit form of its spectral curves is yet unknown except the case of two particles, see the next subsection) the relevant Hamiltonians are those of the elliptic Ruijsenaars system, given by the same (7.180) with

$$\mathcal{Z}_{ij}^{(0)}(a) = \sqrt{1 - \frac{2g^2}{\text{sn}_\tau^2(a_{ij})}} \sim \frac{\sqrt{\theta(\widehat{a}_{ij} + \varepsilon|\widetilde{\tau})\theta(\widehat{a}_{ij} - \varepsilon|\widetilde{\tau})}}{\theta(\widehat{a}_{ij}|\widetilde{\tau})} \quad (7.184)$$

where $\widetilde{\tau}$ is the modulus of the second torus associated with the double elliptic system and $\widehat{a}_{ij}$ are a_{ij} rescaled, [109].

In all these examples $H_0^{(0)} = 1$, the theta-functions $\Theta^{(N)}$ are singular and given by determinant (solitonic) formulas with finite number of items (only terms with $n_i = 0, 1$ survive in the series expansion of the theta-function), and Poisson-commutativity of arising Hamiltonians is analytically checked within the theory of Ruijsenaars integrable systems.

Beyond the perturbative limit. Beyond the perturbative limit, the analytical evaluation of $\Theta^{(N)}$ becomes less straightforward.

The non-perturbative deformation (3.6) of the curve (5.16), is associated with somewhat sophisticated prepotential of the periodic Toda chain,

$$\mathcal{F}(a) = \frac{1}{4\pi i} \sum_{i < j}^N a_{ij}^2 \log \frac{a_{ij}}{\Lambda} + \sum_{k=1}^{\infty} \Lambda^2 \mathcal{F}^{(k)}(a) \quad (7.185)$$

The period matrix is

$$\begin{aligned} \pi i T_{ij} &= \frac{\partial^2 \mathcal{F}}{\partial a_i \partial a_j} = -2\pi i \tau \delta_{ij} + \log \frac{a_{ij}}{\Lambda} + \sum_{k=1}^{\infty} \Lambda^2 \frac{\partial^2 \mathcal{F}^{(k)}(a)}{\partial a_i \partial a_j}, \quad i \neq j, \\ T_{ii} &= -2\tau - \sum_{j \neq i} T_{ij} \end{aligned} \quad (7.186)$$

and τ in this case can be removed by the rescaling of Λ . Then,

$$\begin{aligned} \Theta^{(N)}(p|T) &= \sum_{k=0}^{N-1} e^{2\pi i k \zeta} \Theta_k(p, a) = \sum_{k=0}^{N-1} e^{2\pi i k \zeta} \sum_{n_i, \sum_i n_i = k} e^{-i\pi \sum_{i < j} T_{ij} n_{ij}^2} e^{2\pi i \sum_i n_i p_i} = \\ &= \left(1 + \sum_{i \neq j}^N e^{2\pi i(p_i - p_j)} \mathcal{Z}_{ij}^4 \prod_{k \neq i, j} \mathcal{Z}_{ik} \mathcal{Z}_{jk} + \right. \\ &+ \sum_{i \neq j \neq k \neq l}^N e^{2\pi i(p_i + p_j - p_k - p_l)} \mathcal{Z}_{ik}^4 \mathcal{Z}_{il}^4 \mathcal{Z}_{jk}^4 \mathcal{Z}_{jl}^4 \prod_{m \neq i, j, k, l} \mathcal{Z}_{im} \mathcal{Z}_{jm} \mathcal{Z}_{km} \mathcal{Z}_{lm} + \dots \Big) + \\ &+ e^{2\pi i \zeta} \left(\sum_{i=1}^N e^{2\pi i p_i} \prod_{j \neq i} \mathcal{Z}_{ij} + \sum_{i \neq j \neq k} e^{2\pi i(p_i + p_j - p_k)} \mathcal{Z}_{ik}^4 \mathcal{Z}_{jk}^4 \prod_{l \neq i, j, k} \mathcal{Z}_{il} \mathcal{Z}_{jl} \mathcal{Z}_{kl} + \right. \\ &+ \sum_{i \neq j} e^{2\pi i(2p_i - p_j)} \mathcal{Z}_{ij}^9 \prod_{k \neq i, j} \mathcal{Z}_{ik}^4 \mathcal{Z}_{jk} + \dots \Big) + \\ &+ e^{4\pi i \zeta} \left(\sum_{i \neq j} e^{2\pi i(p_i + p_j)} \prod_{k \neq i, j} \mathcal{Z}_{ik} \mathcal{Z}_{jk} + \sum_i e^{4\pi i p_i} \prod_{k \neq i} \mathcal{Z}_{ik}^4 + \dots \right) + \\ &+ \dots \end{aligned} \quad (7.187)$$

with

$$\mathcal{Z}_{ij} = e^{-i\pi T_{ij}} = \mathcal{Z}_{ij}^{(0)} \left(1 - i\pi \frac{\partial^2 \mathcal{Z}^{(1)}(a)}{\partial a_i \partial a_j} + \dots \right), \quad i \neq j \quad (7.188)$$

The first few corrections $\mathcal{Z}^{(k)}$ to the prepotential are explicitly known in the Toda-chain case [193], for example,

$$\mathcal{Z}^{(1)} = -\frac{1}{2i\pi} \sum_{i=1}^N \prod_{k \neq i} \left(\mathcal{Z}_{ik}^{(0)} \right)^2 \quad (7.189)$$

The coefficients Θ_k in (7.187) are expanded into powers of $(\Lambda/a)^{2N}$ and the leading (zeroth-order) terms are exactly the perturbative expressions (7.180). Thus, the degenerated Ruijsenaars Hamiltonians (7.180) are just the perturbative approximations to the $H_k = \Theta_k/\Theta_0$ – the full Hamiltonians of the integrable system dual to the Toda chain.

The same expansion can be constructed for the other, elliptic systems. Note that the period matrix in these cases is expanded into powers of $q = e^{2\pi i\tau}$:

$$\pi i T_{ij} = \frac{\partial^2 \mathcal{F}}{\partial a_i \partial a_j} = -2\pi i \tau \delta_{ij} + \log \frac{a_{ij}}{\Lambda} + \mu^2 \sum_{k=1}^{\infty} q^k \frac{\partial^2 \mathcal{F}^{(k)}(a)}{\partial a_i \partial a_j}, \quad i \neq j \quad (7.190)$$

and the dimensional constant μ depends on the system. E.g., in the Calogero model $\mu = iM$ etc. The limit to the Toda system corresponds to $q \rightarrow 0$, $q\mu^{2N} = \text{fixed}$.

Note that the first (instanton-gas) correction to the prepotential is indeed known to be of the same universal form (7.189) not only for the Toda chain, but also for the Calogero system [154]. For the Ruijsenaars and double-elliptic systems eq.(7.189) is not yet available in the literature.

In [93] the perturbative expansions for the Hamiltonians were tested for Poisson commutativity, up to few first terms and small number of particles N . In particular, it was checked that (7.189) is true for both the Ruijsenaars and Dell systems.

The main lesson we could get from this consideration is that if the *universal* expressions like (7.189) in terms of perturbative $\mathcal{Z}_{ij}^{(0)}$, the same for all the systems, will be found for higher corrections to the prepotentials⁵⁴, this will immediately give an explicit (although not the most appealing) construction of the Hamiltonians dual to the Calogero and Ruijsenaars models and, especially important, the self-dual double-elliptic system which was explicitly constructed above only for $N = 2$.

Let us note now that the $SU(2)$ Hamiltonian (7.159) has a rather nice form in terms of the p, q variables or their duals P, Q . However, this our general $SU(N)$ construction of the Hamiltonians as ratios of genus g θ -functions uses another kind of canonical variables – angle-action variables p_i^{Jac}, a_i . The *flat* moduli a_i play the central role in Seiberg-Witten theory theory, while Q_i are rather generalizations of the algebraic moduli.

It is an interesting open problem to express the Hamiltonians in terms of P_i and Q_i , perhaps, they can acquire a more transparent form, like it happens for $SU(2)$. Another option is to switch to the “separated” variables $\mathcal{P}_i, \mathcal{Q}_i$ [179] such that $\oint_{A_j} \mathcal{P}_i d\mathcal{Q}_i = a_i \delta_{ij}$ (while generically $a_i = \oint_{A_j} \sum_i \mathcal{P}_i d\mathcal{Q}_i$). Generically, they are different from P_i, Q_i but coincide with P, Q in the case of $SU(2)$.

7.10 Dell systems: 6d adjoint matter

Thus, we have constructed the Dell system and a set of dualities that acts on the Toda-Calogero-Ruijsenaars-Dell family. They are all put together in Fig.2.

⁵⁴ To avoid possible confusion, the recurrent relation [157, 82, 149, 194] for the Toda-chain prepotential,

$$\frac{\partial^2 \mathcal{F}}{\partial \log \Lambda^2} \sim \frac{\partial^2 \mathcal{F}}{\partial \log \Lambda \partial a_i} \frac{\partial^2 \mathcal{F}}{\partial \log \Lambda \partial a_j} \frac{\partial^2}{\partial p_i \partial p_j} \log \Theta_0 \Big|_{p=0}$$

does *not* immediately provide such *universal* expressions. Already for $\mathcal{Z}^{(1)}$ this relation gives

$$\mathcal{Z}^{(1)} \sim \sum_{i < j}^N \left(\mathcal{Z}_{ij}^{(0)} \right)^2 \prod_{k \neq i, j}^N \mathcal{Z}_{ik}^{(0)} \mathcal{Z}_{jk}^{(0)}$$

which coincides with (7.189) for the Toda-chain $\mathcal{Z}_{ij}^{(0)}$, eq.(7.181), but is not true (in variance with (7.189)) for Calogero $\mathcal{Z}_{ij}^{(0)}$, eq.(7.182). Meanwhile, the recurrent relations of ref.[195] for the Calogero system provide more promising expansion.

All this emphasizes the need to study extension of the Whitham theory and the WDVV equations from the Toda chains to the Calogero and Ruijsenaars systems.

coordinate momentum	rational	trigonometric	elliptic
rational	rational Calogero	trigonometric Calogero	elliptic Calogero
trigonometric	rational Ruijsenaars	trigonometric Ruijsenaars	elliptic Ruijsenaars
elliptic	dual Calogero	dual Ruijsenaars	Dell system

Figure 2: Action of the coordinate-momentum duality on the Calogero-Ruijsenaars-Dell family. Hooked arrows mark self-dual systems. The duality leaves the coupling constant ν intact.

Since the theories of this family describes low energy limits of different SUSY gauge theories, this table describes the dualities between different gauge theories too. Say, perturbative limit of the $4d$ theory (trigonometric Calogero) is dual to a special degeneration [148] of the perturbative limit of the $5d$ theory (rational Ruijsenaars), while full, *non-perturbative* $5d$ theory (elliptic Ruijsenaars) is dual to the perturbative limit of the $6d$ theory (Dell system).

To conclude our discussion of dualities and Dell systems, we calculate the perturbative prepotential of the 2-particle Dell system (7.159) and show that it coincides with what is to be expected for the $6d$ theory with adjoint matter, supporting the identification of the Dell system with the $6d$ gauge theory.

Let us start from the full Dell model for 2-particles. Then, it is given by the Hamiltonian (7.159) parametrized by two independent (momentum and coordinate) elliptic curves with elliptic moduli k and $\tilde{k}$. As usual, for two particles we can construct the *full* spectral curve from the Hamiltonian ([37], see also ss.3.2, s.3.7 and (7.127)),

$$H(\zeta, \xi|k, \tilde{k}) = u \quad (= \text{cn}(Q|k)) \quad (7.191)$$

It is characterized by the effective elliptic moduli

$$k_{eff} = \frac{k\alpha(q|\tilde{k})}{\beta(q|k, \tilde{k})}, \quad \tilde{k}_{eff} = \frac{\tilde{k}\alpha(q|k)}{\beta(q|\tilde{k}, k)} \quad (7.192)$$

where the functions α and β are manifestly given in (7.157). Coordinate-momentum duality interchanges $k \leftrightarrow \tilde{k}$, $k_{eff} \leftrightarrow \tilde{k}_{eff}$ (and $q, p \leftrightarrow Q, P$), while in general $SU(N)$ case, the model describes an interplay between the four tori: the two bare elliptic curves and two effective Jacobians of complex dimension $g = N - 1$.

From our previous discussions it should be clear that the generating differential in $6d$ theories is of the form $dS = \zeta d\xi$, with ξ living on the coordinate torus and ζ – on the momentum torus.

Now we calculate the leading order of the prepotential expansion in powers of $\tilde{k}$ when the bare spectral torus degenerates into sphere. In the forthcoming calculation we closely follow the line of s.5.6.

When $\tilde{k} \rightarrow 0$, $\text{sn}(q|\tilde{k})$ degenerates into the ordinary sine. For further convenience, we shall parameterize the coupling constant $2\nu^2 \equiv \text{sn}^2(\epsilon|k)$. Now the spectral curve (7.191) acquires the form

$$\alpha(\xi) \equiv \sqrt{1 - \frac{\text{sn}^2(\epsilon|k)}{\sin^2 \xi}} = \frac{u}{\text{cn}\left(\zeta\beta \middle| k_{eff}\right)} \quad (7.193)$$

Here the variable ξ lives in the cylinder produced after degenerating the bare coordinate torus. So does the variable $x = 1/\sin^2 \xi$. Note that the A -period of the dressed torus shrinks on the sphere to a contour around $x = 0$. Similarly, B -period can be taken as a contour passing from $x = 0$ to $x = 1$ and back.

The next step is to calculate variation of the generating differential $dS = \eta d\xi$ w.r.t. the modulus u in order to obtain a holomorphic differential:

$$dv = \left(-i\text{sn}(\epsilon|k)\sqrt{k'^2 + k^2u^2}\right)^{-1} \frac{dx}{x\sqrt{(x-1)(U^2-x)}} \quad (7.194)$$

where $U^2 \equiv \frac{1-u^2}{\text{sn}^2(\epsilon|k)}$. Since

$$\frac{\partial a}{\partial u} = \oint_{x=0} \frac{\partial dS}{\partial u} = \oint_{x=0} dv = -\frac{1}{\sqrt{(1-u^2)(k'^2 + k^2u^2)}} \quad (7.195)$$

we deduce that $u = \text{cn}(a|k)$ and $U = \frac{\text{sn}(a|k)}{\text{sn}(\epsilon|k)}$. The ratio of the B - and A -periods of dv gives the period matrix

$$T = \frac{U}{\pi} \int_0^1 \frac{dx}{x\sqrt{(x-1)(U^2-x)}} = -\frac{1}{i\pi} \lim_{\kappa \rightarrow 0} \left(\log \frac{\kappa}{4}\right) + \frac{1}{i\pi} \log \frac{U^2}{1-U^2} \quad (7.196)$$

where κ is a small- x cut-off. The U dependent part of this integral is finite and can be considered as the “true” perturbative correction, while the divergent part just renormalizes the bare “classical” coupling constant τ , *i.e.* classical part of the prepotential (see s.5.6 for further details). Therefore, the perturbative period matrix is finally

$$T_{finite} = \frac{i}{\pi} \log \frac{\text{sn}^2(\epsilon|k) - \text{sn}^2(a|k)}{\text{sn}^2(a|k)} + \text{const} \longrightarrow \frac{i}{\pi} \log \frac{\theta_1(a+\epsilon)\theta_1(a-\epsilon)}{\theta_1^2(a)} \quad (7.197)$$

and the perturbative prepotential is the elliptic tri-logarithm. Note that it is again of the form (5.70) and (5.69) but with the function $f^{(6)}$ instead of $f^{(4)}$ and $f^{(5)}$. Remarkably, it lives on the *bare* momentum torus, while the modulus of the perturbative curve (7.193) is the dressed one.

7.11 Quantum duality

Our discussion so far concerned the classical systems. However duality is expected to be a powerful tool to deal with quantum problems too. This point has been first recognized by Ruijsenaars in his early papers. The very idea is that Hamiltonians and their duals have common wave-functions and this fact could be exploited to consider one or another set of eigenvalue problems, with respect to different variables. Since the wave functions in the systems under consideration can be realized in group terms, one could expect that quantum duality can be formulated purely in group terms too. Recent activity in this direction supports such expectations [196].

Here we work out a couple of simple quantum 2-particle examples of dual systems in order to illustrate the general picture. Let us start with the most standard oscillator example with the Hamiltonian

$$\hat{H} = -1/2 \frac{\partial^2}{\partial q^2} + \frac{\omega^2 q^2}{2} \quad (7.198)$$

Its normalized eigenfunctions are

$$\begin{aligned} \hat{H}\psi_n &= \omega(n + 1/2)\psi_n \\ \psi_n(q) &= \left(\frac{\omega}{\pi}\right)^{1/4} \frac{e^{-\frac{\omega q^2}{2}}}{2^{n/2} \sqrt{n!}} H_n(q\sqrt{\omega}) \end{aligned} \quad (7.199)$$

where $H_n(\xi)$ are the Hermite polynomials $H_n(\xi) = e^{\xi^2} (-\partial_\xi)^n e^{-\xi^2}$. Using this representation for the wave-function one can easily obtain a recurrence relation

$$\sqrt{n+1}\psi_{n+1}(q) + \sqrt{n}\psi_{n-1}(q) = \sqrt{2\omega}q\psi_n(q) \quad (7.200)$$

It means that $\psi_n(q)$ is the eigenfunction of the difference operator

$$\hat{H}_D = T_+ \sqrt{n} + \sqrt{n} T_-, \quad T_\pm = e^{\pm \frac{\partial}{\partial n}} \quad (7.201)$$

acting on the subscript n , which is nothing but the quantized dual Hamiltonian.

Less trivial example whose classical version we discussed throughout this whole section is the quantum (2-particle) Hamiltonian of the trigonometric Calogero model

$$\hat{H} = -\frac{1}{2} \frac{\partial^2}{\partial q^2} + \frac{\nu(\nu-1)}{2 \sin^2(q)} \quad (7.202)$$

Its normalized eigenfunctions are

$$\begin{aligned} \hat{H}\psi_n &= \frac{n^2}{2}\psi_n \\ \psi_n(q) &= \sin^\nu(q) \sqrt{n \frac{(n-\nu)!}{(n+\nu-1)!}} \Pi_{n-\frac{1}{2}}^{\nu-\frac{1}{2}}(\cos(q)) \\ \Pi_l^m(x) &= \frac{1}{l!} \partial_x^{l+m} \left(\frac{x^2-1}{2} \right)^l \end{aligned} \quad (7.203)$$

For the sake of simplicity, we take half-integer ν and n . One can make the change $\nu \rightarrow -\nu-1$ and get another eigenfunction with the same eigenvalue. Using the fact that the generating function for Π_l^0 's

$$Z(y, x) = \sum_{l=0}^{\infty} y^l \Pi_l^0 = \frac{1}{\sqrt{1-2xy+y^2}} \quad (7.204)$$

satisfies the two obvious equations

$$\begin{aligned} (x-y)\partial_x Z &= y\partial_y Z \\ (1-2xy+y^2)\partial_y Z &= (x-y)Z \end{aligned} \quad (7.205)$$

which implies

$$(y\partial_y - m)\partial_x^m Z = (x-y)\partial_x^{m+1} Z \quad (7.206)$$

and, hence, yields

$$((1 - 2xy + y^2)\partial_y + y - x)\partial_x^m Z = m(1 + 2y\partial_y)\partial_x^{m-1} Z \quad (7.207)$$

one derives the recurrence relations

$$\begin{aligned} x\Pi_l^m &= \frac{l+1-m}{2l+1}\Pi_{l+1}^m + \frac{l+m}{2l+1}\Pi_{l-1}^m \\ \cos(q)\psi_n &= \frac{1}{2}\left(\sqrt{\frac{1-\nu(\nu-1)}{n(n+1)}}\psi_{n+1} + \sqrt{\frac{1-\nu(\nu-1)}{n(n-1)}}\psi_{n-1}\right) \end{aligned} \quad (7.208)$$

i.e. ψ_n is an eigenfunction of the finite-difference operator acting on the subscript n

$$\hat{H}_D\psi(q) = \cos(q)\psi(q) \quad (7.209)$$

$$\hat{H}_D = T_+ \sqrt{1 - \frac{\nu(\nu-1)}{n(n-1)}} + \sqrt{1 - \frac{\nu(\nu-1)}{n(n-1)}} T_- \quad (7.210)$$

which is a quantum version of the rational Ruijsenaars model.

Summarizing, when the system with trigonometric dependence on momentum is quantized, its Hamiltonian becomes a finite-difference operator. The wave-functions become the functions of discrete variables. The origin of this is in the Bohr-Sommerfeld quantization condition. Indeed, since the trigonometric dependence on momenta implies that the leaves of polarization are compact and, moreover, non-simply connected, covariantly constant sections of the prequantization connection along the polarization fiber generically ceases to exist. It is only for special “quantized” values of the action variables that the section exists. In the elliptic case, the quantum dual Hamiltonian is going to be a difference operator of infinite order.

8 Branes as low energy degrees of freedom in supersymmetric gauge theories

8.1 Field theory on D-brane worldvolume

We begin this section with the description of nonperturbative states in string theory which play the key role for constructing the low-energy degrees of freedom in SUSY gauge theories. In particular, the phase space of the integrable systems, considered in the previous sections is associated with the collective coordinates of branes of different dimensions. Besides it, nonperturbative dualities in string theory suggest several different formulations for the same single field theory.

The central objects introduced into string theory by exact solving the $\mathcal{N} = 2$ SYM theory are the duality (between the strong and weak coupling regimes) and the exact spectrum of BPS solitons. It appeared that in order to consistently describe duality in string theory, one has to introduce solitonic objects of different dimensions, “p-branes” with p space coordinates [43]. In quasi-classical approximation, their masses behave as $m = g^{-2}$, where g is the string coupling constant, and the mass formulae for the BPS states, being related to the central extension of the SUSY algebra remains exact at quantum level. At the strong coupling regime, the solitons become the lightest objects in the theory and define the dynamics in that region. Technically, it is possible to introduce dual variables which yield a weak coupling theory in terms of the solitonic fields.

All this picture is literally inherited from Seiberg-Witten theory. On the other hand, Seiberg-Witten theory itself gets a new description in string/brane terms. Indeed, the appearance of Riemann surfaces in Seiberg-Witten theory can be explained in terms of branes. It is this surface which is the part of the worldvolume of the larger brane. The structure of the brane intersections is fixed by the SUSY and the set of configurations yielding the field theory become finite or even single.

As we already explained in the Introduction, the key point which admits the use of branes for obtaining field theories is the $U(1)$ gauge theory on the brane worldvolume. It is known that branes are charged with respect to the RR sector of the theory [43] and can serve as a space of endpoints of strings. Using this property Witten formulated the new view on the gauge theory as the theory on the worldvolume of coinciding branes which carry $U(1)$ factor each. The open strings connecting branes give rise to non-Abelian theory, rank of the gauge group being defined by the number of branes. In what follows we shall follow this line: SUSY field theories are defined on the brane worldvolume and the parameters are introduced via the geometry of the brane configuration.

Some parameters of $\mathcal{N} = 2$ SUSY theories (condensates of scalars) coordinatize the Coulomb branch of the corresponding vacuum moduli space, while other parameters (bare masses and coupling constants) remain external parameters. All these parameters are defined in the brane terms (see the Introduction): scalar condensates yield the distance between branes (or determine embedding of the single M-theory brane into the target space), coupling constant is just the distance between additional, background branes (or also determines geometry of the M-theory brane), while the hypermultiplet mass is the coordinate of another brane (often of different dimension) in the proper direction. In what follows we shall present examples of particular brane configurations. Coordinates of branes appear to be associated with the degrees of freedom of the integrable system governing the low energy action in the SUSY gauge theory.

Let us emphasize that the gauge theories of the Hitchin type responsible for the integrable many-body systems arise in the “momentum” space or “the space of fields” and should not be thought of as those obtained by the dimensional reduction from the $4d$ SUSY gauge theory. In fact, these theories emerge from different degenerations of the toric geometry.

It is useful to start with the softly broken $\mathcal{N} = 4$ theory. In $\mathcal{N} = 1$ terms, there are 3 adjoint hypermultiplets whose complex scalar components parameterize the embedding of branes into six dimensions. The potential of scalars is

$$V = \frac{1}{g^2} \text{Tr} \sum_{i=1,2,3} [\phi_i, \phi_i^+]^2 \quad (8.1)$$

and is induced by the kinetic term (as a result of integration over auxiliary fields). There is also a non-trivial superpotential, $W = \text{Tr} \Phi_1 [\Phi_2, \Phi_3] + M^2 \text{Tr} (\Phi_2^2 + \Phi_3^2)$, where Φ_i are $\mathcal{N} = 1$ hypermultiplets with lowest components ϕ_i respectively and M is the mass of the hypermultiplet. The minimum of the potential energy amounts to the vacuum configuration with non-trivial commutation relations between the 3 scalars. In the large mass limit, the fluctuations orthogonal to the algebraic surface in the two dimensional complex space

$$\text{Tr} (\Phi_2^2 + \Phi_3^2) = \text{const}, \quad (8.2)$$

are frozen. Note that, since one of the scalars remains massless, it jointly with the gauge superfield forms one $\mathcal{N} = 2$ supermultiplet, while the two massive scalars are the components of the $\mathcal{N} = 2$ adjoint massive hypermultiplet, i.e. the $\mathcal{N} = 4$ SUSY is broken down to the $\mathcal{N} = 2$ SUSY. In this

$\mathcal{N} = 2$ theory, there are valleys for the massless scalar ϕ_1 due to the potential term (8.1), i.e. this scalar may have a condensate which parameterizes the Coloumb branch.

There are two equivalent approaches to $\mathcal{N} = 2$ theories in the string theory framework which start from the theory in ten dimensions. In the first, geometrical engineering approach, one considers the compact Calabi-Yau (CY) threefold near the singularity providing the desired gauge group [48, 54]. In the second approach proposed in [51], the theory on the worldvolume of D4-branes stretched between two NS 5-branes is considered. The location of the branes is summarized in the following table

$$\left\{ \begin{array}{l} NS5 \\ D4 \\ D6 \end{array} \right| \begin{array}{cccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ + & + & + & + & + & + & - & - & - & - \\ + & + & + & + & - & - & + & - & - & - \\ + & + & + & + & - & - & - & + & + & + \end{array} \quad (8.3)$$

where plus stands for the extended direction of each brane and minus for the transverse direction. The fourbranes are finite in the x_6 direction stretching between the two NS branes, therefore, the long range theory is a four-dimensional theory as we need. While in this type IIA picture the original SUSY (32 real supercharges) is broken down to $N = 2$ in $4d$ (8 real supercharges) due to the concrete brane configuration⁵⁵, in the IIB picture (related to CY), the original SUSY is broken via geometry of the CY manifold. D-branes wrapped around shrinking cycles provide the field theory and break half of the supersymmetries indeed. Note that the IIA picture can be lifted to M-theory in $11d$ which has the $11d$ SUGRA as its low-energy limit.

8.2 Branes and degrees of freedom in many-body systems

Let us start with the description of degrees of freedom for the $4d$ $SU(N_c)$ theory without matter in IIB/F picture [197]. It emerges as the worldvolume theory on D7-branes wrapped around an elliptically fibered K3 [54]. The bare coupling of the gauge theory yields the complex structure of the elliptic fiber, therefore, the spectral curve of the Calogero model enters the picture automatically.

To identify the degrees of freedom using the approach of section 2.3, one has to define the holomorphic one-form ϕ as well as the anti-holomorphic vector potential $\overline{A}$ on the spectral curve. The holomorphic form can be identified if one takes into account that the torus is embedded into the K3 space as a supersymmetric cycle. Therefore the embedding of D7 branes is described in terms of the one-form [56] which after all turns out to be the Lax operator for the many-body system. The anti-holomorphic vector field $\overline{A}$ whose eigenvalues are the coordinates of particles in the many-body system comes from the N_c branes wrapped around the spectral curve with the worldvolume gauge fields. Finally the marked point on the spectral curve with nontrivial monodromy is introduced via additional “background” branes localized on the spectral curve. Such “background” branes play the role of regulators in field theory. Let us emphasize that in F-theory the 11-th and 12-th dimensions just provide the bare coupling constant.

Since the background brane does not change the interpretation of degrees of freedom let us consider the simplest case of the free many-body system. The trigonometric Ruijsenaars system, i.e. the $5d$ SUSY theory is the most illustrative, since in this case the brane geometry is in one-to-one correspondence with the Hamiltonian reduction procedure of section 2. Let us remind that the system originates from the Chern-Simons theory given on the product of torus and real (time) line. There is

⁵⁵The presence of D4-branes breaks symmetry down to 16 supercharges and two NS branes breaks it down further to 8 supercharges.

the relation for the monodromies of the gauge field around the torus cycles g_A, g_B

$$g_A g_B g_A^{-1} g_B^{-1} = 1$$

In the free case the matrices can be diagonalized simultaneously and their diagonal elements represent the coordinates and momenta of the dynamical system $g_A = \text{diag}(e^{iq_i})$, $g_B = \text{diag}(e^{ip_i})$. Note that from the F-theory point of view we have degenerated the fiber torus to a cylinder.

Let us compare the geometrical descriptions following from the Chern-Simons Lagrangian and the brane configuration. Two coordinates x_5, x_6 are essential for the description of the five dimensional theory. The typical scale along x_5 is R_5^{-1} , while that along x_6 is g^{-2} in the string units [198]. We shall assume that both coordinates are compactified onto circles of the corresponding radii.

The (coordinate and momentum) circles in the Ruijsenaars model are related to the circles in the brane configuration via the T-duality transformation along the both directions. Therefore, the radii of circles in the Ruijsenaars model are R_5, g^2 , and after the T-duality transformation the eigenvalues of the monodromy matrix transform into the brane positions along the dual circle. There are precisely N_c 5-branes in the IIB theory extended along x_6 and localized at p_i along x_5 . The coordinates in the dynamical system are provided by N_c 1-branes localized at q_i along x_6 . The SUSY gauge theory is defined on $(x_0, x_1, x_2, x_3, x_4, x_6)$, and the coordinates of 5-branes along x_5 define the vacuum expectation values of the real scalar field.

To switch on the interaction in the dynamical system, one has to add a massive hypermultiplet in adjoint in the field theory language, to introduce a marked point with non-trivial monodromy around in the Chern-Simons description or to add the background branes in the string theory approach.

The new relation for the monodromy matrix is

$$g_A g_B g_A^{-1} g_B^{-1} = g_c,$$

where g_c corresponds to inserting the $\mathbb{C}P^{n-1}$ orbit at the marked point. After the T-duality transformation, a brane configuration with non-trivial flux arises. The color orientation of the fluxes follows from the structure of the monodromy matrix g_c .

Another approach to description of field theories, for definiteness let them be $4d$ theories, is based on the consideration of field theory on the worldvolume of M5-brane, the solitonic state in M-theory [51]. We look for an M5-brane configuration that provides a theory with 8 real supercharges, i.e. in $\mathcal{N} = 2$ theory. This can be achieved by considering the product of R^4 and a fixed genus compact surface Σ as the worldvolume of M5-brane. After compactification onto the circle (11-th dimension), the M5-brane in the IIA theory looks as a collection of D4-branes located between two NS-branes.

To get the $SU(N_c)$ theory, one has to introduce N_c D4-branes of finite size along one of the coordinates, say, x_6 . Then, the plane $x_4 + ix_5$ in the perturbative regime is where the momentum variables in the many-body system live. Moreover, there are D0-branes on the D4-worldvolume, one per each D4-brane, which are ordered along the x_6 coordinate. In the low-energy limit, the D0-brane coordinates are defined

$$q_j = j\Delta + \phi_j, \tag{8.4}$$

where Δ fixes their equilibrium locations, while ϕ_i correspond to small fluctuations and coincide with the variables in the Toda system. Let us note that q_i can be also considered as the coordinates in the Calogero system, without any ordering. Comparing with the field theory variables one immediately

arrives at $N_c \Delta = \frac{1}{g^2(M)}$ which perfectly fits the suggested ordering of the D0-branes along the x_6 direction.

A qualitative interpretation of the 2×2 Toda Lax operator is then as follows. The element L_{11} corresponds to the D0-brane confining with the D4-brane so that they have the same $(x_4 + ix_5)$ -coordinate. Since Lax operators are related with fermionic zero modes in a nontrivial background field, the size of the Lax matrix indicates the presence of the two dimensional space of zero modes. Elements L_{12}, L_{21} fix the matrix elements of the spectral parameter between the fermionic zero modes localized on the neighbour D0-branes.

Let us note once again that qualitatively in any brane configuration, there are three essential elements presented. First, there are the branes that provide the worldvolume for the field theory, N_c D7-branes in IIB/F or one M5-brane in IIA/M picture. Second, it is necessary to introduce the branes responsible for the variables in the dynamical system, which are zero size instantons in the field theory language [199]. Finally, the account of the regulators in field theory corresponds to an additional Wilson line in the IIB/F picture or fluxes in the IIA/M description.

Recently the correspondence between integrable systems and branes has been pushed forward even further. Namely, one can start with the $4d$ theory with one compact dimension [200] and map the system to the Hitchin phase space by a chain of dualities. On one hand, such a moduli space coincides with the moduli space of the non-commutative instantons on T^4 or K3 manifolds [201]. On the other hand non-commutative instantons introduced in [202] allow a deformed ADHM description, moreover, the explicit map between the moduli space of non-commutative instantons and the trigonometric Calogero model has been elaborated [203].

8.3 Branes in five dimensional theory

While considering $5d$ gauge theories the role of the brane configuration becomes the most transparent. It appears that in five dimensions the brane configuration fixes the structure of the spectral curve of the integrable system. We shall consider below the configuration of branes in the IIB theory which is related to the toric manifold data providing the compactification of M-theory [54, 48, 68].

We start from the geometrical engineering picture [98]. As a compact manifold we take the CY manifold in the vicinity of singularities. Being interested only in this region of parameters, we discuss exclusively a local model implying that its embedding into a compact Calabi-Yau is always possible. To this end, we consider A_{n-1} ALE spaces fibered over a set of the base spheres, each sphere corresponding to an $SU(n)$ gauge theory⁵⁶. The size of the base sphere $V \propto \frac{1}{g^2}$ is governed by the coupling constant of the corresponding $SU(n)$ factor. Kahler moduli of $\mathbf{P}^1$'s associated with blowing up the ALE space define the Cartan scalars a_i on the Coulomb branch of the moduli space. This non-compact Calabi-Yau space is a hypersurface in the holomorphic quotient and can be represented as a set of Kahler cones, or, in other words, by the toric polyhedron Δ .

For example, one can start from the $\mathcal{N} = (2, 2)$ linear sigma model construction of Calabi-Yau [204] and define $U(1)$ charge vectors of matter fields

$$\mathbf{Q}^a = (q_1^a, \dots, q_k^a) \tag{8.5}$$

From the σ -model point of view these vectors define gauge-invariant variables, while from the Calabi-

⁵⁶Throughout this subsection we consider the gauge theory whose gauge group is a product of $k - 1$ simple factors, typically $SU(n_i)$.

Yau standing point they act on the homogeneous coordinates in projective space and fulfill r linear relations

$$\sum_J Q_J^a \nu_J = 0 \quad (8.6)$$

between vectors $\nu_J \in \mathbf{N} = \mathbf{Z}^d$ which define coordinate patches in each Kahler cone. Vertices ν_J in lattice $\mathbf{N}$ form a toric polyhedron $\Delta = \{\nu_J\}$. For the toric variety to be smooth, Δ must be convex. Thus, having defined the original Calabi-Yau in terms of holomorphic quotients, we now turn to the solution of the model via mirror symmetry.

As it was explained in [48, 54], Kahler moduli of the original Calabi-Yau specify Coulomb moduli of the low-energy effective theory. Under mirror transform they get mapped to the complex structure of the mirror manifold, which is just Riemann surface E having the same moduli space. There is a profound reason in this symmetry, because the Riemann surface is the spectral curve of the associated integrable system, **XXZ** spin chain, section 3.5.

In terms of charge vectors this mirror manifold is

$$\prod_J y_J^{q_J^a} = 1 \quad (8.7)$$

For the models we consider here it is just a spectral curve of underlying integrable system.

Let us now turn to the description of $5d$ theories in terms of brane configurations. One can derive the brane configuration using T-duality along x_4 from the IIA picture [51]. Thus, one obtains the IIB picture [198] with the brane configuration [198]

$$\left\{ \begin{array}{c} NS5 \\ D5 \end{array} \right| \begin{array}{cccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ + & + & + & + & + & + & - & - & - & - \\ + & + & + & + & + & - & + & - & - & - \end{array} \quad (8.8)$$

where the field theory is realized on the D5-brane worldvolume. The holomorphic solution to the $2d$ Laplace equation in the space (x_4, x_5) , $\Delta s = 0$ ($s \equiv x_6 + ix_{10}$): $s = \log(x_4 + ix_5)$ transforms into the solution to the $1d$ Laplace equation $s(\equiv x^6) = \frac{1}{2}|x_5| = \frac{x_5}{2}\text{sign}(x_5)$. If one demands for SUSY to be broken up to $1/4$, a restriction on the brane intersection arises. In particular, when charged n D5-brane meet charged k NS5-branes they must merge into (k, n) bound state with the slope $\tan \phi = \frac{k}{n}$ on (x_5, x_6) plane [198].

As usual we divide moduli of brane configurations into two groups. The first group of parameters, external parameters (e.g. coupling constants, bare quark masses) define the five-dimensional theory, while the rest are dynamical moduli. In brane terms, changing the parameters of the first group one changes the asymptotic (i.e. infinite in all their non-zero dimensions) branes⁵⁷, while changing the parameters of the second group does not.

This is a good point to make a contact with the geometrical engineering approach [48, 54]. In that language each link in the toric polyhedron Δ is associated with a 1-cycle on the corresponding Riemann surface E . Periods along 1-cycles in the compact part of E (i.e. internal links in Δ) define the dynamical moduli whereas bare parameters arise from 1-cycles that wrap the non-compact part of E (i.e. related to the links on the boundary of Δ). One can go further and note that the number of such external links in Δ corresponding to some low-energy effective field theory coincide with the number of infinite fivebranes in the brane construction of the same theory.

⁵⁷In our case, NS5 branes.

The reason for this is that the brane picture and the toric polyhedron corresponding to a five-dimensional field theory are intimately related to each other. To establish other connections let us visualize brane diagrams in other terms. Any such configuration depicts the kinematics of two-dimensional (x_5, x_6) free particles. The charge conservation in the vertices where branes merge together is nothing but the momentum conservation law for free particles. The integer-valued momentum $p^{(5,6)} = (k, n)$ of particle is just the $SL(2, \mathbb{Z})$ charge of the brane. The vertical branes in terms of [198] carry the $(1, 0)$ charge (momentum) vector, and horizontal ones $-(0, 1)$. Notice that the number of vertical (NS5) branes specifies the number of simple gauge factors whereas the number of horizontal (D5) branes defines the rank of each $SU(n_i)$ factor.

For instance, for $SU(n)^k$ gauge theory there are $k + 1$ NS-branes and n D-branes in each stack. And, similarly, the number of vertical links in the toric polyhedron, i.e. the height of Δ equals $k + 1$ and the number of horizontal links, i.e. its horizontal size is n . At generic point on the moduli space, the vertices of Δ lie on a two-dimensional integer lattice. In these conventions we can endow each link in Δ with an integer-valued two-dimensional vector (k_δ, n_δ) .

One can exploit [98] the fact that type IIB brane configurations for five-dimensional field theories are in one-to-one correspondence with toric polyhedrons of Calabi-Yau M-theory compactifications describing these field theories. The exact relation between the two is provided by identification of the (k_δ, n_δ) charge (momentum) vector with a link vector in Δ . This powerful tool allows one to solve any five-dimensional $\prod SU(n)$ gauge theory with arbitrary matter.

The toric polyhedron Δ for the case of pure gauge $SU(n)$ theory can be found in [54]. Adding a massive hypermultiplet, we break one link on the boundary of Δ into two pieces and add one more internal link. Therefore, the number of internal and external links representing massive hypermultiplets increases by one. Within the brane approach, the picture is exactly the same: adding a fundamental matter hypermultiplet to the theory one attaches a new semi-infinite brane. It can be done in four different ways: between the pairs of the four semi-infinite branes already presented in the configuration.. The same possibilities can be found in breaking one of the four external links of the polyhedron into two parts. For the $\prod_{i=1}^k SU(n_i)$ case, one has to take a fibration of n_i spheres over each of k base spheres. The total number of external links is $2(k + 1)$ as the number of semi-infinite branes in the brane configuration. The number of finite branes is $\sum_{i=1}^k (3n_i - 1) - 1$ which matches exactly the number of internal links in Δ . The relation between toric diagrams and brane configurations has been also considered in [205].

8.4 Branes and equations of motion in many-body systems

Our next step in clarification of the role of brane moduli as degrees of freedom in integrable systems is to interpret the equations of motion in the brane terms [197]. To this end, let us relate the brane configuration with the monopole moduli space and identify the equations of motion with the Nahm equations defining the structure of the moduli space.

D0-branes localized on D4-branes can be considered as the degrees of freedom corresponding to the BPS monopoles. They can be related to other D4-branes by open strings. Following [206], the theory on the string worldvolume can be obtained by the dimensional reduction from the $10d$ SYM theory. Moreover, there is one-to-one correspondence between the equation defining the supersymmetric vacuum state in σ -model and the Nahm equation for monopoles. Now, in order to get the equations of motion in the Toda system, one should take into account the connection between the spectral curve in the Toda system and the spectral curve amounted from the Nahm equations for the

charge N_c cyclic monopole [207]. The spectral curve for the cyclic configuration can be derived from the generic monopole spectral curve

$$\lambda^n + \lambda^{n-1}a_1(w) + \dots + a_n(w) = 0 \quad (8.9)$$

where $a_i(w)$ is a polynomial of degree $2i$. Then, assuming the center of mass is fixed at the origin, the overall $U(1)$ phase is unit and imposing the (order N_c) cyclic group condition, one arrives at the Toda curve.

One can also connect the Lax operator for the Toda system with the Nahm matrices T_i known in the monopole theory [208]

$$T(w) = T_+ + 2T_0w + T_-w^2 \quad (8.10)$$

where $T_{\pm}$ are respectively the (strictly) upper and low triangular parts of the Toda Lax operator (3.4), while T_0 is its diagonal part⁵⁸. Adopting the notation $T_{\pm} \equiv T_1 \pm iT_2$, one can identify the Toda Lax equation (i.e. the equations of motion)

$$\frac{dT(w)}{dt} = \epsilon_{ijk}[T(w), M] \quad (8.11)$$

where $M \equiv T_+w^{-1} - T_-w$, with the Nahm equations

$$\frac{dT_i}{dt} = 2i\epsilon_{ijk}[T_j, T_k] \quad (8.12)$$

To develop a more geometrical picture, it is useful to consider fermions in the background monopole field

$$(\sigma D + \phi - t)\Psi = 0. \quad (8.13)$$

where t is an external parameter. The equation has k linear independent solutions for the magnetic charge k . It is the variable t which becomes the time variable in the Toda system and simultaneously plays the role of the space variable in the brane interpretation. The Nahm matrices in terms of the fermionic zero modes are

$$T_j = \int x_j \Psi^+ \Psi d^3x; \quad j = 1, 2, 3 \quad (8.14)$$

These expressions have the structure of the Berry connection if the momentum space is considered as the parameter space. There exists also an inverse construction which allows one to derive the multi-monopole fields starting from the auxiliary one-dimensional problem for the fermions in the external Nahm connection

$$\begin{aligned} (i\frac{d}{dt} + i\vec{\sigma}\vec{x} + \vec{\sigma}\vec{T}^+)v &= 0 \\ A_j &= \int v^+ \partial_j v dt \quad \phi = \int tv^+ v dt \end{aligned} \quad (8.15)$$

where the superscript plus of $\vec{T}$ means Hermitian conjugation. The limits of the integration are fixed by the asymptotics of the scalar field ϕ . Note that the fermions at $x = 0$ are identified with the spectral fermions of the Toda (or, more generally Nahm) system.

⁵⁸In monopole terms, the variable w is the coordinate on the twistor manifold $\mathbb{CP}^1$.

One may also consider the generalization of the Nahm construction for the $SU(n)$ case [206]. Then, one needs to fix asymptotics of the n components of the scalar field to be some (ordered) numbers μ_i , $i = 1, \dots, n$, $\sum_i \mu_i = 0$. It naturally divides the domain where the Nahm system is given onto the $n - 1$ intervals (μ_1, μ_2) , $\dots$, (μ_{n-1}, μ_n) . On each a -th interval the Nahm system for the three $p_k \times p_k$ matrices $\mathcal{T}_{i,a}(t)$ is

$$\frac{d\mathcal{T}_{i,a}}{dt} = 2\epsilon_{ijk} [\mathcal{T}_{j,a}, \mathcal{T}_{k,a}] \quad (8.16)$$

Thus, the $SU(n)$ monopole is characterized by the set of topological charges $p_1, \dots, p_{n-1}$.

The correspondence between solutions to the Nahm equations and the spectral curves of integrable systems is generalized to the $SU(N_c)$ case [206], where the connection with the higher spin magnets emerges [38]. The solution to the Nahm equations is a product of $n - 1$ components, each of them being a spectral curve for the $SU(2)$ monopole of charge p_a , while the whole monopole spectral curve is a special case of the curve for the $sl(p)$ spin magnet with $p = \sum_a p_a$. The configuration of the magnet is specified by a peculiar pair clusterization, each of them describing the $sl(p_a)$ magnet. There are also additional matching conditions for the correlations between the two-site groups.

The peculiarity of the spin chain configuration has a clear explanation in the brane language. The monopole moduli space is described via the configuration of D1- and D3-branes [206]. The clusterization in the brane language means that the D1-branes have their endpoints only on the neighbor D3-branes. Thus, appealing to the spin chain interpretation, one can identify the D3-branes with the sites of the spin chain, while the D1-branes give rise to the $SU(p)$ group, p being the full number of the D1-branes. Let us note that the twisting procedure used in section 3 is equivalent to adding semi-infinite D1-branes to the very left and the very right D3-branes.

8.5 Branes and two Lax representations

Now let us consider two distinct Lax representations of the Toda chain that we discussed in section 3.2 and explain their interpretation in terms of brane diagrams [38]. Remind that in the 2×2 representation the system is formulated in terms of the transfer matrix, while for the $N_c \times N_c$ representation the notion of the transfer matrix is missed.

From the brane pictures point of view the two Lax representations are nothing but the choice of parametrization of the fivebrane worldvolume Σ . Of two holomorphic coordinates (which are Cartan elements of scalars) $\lambda = x_4 + ix_5$ and $s = x_6 + ix_{10}$,⁵⁹ we can regard any we wish as a spectral parameter. In the “standard” 2×2 representation, $w = \exp \frac{-s}{R}$ is a spectral parameter. Another possibility is to choose λ as a spectral parameter. It leads to the $N_c \times N_c$ representation.

Geometrical image of this phenomenon is the base-fiber symmetry of the Calabi-Yau threefold [54]. In the “standard” 2×2 language w parameterizes the base A_m singularity and counts the number m of gauge factors. λ is a coordinate in the fiber. If each coefficient at w^i is a polynomial of degree n , we deal with the $SU(n)^m$ gauge theory with n extra fundamentals at each end of the chain of the gauge factors. The duality $SU(n)^m \leftrightarrow SU(m+1)^{n-1}$ proposed in [54] is nothing but the $m+1 \times m+1 \leftrightarrow n \times n$ symmetry of the Lax representations. Under this symmetry the role of coordinates on the Newton polygon describing the manifold is changed.

Note that the generating differential

⁵⁹Equivalently, $s = x_6 + ix_9$ in the Type IIB language.

$$dS = \lambda \frac{dw}{w} = \lambda ds \approx -sd\lambda \quad (8.17)$$

also suggests this symmetry.

Let us extend this argumentation onto description of the fundamental matter in $4d$ gauge theory. There are two different realizations of the matter in the brane language: via semi-infinite D4-branes and via D6-branes. On the integrable side, the first possibility corresponds to the additionally twisted Toda chain [209], while the second one to the **XXX** magnet.

Indeed, the curve for the theory with the fundamental matter can be also obtained from the $N_c \times N_c$ representation by deforming the Lax operator. To this end, one has to add non-trivial entries to the first and the last columns which corresponds to attaching the semi-infinite branes to the end branes of the configuration. On the contrary, in 2×2 representation D6-branes correspond to the fundamentals which are localized at the points corresponding to the masses. It is in agreement with the relation between the masses and inhomogeneities in the spin chain. The both representations can be related if one takes into account the phenomenon of the brane creation when branes pass through each other [47].

In [47] the authors considered the three dimensional theory in the common longitudinal directions of branes (in the type IIB theory):

$$\left\{ \begin{array}{l} NS5 \\ D3 \\ D5 \end{array} \right| \begin{array}{cccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ + & + & + & + & + & + & - & - & - & - \\ + & + & + & - & - & - & + & - & - & - \\ + & + & + & - & - & - & - & + & + & + \end{array} \quad (8.18)$$

By T-duality in the third direction, this setup gets mapped onto the Type IIA brane picture [51] so that the statement of creating a new D3-brane when the D5-brane passes through the NS5-brane transforms into the claim that, moving the D6-brane through the NS5-brane creates a new D4-brane stretched between them.

8.6 Whitham dynamics via branes

So far we discussed the brane interpretation of the Toda/spin chains. Now we are going to say some words about the other, Whitham integrability of Seiberg-Witten theory and start with the brane configuration in F-theory, i.e. on the elliptically fibered K3 manifold or, equivalently, on orientifold in IIB theory.

According to [210], $\mathcal{N} = 2$ theory in $4d$ can be considered as a theory on the worldvolume of 3-branes in the background of four 7-branes located at the points $\pm\Lambda$ on the complex plane of $u = \text{Tr}\phi^2$ for the $SU(2)$ gauge group. We assume that masses of all fundamentals are large and, therefore, we can work with the pure gauge theory.

Consider now the dynamics of the 3-branes along the directions transverse to the background 7-branes. We shall restrict ourselves to the $SU(2)$ gauge group. However, since the whole picture is very transparent, the generalization to higher rank groups is straightforward.

We are going to study the Whitham dynamics of moduli parametrized by branching points. In the $SU(2)$ case, one can write the Seiberg-Witten curve in the hyperelliptic form (3.8)

$$Y^2 = (\lambda^2 - u)^2 - \Lambda_{QCD}^4, \quad dS = 2\lambda^2 \frac{d\lambda}{Y} \quad (8.19)$$

i.e. there are 4 branching points $\pm\sqrt{u \pm \Lambda_{QCD}^2}$. Since the Whitham dynamics governs the u -behaviour, all the four points move under the Whitham flows. Therefore, it is more convenient to use another parameterization that goes back to the original paper [1]

$$Y^2 = (x + u)(x - \Lambda_{QCD}^2)(x + \Lambda_{QCD}^2), \quad dS = (x + u) \frac{dx}{Y} \quad (8.20)$$

This parameterization is obtained from (8.19) by change of variables $x = \lambda^2 - u$. This is a different curve but the new differential dS has on this curve the same periods as the differential (8.19).

The parameterization (8.20) has the evident advantage that only one of its branching points is moved by the Whitham flows. This Whitham dynamics is nothing but the Gurevich-Pitaevskii solution first emerged in the problem of modulation of the one-zone solution of the KdV equation [211]. In the brane language this solution looks as follows.

Initially, the 3-branes coincide with a pair of 7-branes and then they scatter during the evolution process. After all, they are absorbed by the remaining pair of 7-branes. Therefore, the exact metric on the moduli space is provided by the 3-brane exchange process which generalizes the closed string exchange picture suggested in [212].

Returning back to the Whitham dynamics, the coordinate u defines positions of the two 3-branes located at the points $\pm\sqrt{u}$ on the a -plane⁶⁰, while the other branching points fix locations of the background branes. The branching point corresponding to the dynamical 3-branes moves in accordance with the Whitham flows for the one-zone solution from $u = \Lambda_{QCD}^2$ to $u = -\Lambda_{QCD}^2$, while 7 branes remain at rest.

Let us emphasize the close analogy between the dynamics of branes and the low-energy monopole scattering. Indeed, there is a geodesics which corresponds to the transformation of the monopoles into dyons [213]. The brane evolution above is equivalent to this process in the following sense. At the initial moment, when branes are located at the points $\pm\Lambda_{QCD}^2$, there are massless monopoles in the theory in accordance with the exact formula (1.5) for the BPS spectrum. After the scattering process at the right angle at $u = 0$, the branes acquire an electric charge which can be explained as follows. At the endpoint of the evolution when branes are at $u = -\Lambda_{QCD}^2$, there are massless dyons in the theory. However, the dyons can be treated as (1,1) strings stretched between the dynamical and background branes [214]. Therefore, the dynamical branes have to acquire the electric charge necessarily.

9 Conclusion

We have tried to clarify the role and interpretation of classical integrability in the vacuum sector of supersymmetric gauge theories which we basically understand as obtained via string compactifications. Briefly this can be formulated as follows. The vacuum sector is described in terms of moduli spaces which describe the phase spaces of two different integrable systems. An integrable many-body system defines a dynamics on a moduli space which can be formulated in terms of the geometry of the compactification manifold. On the other hand, the second integrable dynamics, the Whitham dynamics can be interpreted as governing the renormalization group flows in the theory and develops

⁶⁰Since the whole our consideration here is done at the quasi-classical (perturbative) level, $u = a^2$.

on the Coulomb branch of the moduli space. Both integrable systems can be interpreted in terms of the dynamics of collective coordinates of branes in different dimensions.

The effectiveness and the validity of the approach is out of doubts at the moment, despite the lack of its derivation from the first principles. The most important problem to be solved seems to be to sum up the infinite number of instantons into the finite number of degrees of freedom that provide the phase space for the integrable system. One more question essential from the conceptual point of view is to clarify the meaning of quantization of the integrable system. Role of the corresponding “Planck constant” is uncertain at the moment.

It is absolutely clear that the branes are extremely important for dealing with nonperturbative contributions and play the role analogous to the Feynman diagrams in the perturbation theory. We have shown that, in simplest situations, integrability can be described in brane terms, however, the extensive formulation of these relations is still missed and, therefore, further work in this direction is desired.

The authors are grateful to H.W.Braden, V.Fock, S.Gukov, A.Marshakov, A.Morozov, I.Krichever, N.Nekrasov and B.Rubtsov for discussions and collaboration and to A.Gerasimov, S.Kharchev, A.Losev, M.Olshanetsky, A.Rosly, K.Selivanov and A.Zabrodin for useful discussions. A.G. thanks University of Angers where the work was partially done for the hospitality. A.M. thanks T.Takebe for discussions on duality and kind hospitality at the Ochanomizu University, Tokyo, where this work was partially done. The work was supported in part by grants INTAS-99-1705, RFBR-98-01-00327, CRDF-RP1-2108 and CNRS grant for research 2000 (A.G.) and by grants INTAS 99-0590, CRDF #6531, RFBR 00-02-16101a and the JSPS fellowship for research in Japan (A.M.).

References

- [1] N.Seiberg and E.Witten, Nucl.Phys., **B426** (1994) 19-52
- [2] N.Seiberg and E.Witten, Nucl.Phys., **B431** (1994) 484-550
- [3] B.Dubrovin, I.Krichever and S.Novikov, “Integrable systems - I”, *Sovremennye problemy matematiki (VINITI), Dynamical systems - 4*, (1985) 179
B.Dubrovin, UMN, **36** (1981) 11
- [4] I.Krichever, Uspekhi Mat.Nauk, **44:2** (1989) 121
- [5] B.Dubrovin,S.Novikov, Math.Phys.Rev., **9** (1992) 105-213
I.Krichever, Math.Phys.Rev., **9** (1992) 3-105
- [6] A.Bilal, hep-th/96010010
W.Lerche, Nucl.Phys.Proc.Suppl., 55B (1997) 83-117; Fortsch.Phys., 45 (1997) 293-340
L.Alvarez-Gaume and S.F.Hassan, Fortsch.Phys., 45 (1997) 159-236
- [7] A. Giveon and D. Kutasov, Rev.Mod.Phys., 71 (1999) 983-1084
J.Schwarz, hep-th/9601077
W.Taylor, hep-th/9801182
- [8] A.Klemm, hep-th/9705131
- [9] A.Morozov, UFN, **162** (1992) 84; UFN, **164** (1994) 3 -62
- [10] H.W.Braden and I.M.Krichever (Eds.), *sl Integrability: The Seiberg-Witten and Whitham Equations*, Gordon and Beach, 2000.
- [11] E.D’Hoker and D.H.Phong, hep-th/9912271
- [12] M.Shifman, Phys.Rep., **209** (1991) 343-380
- [13] A.Migdal and M.Shifman, Phys.Lett., **B114** (1982) 445
G.Veneziano and S.Yankielowicz, Phys.Lett., **B113** (1982) 231-236
- [14] A.Mironov, Theor.Math.Phys., **114** (1998) 127
- [15] C.Vafa and S.Cecotti, Comm.Math.Phys., **158** (1993) 569-644
- [16] M.Douglas, Phys.Lett., **B238** (1990) 176
- [17] M.Fukuma, H.Kawai and R.Nakayama, Comm.Math.Phys., **143** (1992) 371-403
R.Dijgraaf, E.Verlinde and H.Verlinde, Nucl.Phys., **B352** (1991) 59-86
- [18] A.Gerasimov, A.Marshakov, A.Mironov, A.Morozov and A.Orlov, Nucl.Phys., **B357** (1991) 565
- [19] S.Kharchev, A.Marshakov, A.Mironov, A.Morozov and A.Zabrodin, Nucl.Phys., **B380** (1992) 181
- [20] S.Kharchev, A.Marshakov, A.Mironov and A.Morozov, Nucl.Phys., **B397** (1993) 339-378
- [21] F.David, Mod.Phys.Lett., **A5** (1990) 1019

- [22] A.Mironov and A.Morozov, Phys.Lett., **B252** (1990) 47-52
J.Ambjorn, J.Jurkiewicz and Yu.Makeenko, Phys.Lett., **251B** (1990) 517-524
H.Ityama and Y.Matsuo, Phys.Lett., **255B** (1991) 202
Y.Makeenko, A.Marshakov, A.Mironov and A.Morozov, Nucl.Phys., **B356** (1991) 574
- [23] M.Atiyah, V.Drinfeld, N.Hitchin and Yu.Manin, Phys.Lett., **A65** (1978)
- [24] M.Douglas, J.Geom.Phys., 28 (1998) 255-262
- [25] E.Witten, Nucl.Phys., **B340** (1990) 281-332; Surveys Dif.Geom., **1** (1991) 243-310
- [26] M.Kontsevich, Func.Analiz i Prilozh., **25** (1990) 50-64
- [27] E.Witten, Comm.Math.Phys., **117** (1988) 353
- [28] S.Donaldson, Duke.Math.J., **54** (1987) 231
- [29] K.Li, Nucl.Phys., **B354** (1991) 711
- [30] V.Novikov, M.Shifman, A.Vainshtein and V.Zakharov, Nucl.Phys., **B229** (1983) 381
- [31] V.Novikov, M.Shifman, A.Vainshtein and V.Zakharov, Nucl.Phys., **B229** (1983) 407
- [32] A.Marshakov, A.Mironov and A.Morozov, Phys.Lett., **B389** (1996) 43
- [33] M.Shifman, A.Vainshtein, Nucl.Phys., **B359** (1991) 571-580
N.Seiberg, hep-th/9408013
- [34] C.Montonen and K.Olive, Phys.Lett., **B78** (1977) 117
- [35] K.Olive and E.Witten, Phys.Lett., **B78** (1979) 97
- [36] A.Klemm, W.Lerche, S.Theisen and S.Yankielowicz, Phys.Lett., **344B** (1995) 169
P.Argyres and A.Faraggi, Phys.Rev.Lett., **73** (1995) 3931
- [37] A.Gorsky, I.Krichever, A.Marshakov, A.Mironov and A.Morozov, Phys.Lett., **B355** (1995) 466-477
- [38] A.Gorsky, S.Gukov and A.Mironov, Nucl.Phys., **B517** (1998) 409-461
- [39] A.Marshakov, A.Mironov and A.Morozov, Int.J.Mod.Phys., **A15** (2000) 1157-1206
- [40] I.Krichever, Comm.Pure Appl.Math., **47** (1994) 437;
B.Dubrovin, Nucl.Phys., **B379** (1992) 627; hep-th/9407018
- [41] E.Witten, Surv.Diff.Geom., **1** (1991) 243
R.Dijkgraaf, E.Verlinde and H.Verlinde, Nucl.Phys., **B352** (1991) 59
- [42] Yu.Manin, *Frobenius manifolds, quantum cohomology and moduli spaces*, MPI, 1996
M.Kontsevich and Yu.Manin, Comm.Math.Phys., **164** (1994) 525
- [43] J.Polchinski, Phys.Rev.Lett., **75** (1995) 4724-4727; hep-th/9611050
- [44] C.Hall and P.Townsend, Nucl.Phys., **B438** (1995) 109-137
P.Townsend, hep-th/9507048
- [45] E.Witten, Nucl.Phys., **B430** (1996) 335-350

- [46] S.Kachru and C.Vafa, Nucl.Phys., **B450** (1995) 69-89
- [47] A.Hanany and E.Witten, Nucl.Phys., **B492** (1997) 152
- [48] S.Katz, A.Klemm and C.Vafa, Nucl.Phys., **B497** (1997) 173-195
- [49] H.Ooguri and C.Vafa, Nucl.Phys., **B463** (1996) 55-72
- [50] A.Klemm, W.Lerche, P.Mayr, C.Vafa and N.Warner, Nucl.Phys. **B477** (1996) 746-766
- [51] E.Witten, Nucl.Phys., **B500** (1997) 3-42
- [52] P.Townsend, Phys.Lett., **B373** (1996) 68-75; Phys.Lett., **B350** (1995) 184-187
- [53] K.Hori, H.Ooguri and Y.Oz, Adv.Theor.Math.Phys., **1** (1998) 1
E.Witten, Nucl.Phys., **B507** (1997) 658
- [54] S.Katz, P.Mayr and C.Vafa, Adv.Theor.Math.Phys., **1** (1998) 53-114
- [55] C.Vafa and H.Ooguri, Nucl.Phys., **B500** (1997) 62-74
- [56] M.Bershadsky, V.Sadov and C.Vafa, Nucl.Phys., **B463** (1996) 420;
- [57] C.Vafa, Nucl.Phys., **B469** (1996) 403-417
- [58] F.Calogero, J.Math.Phys., **12** (1971) 419
J.Moser, Adv.Math., **16** (1975) 197-220
- [59] B.Sutherland, Phys.Rev., **A5** (1972) 1372
- [60] M.Olshanetsky and A.Perelomov, Phys.Peps., **71** (1981) 313; Phys.Rep., **94** (1983) 6
D.Kazhdan, B.Kostant and S.Sternberg, Comm. on Pure and Appl.Math., Vol.XXXI (1978) 481-507
- [61] S.N.M.Ruijsenaars and H.Schneider, Ann.Phys. (NY), **170** (1986) 370
S.N.M.Ruijsenaars, Comm.Math.Phys., **110** (1987) 191-213
- [62] S.N.Ruijsenaars, Comm.Math.Phys., **115** (1988) 127-165
- [63] A.Gorsky and N.Nekrasov, Nucl.Phys., **B414** (1994) 213-231
- [64] J.A.Minahan and A.P.Polychronakos, Phys.Lett., **B326** (1994) 288;
A.Kovner, E.Langmann and M.Salmhofer, Mod.Phys.Lett., **A9** (1994) 2913-2926;
J.Blom and E.Langmann, Nucl.Phys., **B563** (1999) 506-532
- [65] A.Gorsky and N.Nekrasov, Nucl.Phys., **B436** (1995) 582-608
- [66] O.Babelon and D.Bernard, Phys.Lett., **B317** (1993) 363
- [67] N.Nekrasov, Nucl.Phys., **B531** (1998) 323-344
- [68] N.Nekrasov and A.Lowrence, Nucl.Phys., **B513** (1998) 239-265
- [69] H.W.Braden, A.Marshakov, A.Mironov and A.Morozov, Nucl.Phys., **B558** (1999) 371
- [70] A.Gorsky and N.Nekrasov, hep-th/9401021
- [71] J.Gibbons and T.Hermesen, Physica, **D11** (1984) 337

- [72] I.Krichever, O.Babelon, E.Billey and M.Talon, hep-th/9411160
- [73] I.Krichever, Func.Anal. & Appl., **14** (1980) 282
- [74] G.Felder and A.Varchenko, q-alg/9605024
- [75] P.I.Etingof, J.Math.Phys., **36** (1995) 2636
P.I.Etingof and A.A.Kirillov Jr., hep-th/9403168
P.I.Etingof, I.B.Frenkel and A.A.Kirillov, hep-th/9407047
- [76] N.Hitchin, Duke.Math.J., **54** (1987) 91
- [77] E.Markman, Comp.Math., **93** (1994) 255
- [78] N.Nekrasov, Comm.Math.Phys., **180** (1996) 587-604
- [79] B.Enriquez and V.Rubtsov, Math.Phys.Lett., **3** (1996) 343
- [80] G.E.Arutyunov, S.A.Frolov and P.B.Medvedev, J.Math.Phys., **38** (1997) 5682; J.Phys., **A30** (1997) 5051
- [81] J. Hurtusise and E. Markman, math/9919161
- [82] A.Gorsky, A.Mironov, A.Marshakov and A.Morozov, Nucl.Phys., **B527** (1998) 690-716
- [83] K.Landsteiner, J.M.Pierre and S.B.Giddings, Phys.Rev., **D55** (1997) 2367
- [84] A.Yung, Nucl.Phys., **B485** (1997) 38-62
V.V.Khoze, M.P.Mattis and M.J.Slater, Nucl.Phys., **B536** (1998) 69-109
N.Dorey, V.V.Khoze and M.P.Mattis, Phys.Lett., **B390** (1997) 205
N.Dorey, V.V.Khoze and M.P.Mattis, Phys.Lett., **B388** (1996) 324
- [85] K.Landsteiner, E.Lopez and D.A.Lowe, JHEP, **9807** (1998) 011
- [86] I.Krichever and D.H.Phong, hep-th/9912180
- [87] R.Donagi and E.Witten, Nucl.Phys., **B460** (1996) 299-334
- [88] E.Martinec, Phys.Lett., **B367** (1996) 91-96
- [89] A.Gorsky, A.Marshakov, Phys.Lett., **B374** (1996) 218-224.
- [90] H.Itoyama and A.Morozov, Nucl.Phys., **B477** (1996) 855-877
- [91] H.Itoyama and A.Morozov, Nucl.Phys., **B491** (1997) 529-573
- [92] H.W.Braden, A.Marshakov, A.Mironov and A.Morozov, Nucl.Phys., **B573** (2000) 553
- [93] A.Mironov and A.Morozov, Phys.Lett., **B475** (2000) 71
- [94] A.Mironov and A.Morozov, hep-th/0001168
- [95] A.Hanany and Y.Oz, Nucl.Phys., **B452** (1995) 283
P.Argyres, D.Plesser and A.Shapere, Phys.Rev.Lett., **75** (1995) 283
J.Minahan and D.Nemeschansky, Nucl.Phys., **B464** (1996) 3-17
- [96] A.Gorsky, A.Marshakov, A.Mironov and A.Morozov, Phys.Lett., **B380** (1996) 75-80

- [97] A.Gorsky, A.Marshakov, A.Mironov and A.Morozov, hep-th/9604078
- [98] A.Gorsky, S.Gukov and A.Mironov, Nucl.Phys., **B518** (1998) 689
- [99] A.Marshakov, A.Mironov, Nucl.Phys., **B518** (1998) 59-91
- [100] V.Inozemtsev, Comm.Math.Phys., **121** (1989) 629
- [101] V.E.Adler and A.B.Shabat, Theor.Math.Phys., **111** (1997) 323; **112** (1997) 179
I.Krichever, hep-th/9909224
- [102] E.Martinec and N.Warner, Nucl.Phys., **459** (1996) 97
- [103] T.Nakatsu and K.Takasaki, Mod.Phys.Lett., **A11** (1996) 157-168
T.Eguchi and S.Yang, Mod.Phys.Lett., **A11** (1996) 131-138
- [104] I.Krichever and D.Phong, hep-th/9708170 and references therein
- [105] M.Toda, *Hi-senkei koushi rikigaku*, Iwanami Shoten, Tokyo, 1978, 1987; *Theory of nonlinear lattices*, Springer-Verlag, Berlin-New York, 1981
- [106] O.Bogoyavlensky, Comm.Math.Phys., **51** (1976) 201-209
- [107] E.Sklyanin, J.Phys., **A21** (1988) 2375
V.Kuznetsov, J.Math.Phys., **31** (1990) 1167
V.Kuznetsov and A.Tsiganov, J.Sov.Math., **59** (1992) 1085-1092
V.Kuznetsov, M.Jorgenson and P.Christiansen, J.Phys., **A28** (1995) 4639
- [108] A.Gorsky and A.Mironov, Nucl.Phys., **B550** (1999) 513
- [109] H.Bateman and A.Erdelyi, *Higher transcendental functions*, vol.1, London 1953
E.T. Whittaker and G.N Watson, *A course in modern analysis*, Cambridge University Press, 4th ed. 1988
- [110] E.D'Hoker and D.H.Phong, hep-th/9808156
- [111] E.D'Hoker and D.H.Phong, Nucl.Phys., **B530** (1998) 537-610; 611-640
A.J.Bodner, E.Corrigan and R.Sasaki, Prog.Theor.Phys., **100** (1998) 1107-1129; **102** (1999) 499-529
A.J.Bodner, R.Sasaki and K.Takasaki, Prog.Theor.Phys., **101** (1999) 487-518
A.J.Bodner and R.Sasaki, Prog.Theor.Phys., **101** (1999) 799-829
- [112] E.D'Hoker and D.H.Phong, Nucl.Phys., **B534** (1998) 697-719
- [113] L.Faddeev and L.Takhtadjan, *Hamiltonian Approach to the Theory of Solitons*, 1986
- [114] P.C.Argyres and A.D.Shapere, Nucl.Phys., **B461** (1996) 437
- [115] S.Kharchev, "Twisted systems of the **XXZ** type", preprint ITEP, 1996; unpublished
- [116] A.Gorsky, Mod.Phys.Lett., **A12** (1997) 719
- [117] A.Hanany, Nucl.Phys., **466** 85
U.Danielsson and B.Sundborg, Phys.Lett., **B358** (1995) 273
A.Brandhuber and K.Landsteiner, Phys.Lett., **B358** (1995) 73

- [118] A.Brandhuber, N.Itzhaki, J.Sonnenschain, S.Theisen and S.Yankielowicz, Phys.Lett., **B415** (1997) 127-134
- [119] A.Kundu, hep-th/9403001
- [120] O.Aharony, A.Hanany and B.Kol, JHEP, **9801** (1998) 002
- [121] S.N.Ruijsenaars, Comm.Math.Phys., **133** (1990) 217-247
See a review in:
S.Kharchev, A.Mironov and A.Zhedanov, Int.J.Mod.Phys., **A12** (1997) 2675-2724
- [122] H.W. Braden, A.Marshakov, A.Mironov and A.Morozov, Phys.Lett., **B448** (1999) 195
- [123] S.N.M.Ruijsenaars, in: *Proceedings of the 1994 Banff Summer School Particles and Fields*, G. Semenoff and L. Vinet Eds, to appear
S.N.M.Ruijsenaars, In: *Integrable and superintegrable systems*, ed. B.Kupershmidt, World Scientific, 165-206, 1990
S.N.M.Ruijsenaars, Publ. RIMS, Kyoto Univ. 30 (1994), 865
H.W. Braden and R. Sasaki, Progr.Theor.Phys., **97** (1997) 1003-1018 and references therein
- [124] S.N.M.Ruijsenaars, Publ. RIMS, Kyoto Univ. 31 (1995), 247
- [125] M.Bruschi and F.Calogero, Commun.Math.Phys., **109** (1987) 481-492
- [126] I.Krichever and A.Zabrodin, Russ.Math. Surveys, **59** (1995) 1101
- [127] V.M.Buchstaber and I.M.Krichever, Teor.Mat.Fiz., **94** (1993) 200-212
H.W.Braden and V.M.Buchstaber, Reviews in Mathematics and Mathematical Physics, **10** No.2 (1997)
H.W.Braden and V.M.Buchstaber, SIAM J.Math.Anal., **28** (1997) 903
H.W.Braden and J.G.B.Byatt-Smith, Bull.Lond.Math.Soc., **31** (1999) 463-470
- [128] E.K.Sklyanin, Prog.Theor.Phys.Suppl., **118** (1995), 35-60
V.B.Kuznetsov and E.K.Sklyanin, J.Phys. **A29** (1996) 2779
V.B.Kuznetsov, F.W.Nijhoff and E.K.Sklyanin, Commun.Math.Phys., **189** (1997) 855
- [129] S.N.M.Ruijsenaars, Annals of Physics, **256** (1997) 226
H.W.Braden and Andrew N.W.Hone, Phys.Lett., **B380** (1996) 296
- [130] E.Date, M.Jimbo, T.Miwa and M.Kashiwara, In: *Nonlinear Integrable Systems - Classical Theory and Quantum Theory*, Ed. by M. Jimbo, T. Miwa, World Scientific, Singapore, 1983
M.Jimbo and T.Miwa, Publ. RIMS, Kyoto Univ., **19** (1983) 943-1001
- [131] Kai Chen, Bo-yu Hou and Wen-Li Yang, J.Math.Phys., **41** (2000) 8132-8147; hep-th/0011145
- [132] E.Sklyanin, Func.Anal. & Apps., **16** (1982) 27
E.Sklyanin, Func.Anal. & Apps., **17** (1983) 34
- [133] M.Shifman and A.Vainshtein, In *M.A.Shifman: *ITEP lectures on particle physics and field theory*,* vol.2, World Scientific, Singapore, 1999 [hep-th/9902018].
K.Intriligator and N.Seiberg, hep-th/9506084
- [134] G.Dvali and M.Shifman, Phys.Lett., **B396** (1997) 64

- [135] G.W.Gibbons and P.K.Townsend, Phys.Rev.Lett., **83** (1999) 1727
S.M.Carroll, S.Hellerman and M.Trodden, Phys.Rev., **D61** (2000) 065001
A.Gorsky and M.Shifman, Phys.Rev., **D61** (2000) 085001
- [136] V.A.Novikov, M.A.Shifman, A.I.Vainshtein and V.I.Zakharov, Nucl.Phys., **260B** (1985) 157
- [137] S.Elitzur, A.Forger, A.Giveon and E.Rabinovici, Phys.Lett., **B353** (1995) 79; Nucl.Phys., **B459** (1996) 160
S.Elitzur, A.Forger, A. Giveon, K.Intrilligator and E.Rabinovici, Phys.Lett., **B379** (1996) 121
S.Terashima and S.Yang, Phys.Lett., **B391** (1997) 107
T.Kitao, Phys. Lett., **B402** (1997) 290
T.Kitao, S.Terashima, and S.Yang, Phys.Lett., **B399** (1997) 75
K.Konishi and H.Terao, Nucl.Phys., **B511** (1998) 264
- [138] A.Giveon, hep-th/9611152
- [139] A.Gorsky, A.Vainshtein and A.Yung, Nucl.Phys., **B584** (2000) 197
- [140] M.R.Douglas and S.H.Shenker, Nucl.Phys., **B447** (1995) 271
- [141] A.Hanany, M.Strassler and A.Zaffaroni, Nucl.Phys., **B513** (1998) 87
- [142] P.C.Argyres and M.R.Douglas, Nucl.Phys., **B448** (1995) 93
P.C.Argyres, M.R.Plesser, N.Seiberg and E.Witten, Nucl.Phys., **B461** (1996) 71
- [143] T.Eguchi, K.Hori, K.Ito and S.Yang, Nucl.Phys., **B471** (1996) 430
- [144] N.Dorey, JHEP, **9907** (1999) 021
- [145] N.M.Davies, T.J.Hollowood, V.V.Khoze and M.P.Mattis, Nucl.Phys., **B559** (1999) 123
- [146] S.Katz and C.Vafa, Nucl.Phys., **B497** (1997) 196
- [147] N.Seiberg and E.Witten, hep-th/9607163
- [148] H.Braden and A.Marshakov, hep-th/0009060
- [149] J.D.Edlestein, M.Mariño and J.Mas, Nucl.Phys., **B541** (1999) 671-697
- [150] E.D'Hoker, I.M.Krichever and D.H.Phong, Nucl.Phys., **B489** (1997) 179-210; Nucl.Phys., **B489** (1997) 211-222
- [151] A.Beilinson and A.Levin, *Proceedings of Symposia in Pure Mathematics*, Vol.55, Part 2 (1994) 126-196
V.Kuznetsov, F.Nijhoff and E.Sklyanin, Comm.Math.Phys., **189** (1997) 855-877
J.Harvey and G.Moore, Nucl.Phys., **B463** (1996) 315-368
- [152] N.Seiberg, Five-dimensional SUSY field theories, nontrivial fixed points, and string dynamics, Phys.Lett. **B388** (1996) 753
K.Intriligator, D.R.Morrison and N.Seiberg, Five-dimensional supersymmetric gauge theories and degenerations of Calabi-Yau Spaces, Nucl.Phys. **B497** (1997) 56
N.Seiberg and D.Morrison, Nucl.Phys., **B483** (1997) 229
- [153] K.L.Vaninsky, solv-int/9808018

- [154] E.D'Hoker and D.Phong, Nucl.Phys., **513** (1998) 405-444
- [155] I.Krichever, hep-th/9205110; Comm.Math.Phys., **143** (1992) 415
- [156] G.Moore and E.Witten, hep-th/9709193
K.Takasaki, Int.J.Mod.Phys., **A14** (1999) 1001-1014
- [157] A.Losev, N.Nekrasov, S.Shatashvili, Nucl.Phys., **B534** (1998) 549-611; hep-th/9801061
- [158] M.Mariño and G.Moore, Comm.Math.Phys., **199** (1998) 25-69
- [159] H.Flashka, G.Forest and D.W.McLaughlin, Comm.Pure Appl.Math., **33** (1979) 299-329
- [160] M.Mineev-Weinstein, P.B.Wiegmann and A.Zabrodin, Phys.Rev.Lett., **84** (2000) 5106
- [161] A.S.Gorsky, hep-th/0010068
- [162] M.Matone, Phys.Lett., **B357** (1995) 342; Phys.Rev., **D53** (1996) 7354
J.Sonnenschein, S.Theisen and S.Yankielowicz, Phys.Lett., **B367** (1996) 145-150
T.Eguchi and S.K.Yang, Mod.Phys.Lett., **A11** (1996) 131
P.Howe and P.West, Nucl.Phys., **B486** (1997) 425
E.D'Hoker, I.Krichever and D.H.Phong, Nucl.Phys., **B494** (1997) 89
- [163] S.Kharchev, A.Marshakov, A.Mironov and A.Morozov, Mod.Phys.Lett., **A8** (1993) 1047-1061
S.Kharchev, hep-th/9810091
- [164] J.D.Edlestein, M.Gomez-Reino and J.Mas, Nucl.Phys., **B561** (1999) 273-292
J.D.Edlestein, M.Gomez-Reino, M.Mariño and J.Mas, Nucl.Phys., **B574** (2000) 587-619
- [165] K.Takasaki, Int.J.Mod.Phys., **A15** (2000) 3635-3666; Prog.Theor.Phys.Suppl., **135** (1999) 53-74
- [166] J.D.Edlestein, M.Gomez-Reino and M.Mariño, hep-th/0006113
- [167] H.Braden, A.Mironov and M.Olshanetsky, unpublished
- [168] A.Marshakov, A.Mironov and A.Morozov, Mod.Phys.Lett., **A12** (1997) 773
- [169] K.Ito and S.-K.Yang, Phys.Lett., **B433** (1998) 56-62
- [170] A.Veselov, Phys.Lett., **A261** (1999) 297-302
- [171] G.Bertoldi and M.Matone, Phys.Rev., **D57** (1998) 6483-6485
- [172] J.M.Isidro, Nucl.Phys., **B539** (1999) 379-402
- [173] A.Morozov, Phys.Lett., **B427** (1998) 93-96
- [174] A.Losev, JETP Lett., **65** (1997) 374
- [175] A.Mironov and A.Morozov, Phys.Lett., **B424** (1998) 48-52
- [176] L.Caporaso, J.Harris, alg-geom/9608025
E.Getzler, alg-geom/9612004
for the latest review and further references, see:
E.Getzler, math/9812026
- [177] A.Gorsky, N.Nekrasov and V.Roubtsov, hep-th/9901089

- [178] V.Fock, A.Gorsky, N.Nekrasov and V.Roubtsov, JHEP, **0007** (2000) 028
- [179] E. Sklyanin, solv-int/9505003
- [180] B.Enriquez, B.Feigin and V.Rubtsov, Compositio Math., **110** (1998) 1
- [181] K.Takasaki, math/0007073
- [182] J.Hurtubise, Duke.Math.Journal, **83** (1996) 19-50
- [183] G.Wilson, Inv.Math., **133** (1998) 1-41
- [184] H.Nakajima, Lectures on Hilbert Schemes of Points on Surfaces, 1996
- [185] E.Frenkel, q-alg/9506003
- [186] K.Intrilligator and N.Seiberg, Phys.Lett., **B387** (1996) 513
- [187] A.Perelomov, *Integrable Systems of Classical Mechanics and Lie Algebras*, Vol.I, Birkhauser, 1990
L.Gavrilov and A.Perelomov, solv-int/9905011
- [188] to appear
- [189] H.Airault, H.McKean and J.Moser, Comm.Pure and Applied Math., **30** (1977) 95
T.Shiota, J.Math.Phys., **35** (1994) 5844-5849
- [190] I.Krichever and A.Zabrodin, Uspekhi Mat.Nauk, **50** (1995) 3-56
- [191] S.Kharchev, unpublished
- [192] V.Fock and A.Rosly, math/9802054
- [193] See, e.g., A.Klemm, W.Lerche and S.Theisen, Int.J.Mod.Phys., **A11** (1996) 1929
K.Ito and N.Sasakura, Nucl.Phys., **B484** (1997) 141
- [194] J.D.Edelstein and J.Mas, hep-th/9902161
- [195] J.A.Minahan, D.Nemeschansky and N.P.Warner, Nucl.Phys., **B528** (1998) 109
- [196] P.Etingof and A.Varchenko, math/9907181
G.Felder, Y.Markov, V.Tarasov and A.Varchenko, math/0001184
V.Tarasov and A.Varchenko, math/0002132
- [197] A.Gorsky, Phys.Lett., **B410** (1997) 22-27
- [198] O.Aharony and A.Hanany, Nucl.Phys., **B504** (1997) 239-271
- [199] J.Barbon and A.Pasquinucci, Nucl.Phys., **B517** (1998) 125; Fortsch.Phys., **47** (1999) 255-262
- [200] A.Kapustin and S.Sethi, Adv.Theor.Math.Phys., **2** (1998) 571 A.Kapustin, Nucl.Phys., **B534** (1998) 531
- [201] O.J.Ganor, A.Yu.Mikhailov and N.Saulina, hep-th/0007236
Y.E.Cheung, O.J.Ganor, M.Krogh and A.Yu.Mikhailov, Nucl.Phys., **B564** (2000) 259
- [202] N.Nekrasov and A.Schwarz, Commun.Math.Phys., **198** (1998) 689

- [203] H.W.Braden and N.A.Nekrasov, hep-th/991201
- [204] P.Aspinwall, B.Greene and D.Morrison, Nucl.Phys., **B416** (1993) 414
- [205] N.C.Leung and C.Vafa, Adv.Theor.Math.Phys., **2** (1998) 91
- [206] D.-E.Diaconescu, Nucl.Phys., **B503** (1997) 220-238
- [207] P.M.Sutcliffe, Phys.Lett., **B381** (1996) 129
- [208] W.Nahm, in: *Monopoles in quantum field theories*, World Scientific, edt. Cragie et al. 440, 1982
- [209] I.M.Krichever and D.H.Phong, J.Diff.Geom., **45** (1997) 349-389
- [210] A.Sen, Nucl.Phys., **B475** (1996) 562-578
- [211] A. Gurevich and L. Pitaevskii, JETP, **65** (1973) 65
- [212] M.Douglas, M.Li, hep-th/9604041
- [213] M.Atiyah and N.Hitchin, *The geometry and dynamics of magnetic monopoles*, Princeton University Press, Princeton, NJ, 1988
- [214] A.Sen, Phys.Rev., **D55** (1997) 2501-2509